



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 06:33 AM GMT

PDB ID : 2XH0  
Title : ENGINEERING THE ENOLASE ACTIVE SITE POCKET: CRYSTAL  
STRUCTURE OF THE S39N Q167K D321R MUTANT OF YEAST ENO-  
LASE 1  
Authors : Schreier, B.; Hocker, B.  
Deposited on : 2010-06-08  
Resolution : 1.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3190 (1.70-1.70)
Clashscore	102246	3585 (1.70-1.70)
Ramachandran outliers	100387	3527 (1.70-1.70)
Sidechain outliers	100360	3527 (1.70-1.70)
RSRZ outliers	91569	3200 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	443	<div> <div>2%</div> <div>87%</div> <div>12%</div> <div>.</div> </div>
1	B	443	<div> <div>2%</div> <div>87%</div> <div>12%</div> <div>.</div> </div>
1	C	443	<div> <div>88%</div> <div>10%</div> <div>.</div> </div>
1	D	443	<div> <div>2%</div> <div>88%</div> <div>10%</div> <div>.</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	MG	C	1439	-	-	-	X
2	MG	D	1439	-	-	-	X

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 14787 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ENOLASE 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	438	Total	C	N	O	S	0	4	1
			3310	2092	578	636	4			
1	B	438	Total	C	N	O	S	0	2	1
			3272	2067	569	630	6			
1	C	438	Total	C	N	O	S	0	3	1
			3305	2089	577	635	4			
1	D	438	Total	C	N	O	S	0	2	1
			3287	2077	573	631	6			

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	437	LEU	-	EXPRESSION TAG	UNP P00924
A	438	HIS	-	EXPRESSION TAG	UNP P00924
A	439	HIS	-	EXPRESSION TAG	UNP P00924
A	440	HIS	-	EXPRESSION TAG	UNP P00924
A	441	HIS	-	EXPRESSION TAG	UNP P00924
A	442	HIS	-	EXPRESSION TAG	UNP P00924
A	443	HIS	-	EXPRESSION TAG	UNP P00924
A	39	ASN	SER	ENGINEERED MUTATION	UNP P00924
A	167	LYS	GLN	ENGINEERED MUTATION	UNP P00924
A	241	ILE	VAL	CONFLICT	UNP P00924
A	321	ARG	ASP	ENGINEERED MUTATION	UNP P00924
B	437	LEU	-	EXPRESSION TAG	UNP P00924
B	438	HIS	-	EXPRESSION TAG	UNP P00924
B	439	HIS	-	EXPRESSION TAG	UNP P00924
B	440	HIS	-	EXPRESSION TAG	UNP P00924
B	441	HIS	-	EXPRESSION TAG	UNP P00924
B	442	HIS	-	EXPRESSION TAG	UNP P00924
B	443	HIS	-	EXPRESSION TAG	UNP P00924
B	39	ASN	SER	ENGINEERED MUTATION	UNP P00924
B	167	LYS	GLN	ENGINEERED MUTATION	UNP P00924
B	241	ILE	VAL	CONFLICT	UNP P00924

*Continued on next page...*


*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	321	ARG	ASP	ENGINEERED MUTATION	UNP P00924
C	437	LEU	-	EXPRESSION TAG	UNP P00924
C	438	HIS	-	EXPRESSION TAG	UNP P00924
C	439	HIS	-	EXPRESSION TAG	UNP P00924
C	440	HIS	-	EXPRESSION TAG	UNP P00924
C	441	HIS	-	EXPRESSION TAG	UNP P00924
C	442	HIS	-	EXPRESSION TAG	UNP P00924
C	443	HIS	-	EXPRESSION TAG	UNP P00924
C	39	ASN	SER	ENGINEERED MUTATION	UNP P00924
C	167	LYS	GLN	ENGINEERED MUTATION	UNP P00924
C	241	ILE	VAL	CONFLICT	UNP P00924
C	321	ARG	ASP	ENGINEERED MUTATION	UNP P00924
D	437	LEU	-	EXPRESSION TAG	UNP P00924
D	438	HIS	-	EXPRESSION TAG	UNP P00924
D	439	HIS	-	EXPRESSION TAG	UNP P00924
D	440	HIS	-	EXPRESSION TAG	UNP P00924
D	441	HIS	-	EXPRESSION TAG	UNP P00924
D	442	HIS	-	EXPRESSION TAG	UNP P00924
D	443	HIS	-	EXPRESSION TAG	UNP P00924
D	39	ASN	SER	ENGINEERED MUTATION	UNP P00924
D	167	LYS	GLN	ENGINEERED MUTATION	UNP P00924
D	241	ILE	VAL	CONFLICT	UNP P00924
D	321	ARG	ASP	ENGINEERED MUTATION	UNP P00924

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Mg 1 1	0	0
2	A	1	Total Mg 1 1	0	0
2	D	1	Total Mg 1 1	0	0
2	C	1	Total Mg 1 1	0	0

- Molecule 3 is PHOSPHOENOLPYRUVATE (three-letter code: PEP) (formula: C<sub>3</sub>H<sub>5</sub>O<sub>6</sub>P).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	O	P	0	0
			10	3	6	1		
3	B	1	Total	C	O	P	0	0
			10	3	6	1		
3	C	1	Total	C	O	P	0	0
			10	3	6	1		
3	D	1	Total	C	O	P	0	0
			10	3	6	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	449	Total	O	0	0
			449	449		
4	B	411	Total	O	0	0
			411	411		
4	C	405	Total	O	0	0
			405	405		
4	D	304	Total	O	0	0
			304	304		







## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	55.33Å 60.80Å 120.67Å 89.89° 89.90° 65.84°	Depositor
Resolution (Å)	37.85 – 1.70 37.84 – 1.70	Depositor EDS
% Data completeness (in resolution range)	93.2 (37.85-1.70) 76.1 (37.84-1.70)	Depositor EDS
$R_{merge}$	0.16	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.72 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.197 , 0.246 0.198 , 0.247	Depositor DCC
$R_{free}$ test set	6059 reflections (5.31%)	DCC
Wilson B-factor (Å <sup>2</sup> )	12.6	Xtriage
Anisotropy	0.157	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 42.2	EDS
Estimated twinning fraction	0.147 for h,h-k,-l 0.459 for -h,-k,l 0.147 for -h,-h+k,-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.44$ , $\langle L^2 \rangle = 0.27$	Xtriage
Outliers	0 of 120183 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	14787	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.82% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, PEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.53	0/3382	0.66	2/4579 (0.0%)
1	B	0.52	0/3338	0.65	0/4528
1	C	0.53	0/3374	0.65	1/4568 (0.0%)
1	D	0.52	0/3353	0.65	1/4544 (0.0%)
All	All	0.52	0/13447	0.66	4/18219 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	119	ARG	NE-CZ-NH2	-5.33	117.63	120.30
1	D	12	ASP	CB-CG-OD1	5.14	122.93	118.30
1	A	119	ARG	NE-CZ-NH1	5.12	122.86	120.30
1	C	119	ARG	NE-CZ-NH2	-5.03	117.78	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3310	0	3323	34	0
1	B	3272	0	3238	37	0
1	C	3305	0	3317	36	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	3287	0	3275	27	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	10	0	2	0	0
3	B	10	0	2	0	0
3	C	10	0	2	0	0
3	D	10	0	2	0	0
4	A	449	0	0	5	0
4	B	411	0	0	10	1
4	C	405	0	0	10	0
4	D	304	0	0	3	0
All	All	14787	0	13161	126	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 126 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:369:VAL:O	1:C:392:THR:CG2	2.10	0.98
1:C:369:VAL:O	1:C:392:THR:HG23	1.69	0.90
1:C:300:GLU:HG2	1:C:321:ARG:HB3	1.57	0.86
1:B:194:LYS:HE3	4:B:2239:HOH:O	1.78	0.81
1:A:131:LYS:HE2	4:A:2227:HOH:O	1.86	0.74

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:2124:HOH:O	4:B:2263:HOH:O[1_565]	1.93	0.27

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	440/443 (99%)	432 (98%)	7 (2%)	1 (0%)	52	32
1	B	438/443 (99%)	429 (98%)	8 (2%)	1 (0%)	52	32
1	C	439/443 (99%)	431 (98%)	7 (2%)	1 (0%)	52	32
1	D	438/443 (99%)	427 (98%)	10 (2%)	1 (0%)	52	32
All	All	1755/1772 (99%)	1719 (98%)	32 (2%)	4 (0%)	52	32

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	402	ARG
1	B	402	ARG
1	C	402	ARG
1	D	402	ARG

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	345/351 (98%)	341 (99%)	4 (1%)	78	65
1	B	335/351 (95%)	331 (99%)	4 (1%)	78	65
1	C	344/351 (98%)	340 (99%)	4 (1%)	78	65
1	D	339/351 (97%)	336 (99%)	3 (1%)	84	76
All	All	1363/1404 (97%)	1348 (99%)	15 (1%)	80	69

5 of 15 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	373	HIS
1	B	392	THR
1	D	373	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	347	ASN
1	C	437	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	155	ASN
1	C	192	ASN
1	D	316	GLN
1	C	152	ASN
1	D	277	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	PEP	A	1440	2	5,9,9	1.37	1 (20%)	8,13,13	1.51	2 (25%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	PEP	B	1440	2	5,9,9	1.53	1 (20%)	8,13,13	1.30	1 (12%)
3	PEP	C	1440	2	5,9,9	1.44	1 (20%)	8,13,13	1.74	4 (50%)
3	PEP	D	1440	2	5,9,9	1.40	1 (20%)	8,13,13	1.16	1 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PEP	A	1440	2	-	0/5/9/9	0/0/0/0
3	PEP	B	1440	2	-	0/5/9/9	0/0/0/0
3	PEP	C	1440	2	-	0/5/9/9	0/0/0/0
3	PEP	D	1440	2	-	0/5/9/9	0/0/0/0

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1440	PEP	C3-C2	2.33	1.37	1.33
3	C	1440	PEP	C3-C2	2.46	1.37	1.33
3	D	1440	PEP	C3-C2	2.52	1.38	1.33
3	B	1440	PEP	C3-C2	2.73	1.38	1.33

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1440	PEP	O2-C2-C3	-3.18	118.30	124.73
3	C	1440	PEP	O2-C2-C3	-2.97	118.73	124.73
3	B	1440	PEP	O2-C2-C3	-2.39	119.89	124.73
3	D	1440	PEP	O2-C2-C3	-2.22	120.23	124.73
3	C	1440	PEP	O3P-P-O2	-2.18	98.14	105.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	438/443 (98%)	0.08	3 (0%) 89 91	6, 13, 21, 32	0
1	B	438/443 (98%)	0.22	11 (2%) 61 65	6, 14, 26, 44	0
1	C	438/443 (98%)	0.04	2 (0%) 91 93	6, 13, 21, 31	0
1	D	438/443 (98%)	0.17	11 (2%) 61 65	6, 13, 24, 45	0
All	All	1752/1772 (98%)	0.13	27 (1%) 76 80	6, 13, 23, 45	0

The worst 5 of 27 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	437	LEU	4.4
1	D	265	PRO	3.7
1	D	266	ASN	3.7
1	B	159	HIS	3.7
1	C	1	ALA	3.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors



of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	MG	C	1439	1/1	1.00	0.16	5.44	2,2,2,2	0
2	MG	D	1439	1/1	0.98	0.14	2.49	2,2,2,2	0
2	MG	A	1439	1/1	0.99	0.14	1.92	2,2,2,2	0
2	MG	B	1439	1/1	0.99	0.14	1.73	2,2,2,2	0
3	PEP	B	1440	10/10	0.97	0.10	-0.23	10,14,17,17	0
3	PEP	D	1440	10/10	0.97	0.10	-0.27	12,18,22,23	0
3	PEP	A	1440	10/10	0.97	0.06	-1.46	10,13,15,16	0
3	PEP	C	1440	10/10	0.98	0.08	-1.48	8,12,15,17	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.