



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 11:28 PM GMT

PDB ID : 1XIS  
Title : A METAL-MEDIATED HYDRIDE SHIFT MECHANISM FOR XYLOSE ISOMERASE BASED ON THE 1.6 ANGSTROMS STREPTOMYCES RUBIGINOSUS STRUCTURES WITH XYLITOL AND D-XYLOSE  
Authors : Whitlow, M.; Howard, A.J.  
Deposited on : 1991-03-25  
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	2732 (1.60-1.60)
Ramachandran outliers	100387	2654 (1.60-1.60)
Sidechain outliers	100360	2653 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	387	 81% 15% .

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3506 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called XYLOSE ISOMERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	387	Total	C	N	O	S	0	16	1
			3097	1937	575	576	9			

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Mn	0	0
			2	2		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	400	Total	O	0	7
			407	407		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

#### • Molecule 1: XYLOSE ISOMERASE



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	94.64Å 99.97Å 103.97Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 1.60	Depositor
% Data completeness (in resolution range)	(Not available) (10.00-1.60)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	PROLSQ	Depositor
R, $R_{free}$	0.141 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3506	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	16.0	wwPDB-VP

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.29	8/3258 (0.2%)	1.82	95/4403 (2.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	132[A]	GLU	CG-CD	-7.47	1.40	1.51
1	A	132[B]	GLU	CG-CD	-7.47	1.40	1.51
1	A	217	GLU	CD-OE1	-7.18	1.17	1.25
1	A	132[A]	GLU	CD-OE2	6.53	1.32	1.25
1	A	132[B]	GLU	CD-OE2	6.53	1.32	1.25
1	A	113	ARG	CZ-NH2	5.72	1.40	1.33
1	A	340[A]	ARG	CD-NE	-5.49	1.37	1.46
1	A	340[B]	ARG	CD-NE	-5.49	1.37	1.46

All (95) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	113	ARG	NE-CZ-NH1	18.39	129.50	120.30
1	A	132[A]	GLU	CB-CG-CD	15.36	155.66	114.20
1	A	132[B]	GLU	CB-CG-CD	15.36	155.66	114.20
1	A	331	ARG	NE-CZ-NH1	14.95	127.77	120.30
1	A	113	ARG	NE-CZ-NH2	-13.72	113.44	120.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	331	ARG	NE-CZ-NH2	-12.87	113.86	120.30
1	A	266[A]	ARG	NE-CZ-NH1	12.57	126.59	120.30
1	A	266[B]	ARG	NE-CZ-NH1	12.57	126.59	120.30
1	A	157	ARG	NE-CZ-NH2	10.95	125.78	120.30
1	A	381	ASP	CB-CG-OD1	10.71	127.94	118.30
1	A	340[A]	ARG	CD-NE-CZ	10.64	138.49	123.60
1	A	340[B]	ARG	CD-NE-CZ	10.64	138.49	123.60
1	A	74	ARG	NE-CZ-NH1	10.46	125.53	120.30
1	A	266[A]	ARG	NE-CZ-NH2	-10.41	115.10	120.30
1	A	266[B]	ARG	NE-CZ-NH2	-10.41	115.10	120.30
1	A	340[A]	ARG	NE-CZ-NH2	-10.26	115.17	120.30
1	A	340[B]	ARG	NE-CZ-NH2	-10.26	115.17	120.30
1	A	368	ARG	NE-CZ-NH1	10.09	125.34	120.30
1	A	188	ARG	NE-CZ-NH2	-9.52	115.54	120.30
1	A	321	ARG	NE-CZ-NH1	9.43	125.01	120.30
1	A	104	PHE	CB-CG-CD2	-9.39	114.22	120.80
1	A	23[A]	ARG	CG-CD-NE	9.04	130.79	111.80
1	A	23[B]	ARG	CG-CD-NE	9.04	130.79	111.80
1	A	74	ARG	NE-CZ-NH2	-9.02	115.79	120.30
1	A	74	ARG	CD-NE-CZ	9.01	136.21	123.60
1	A	124	ASP	CB-CG-OD2	-8.98	110.22	118.30
1	A	121	ARG	NE-CZ-NH2	-8.64	115.98	120.30
1	A	334	ARG	NE-CZ-NH1	8.54	124.57	120.30
1	A	10	ARG	NE-CZ-NH1	8.30	124.45	120.30
1	A	140	ARG	NE-CZ-NH2	-8.17	116.22	120.30
1	A	208	ARG	NE-CZ-NH2	-7.99	116.31	120.30
1	A	31	ARG	NE-CZ-NH1	7.97	124.28	120.30
1	A	23[A]	ARG	CD-NE-CZ	7.86	134.60	123.60
1	A	23[B]	ARG	CD-NE-CZ	7.86	134.60	123.60
1	A	242	PHE	CB-CG-CD2	-7.57	115.50	120.80
1	A	295	ASP	CB-CG-OD1	7.51	125.06	118.30
1	A	76[A]	ARG	NE-CZ-NH1	7.47	124.04	120.30
1	A	76[B]	ARG	NE-CZ-NH1	7.47	124.04	120.30
1	A	284	ARG	NE-CZ-NH1	7.21	123.90	120.30
1	A	109	ARG	NE-CZ-NH2	-7.16	116.72	120.30
1	A	340[A]	ARG	NE-CZ-NH1	7.06	123.83	120.30
1	A	340[B]	ARG	NE-CZ-NH1	7.06	123.83	120.30
1	A	323	ASP	CB-CG-OD2	7.04	124.64	118.30
1	A	284	ARG	NE-CZ-NH2	-6.99	116.80	120.30
1	A	177	ARG	NE-CZ-NH1	6.99	123.80	120.30
1	A	152	ARG	NE-CZ-NH1	6.97	123.79	120.30
1	A	41	ARG	NE-CZ-NH1	6.94	123.77	120.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	205[A]	ARG	NE-CZ-NH1	6.89	123.75	120.30
1	A	205[B]	ARG	NE-CZ-NH1	6.89	123.75	120.30
1	A	266[A]	ARG	CD-NE-CZ	6.89	133.24	123.60
1	A	266[B]	ARG	CD-NE-CZ	6.89	133.24	123.60
1	A	76[A]	ARG	NE-CZ-NH2	-6.84	116.88	120.30
1	A	76[B]	ARG	NE-CZ-NH2	-6.84	116.88	120.30
1	A	205[A]	ARG	CD-NE-CZ	6.84	133.17	123.60
1	A	205[B]	ARG	CD-NE-CZ	6.84	133.17	123.60
1	A	387[A]	ARG	NE-CZ-NH1	6.69	123.64	120.30
1	A	387[B]	ARG	NE-CZ-NH1	6.69	123.64	120.30
1	A	336	ASP	CB-CG-OD2	6.49	124.14	118.30
1	A	347	LEU	CB-CG-CD2	6.48	122.02	111.00
1	A	320	PHE	CB-CG-CD2	-6.37	116.34	120.80
1	A	152	ARG	NE-CZ-NH2	-6.36	117.12	120.30
1	A	212	TYR	CB-CG-CD1	-6.32	117.21	121.00
1	A	287	ASP	CB-CG-OD1	6.32	123.98	118.30
1	A	124	ASP	CB-CG-OD1	6.24	123.92	118.30
1	A	132[A]	GLU	OE1-CD-OE2	-6.24	115.81	123.30
1	A	132[B]	GLU	OE1-CD-OE2	-6.24	115.81	123.30
1	A	212	TYR	CB-CG-CD2	6.20	124.72	121.00
1	A	42	ARG	NE-CZ-NH1	6.06	123.33	120.30
1	A	321	ARG	NE-CZ-NH2	-6.03	117.29	120.30
1	A	104	PHE	CB-CG-CD1	5.94	124.96	120.80
1	A	242	PHE	CB-CG-CD1	5.93	124.95	120.80
1	A	24	ASP	CB-CA-C	5.91	122.21	110.40
1	A	245	ASP	CB-CG-OD1	5.89	123.61	118.30
1	A	121	ARG	NE-CZ-NH1	5.89	123.24	120.30
1	A	32	ARG	NE-CZ-NH1	5.85	123.22	120.30
1	A	337	GLU	CG-CD-OE1	5.81	129.93	118.30
1	A	10	ARG	NE-CZ-NH2	-5.72	117.44	120.30
1	A	109	ARG	NE-CZ-NH1	5.67	123.13	120.30
1	A	168	TYR	CB-CG-CD1	-5.59	117.64	121.00
1	A	132[A]	GLU	CG-CD-OE1	5.50	129.30	118.30
1	A	132[B]	GLU	CG-CD-OE1	5.50	129.30	118.30
1	A	168	TYR	CB-CG-CD2	5.41	124.25	121.00
1	A	345	ASP	CB-CG-OD1	5.40	123.16	118.30
1	A	65	ASP	CB-CG-OD1	5.39	123.15	118.30
1	A	269	PHE	CB-CG-CD1	-5.36	117.05	120.80
1	A	99	PHE	CB-CG-CD2	-5.31	117.08	120.80
1	A	174	TYR	CB-CG-CD1	5.28	124.17	121.00
1	A	42	ARG	NE-CZ-NH2	-5.22	117.69	120.30
1	A	264	ASP	CB-CG-OD1	5.21	122.99	118.30

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	205[A]	ARG	NE-CZ-NH2	-5.20	117.70	120.30
1	A	205[B]	ARG	NE-CZ-NH2	-5.20	117.70	120.30
1	A	292	ARG	NE-CZ-NH2	5.15	122.88	120.30
1	A	117	ARG	NE-CZ-NH1	5.11	122.86	120.30
1	A	210	GLU	CG-CD-OE1	5.05	128.40	118.30
1	A	286	PHE	CB-CG-CD1	-5.04	117.27	120.80

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	208	ARG	Sidechain
1	A	331	ARG	Sidechain
1	A	41	ARG	Sidechain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3097	0	2947	22	0
2	A	2	0	0	0	0
3	A	407	0	0	9	0
All	All	3506	0	2947	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:96:HIS:HD2	1:A:98:VAL:H	1.30	0.78
1:A:92:ASN:HD21	1:A:95:THR:H	1.36	0.72
1:A:93:LEU:HD21	3:A:618:HOH:O	1.97	0.63
1:A:24:ASP:HB2	1:A:25:PRO:HD2	1.79	0.63
1:A:248:GLY:HA2	3:A:608:HOH:O	2.00	0.62

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:219:GLY:H	1:A:247:ASN:HD21	1.51	0.59
1:A:218:VAL:HG11	3:A:608:HOH:O	2.02	0.58
1:A:96:HIS:CD2	1:A:98:VAL:H	2.18	0.55
1:A:92:ASN:C	1:A:92:ASN:HD22	2.12	0.52
1:A:92:ASN:ND2	1:A:94:PHE:H	2.10	0.50
1:A:91:THR:HG21	3:A:618:HOH:O	2.14	0.48
1:A:337:GLU:HG2	1:A:340[A]:ARG:NH2	2.30	0.46
1:A:92:ASN:HD22	1:A:94:PHE:H	1.65	0.45
1:A:41:ARG:NH2	3:A:564:HOH:O	2.48	0.45
1:A:262:ALA:HA	3:A:801:HOH:O	2.16	0.45
1:A:119:THR:N	3:A:618:HOH:O	2.49	0.44
1:A:218:VAL:HB	1:A:247:ASN:ND2	2.33	0.44
1:A:92:ASN:ND2	1:A:95:THR:H	2.08	0.43
1:A:266[A]:ARG:HG3	3:A:454:HOH:O	2.17	0.43
1:A:24:ASP:CB	1:A:25:PRO:HD2	2.49	0.41
1:A:132[A]:GLU:HG3	1:A:133:THR:N	2.35	0.41
1:A:118:LYS:HB3	3:A:618:HOH:O	2.21	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	401/387 (104%)	388 (97%)	12 (3%)	1 (0%)	52 28

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	186	GLU

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	316/303 (104%)	308 (98%)	8 (2%)	55 26

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	ARG
1	A	23[A]	ARG
1	A	23[B]	ARG
1	A	92	ASN
1	A	185	ASN
1	A	230	HIS
1	A	247	ASN
1	A	347	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	77	GLN
1	A	92	ASN
1	A	96	HIS
1	A	215	ASN
1	A	230	HIS
1	A	247	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

### 6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

### 6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

### 6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.