



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2016 – 08:21 PM GMT

PDB ID : 4XJ0  
Title : Crystal structure of ERK2 in complex with an inhibitor 14K  
Authors : Yin, J.; Wang, W.  
Deposited on : 2015-01-08  
Resolution : 2.58 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026982  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20026982


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.58 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2636 (2.60-2.56)
Clashscore	102246	3003 (2.60-2.56)
Ramachandran outliers	100387	2956 (2.60-2.56)
Sidechain outliers	100360	2956 (2.60-2.56)
RSRZ outliers	91569	2642 (2.60-2.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	349	<div> <div>7%</div> <div>86%</div> <div>13%</div> <div>..</div> </div>
1	B	349	<div> <div>16%</div> <div>86%</div> <div>11%</div> <div>.</div> </div>

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 5788 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Mitogen-activated protein kinase 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	346	Total	C	N	O	P	S	0	0	0
			2831	1815	483	518	1	14			
1	B	339	Total	C	N	O	P	S	0	0	0
			2775	1778	474	508	1	14			

- Molecule 2 is 1-[(1S)-1-(4-chloro-3-fluorophenyl)-2-hydroxyethyl]-4-[2-(tetrahydro-2H-pyran-4-ylamino)pyrimidin-4-yl]pyridin-2(1H)-one (three-letter code: 41B) (formula: C<sub>22</sub>H<sub>22</sub>ClFN<sub>4</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	Cl	F	N	O	0	0
			31	22	1	1	4	3		
2	B	1	Total	C	Cl	F	N	O	0	0
			31	22	1	1	4	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	78	Total 78	O 78	0	0
3	B	42	Total 42	O 42	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Mitogen-activated protein kinase 1



#### • Molecule 1: Mitogen-activated protein kinase 1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	83.19 Å 83.19 Å 274.49 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.82 – 2.58 44.66 – 2.58	Depositor EDS
% Data completeness (in resolution range)	99.3 (45.82-2.58) 99.3 (44.66-2.58)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.54 (at 2.58 Å)	Xtriage
Refinement program	BUSTER 2.11.4	Depositor
R, $R_{free}$	0.208 , 0.244 0.205 , 0.239	Depositor DCC
$R_{free}$ test set	1538 reflections (5.19%)	DCC
Wilson B-factor (Å <sup>2</sup> )	59.1	Xtriage
Anisotropy	0.525	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 59.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtriage
Outliers	1 of 31184 reflections (0.003%)	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5788	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	82.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.36% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NEP, 41B

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.48	0/2882	0.65	0/3900
1	B	0.46	0/2826	0.63	0/3825
All	All	0.47	0/5708	0.64	0/7725

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2831	0	2816	22	0
1	B	2775	0	2757	18	0
2	A	31	0	22	1	0
2	B	31	0	22	2	0
3	A	78	0	0	0	0
3	B	42	0	0	0	0
All	All	5788	0	5617	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:80:HIS:HB3	1:B:83:ILE:HG12	1.80	0.62
1:A:83:ILE:HD12	1:A:140:ILE:HD11	1.81	0.61
1:B:148:ARG:HG2	1:B:209:ILE:HD11	1.82	0.61
1:A:80:HIS:HB3	1:A:83:ILE:HG12	1.82	0.60
1:B:149:ASP:HB2	1:B:170:LEU:HD12	1.85	0.59
1:B:54:LYS:HD3	2:B:401:41B:H1	1.84	0.59
1:B:53:ILE:HG12	1:B:104:VAL:HG22	1.85	0.59
1:A:148:ARG:HG2	1:A:209:ILE:HD11	1.86	0.56
1:A:71:GLU:HG3	1:A:168:PHE:H	1.70	0.56
1:A:149:ASP:HB2	1:A:170:LEU:HD12	1.88	0.55
1:A:106:ASP:O	2:A:401:41B:H5	2.08	0.54
1:B:130:LEU:HD22	1:B:218:LEU:HD13	1.88	0.54
1:A:91:ARG:HG3	1:A:98:MET:SD	2.50	0.52
1:B:148:ARG:HD3	1:B:170:LEU:O	2.09	0.51
1:A:83:ILE:CD1	1:A:140:ILE:HD11	2.42	0.50
1:A:90:ILE:HB	1:A:102:TYR:HB2	1.94	0.50
1:A:148:ARG:HD3	1:A:170:LEU:O	2.12	0.50
1:A:257:ASN:HB3	1:A:260:ALA:HB2	1.94	0.50
1:A:66:GLN:O	1:A:70:ARG:HB2	2.13	0.49
1:B:83:ILE:HD12	1:B:140:ILE:HD11	1.94	0.49
1:B:345:GLU:O	1:B:349:GLU:HG2	2.14	0.48
1:A:89:ILE:HD13	1:A:351:THR:HG22	1.94	0.48
1:A:201:ASN:HB3	1:A:256:ILE:H	1.81	0.46
1:B:69:LEU:HD23	1:B:335:LEU:HD21	1.98	0.46
1:B:275:TRP:HB3	1:B:287:LEU:HD22	1.97	0.46
1:B:89:ILE:HD13	1:B:351:THR:HG22	1.99	0.44
1:A:13:MET:HB3	1:A:18:VAL:HA	1.99	0.44
1:A:257:ASN:HB3	1:A:260:ALA:CB	2.47	0.44
1:A:54:LYS:HD2	1:A:56:ILE:HD11	2.00	0.43
1:A:39:VAL:HG22	1:A:54:LYS:HG2	2.00	0.43
1:B:183:PHE:CZ	1:B:259:LYS:HD3	2.54	0.42
1:B:106:ASP:O	2:B:401:41B:H5	2.19	0.42
1:B:335:LEU:HA	1:B:338:LEU:HD12	2.01	0.42
1:A:330:LYS:HD2	1:A:330:LYS:HA	1.93	0.42
1:A:288:ASP:O	1:A:292:LYS:HG2	2.20	0.41
1:A:267:LEU:HD23	1:A:268:PRO:HD2	2.02	0.41
1:B:262:ASN:HD21	1:B:265:LEU:HG	1.85	0.41
1:B:200:LEU:O	1:B:259:LYS:HB2	2.20	0.40
1:A:91:ARG:NH2	1:A:355:GLN:HG3	2.36	0.40
1:B:136:GLY:O	1:B:140:ILE:HG12	2.20	0.40



There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	339/349 (97%)	320 (94%)	15 (4%)	4 (1%)	16	33
1	B	334/349 (96%)	305 (91%)	25 (8%)	4 (1%)	16	33
All	All	673/698 (96%)	625 (93%)	40 (6%)	8 (1%)	16	33

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	248	SER
1	A	181	THR
1	A	188	VAL
1	B	181	THR
1	A	15	ARG
1	B	15	ARG
1	A	180	HIS
1	B	46	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	311/315 (99%)	299 (96%)	12 (4%)	39	66
1	B	304/315 (96%)	295 (97%)	9 (3%)	48	75

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	615/630 (98%)	594 (97%)	21 (3%)	44 71

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	17	GLN
1	A	19	PHE
1	A	91	ARG
1	A	184	LEU
1	A	201	ASN
1	A	202	SER
1	A	227	ILE
1	A	233	TYR
1	A	252	LEU
1	A	263	TYR
1	A	278	LEU
1	A	337	ASP
1	B	15	ARG
1	B	19	PHE
1	B	91	ARG
1	B	179	ASP
1	B	233	TYR
1	B	251	ASP
1	B	263	TYR
1	B	278	LEU
1	B	336	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link

column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	NEP	A	125	1	8,14,15	4.44	3 (37%)	5,20,22	2.24	1 (20%)
1	NEP	B	125	1	8,14,15	2.45	2 (25%)	5,20,22	2.21	2 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	NEP	A	125	1	-	0/4/12/14	0/1/1/1
1	NEP	B	125	1	-	0/4/12/14	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	125	NEP	P-O1P	-5.63	1.42	1.54
1	B	125	NEP	P-O3P	-5.28	1.42	1.47
1	A	125	NEP	CD2-NE2	-4.37	1.31	1.39
1	B	125	NEP	CD2-NE2	-3.87	1.32	1.39
1	A	125	NEP	P-O3P	10.13	1.56	1.47

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	125	NEP	CG-CD2-NE2	-4.25	103.27	108.96
1	A	125	NEP	CG-CD2-NE2	-4.19	103.35	108.96
1	B	125	NEP	O-C-CA	-2.01	120.34	125.72

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	41B	A	401	-	31,34,34	1.13	1 (3%)	40,47,47	2.51	11 (27%)
2	41B	B	401	-	31,34,34	1.09	1 (3%)	40,47,47	2.37	10 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	41B	A	401	-	-	0/16/26/26	0/4/4/4
2	41B	B	401	-	-	0/16/26/26	0/4/4/4

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	41B	C22-N24	4.40	1.39	1.34
2	A	401	41B	C22-N24	4.51	1.40	1.34

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	41B	N21-C22-N23	-6.83	119.38	126.66
2	B	401	41B	N21-C22-N23	-6.27	119.98	126.66
2	B	401	41B	C19-C20-N21	-3.29	120.03	123.88
2	A	401	41B	C13-C14-C18	-2.56	117.08	121.30
2	A	401	41B	C19-C20-N21	-2.52	120.93	123.88

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401	41B	C19-C18-N23	-2.38	118.54	121.98
2	A	401	41B	C4-C3-CL1	-2.27	117.90	120.03
2	A	401	41B	C19-C18-N23	-2.11	118.93	121.98
2	B	401	41B	C29-O28-C27	2.03	116.81	109.89
2	A	401	41B	C29-O28-C27	2.12	117.13	109.89
2	B	401	41B	C15-C14-C18	2.13	123.26	120.57
2	A	401	41B	C15-C14-C18	2.53	123.77	120.57
2	B	401	41B	C2-C3-C4	2.62	120.93	119.07
2	B	401	41B	C14-C18-N23	2.63	119.72	116.00
2	A	401	41B	N24-C22-N23	2.76	120.92	116.97
2	B	401	41B	N24-C22-N23	3.24	121.60	116.97
2	A	401	41B	C2-C3-C4	3.33	121.43	119.07
2	B	401	41B	C18-N23-C22	7.13	121.92	116.45
2	A	401	41B	C20-N21-C22	7.92	121.54	115.42
2	B	401	41B	C20-N21-C22	7.96	121.57	115.42
2	A	401	41B	C18-N23-C22	8.06	122.64	116.45

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	41B	1	0
2	B	401	41B	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	345/349 (98%)	0.45	26 (7%) 17 13	45, 70, 123, 169	0
1	B	338/349 (96%)	0.78	55 (16%) 2 1	48, 83, 144, 185	0
All	All	683/698 (97%)	0.61	81 (11%) 6 4	45, 77, 138, 185	0

All (81) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	21	VAL	6.1
1	B	333	MET	6.0
1	A	264	LEU	5.9
1	B	13	MET	5.7
1	B	352	ALA	4.7
1	B	180	HIS	4.3
1	A	256	ILE	4.3
1	B	92	ALA	4.3
1	B	93	PRO	4.3
1	B	342	LYS	4.3
1	B	94	THR	4.0
1	B	338	LEU	4.0
1	B	183	PHE	3.9
1	B	99	LYS	3.9
1	B	14	VAL	3.8
1	B	16	GLY	3.7
1	A	265	LEU	3.7
1	B	19	PHE	3.6
1	B	95	ILE	3.6
1	B	349	GLU	3.6
1	B	332	ASP	3.6
1	A	254	CYS	3.4
1	B	296	PHE	3.4
1	B	347	ILE	3.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	348	PHE	3.3
1	B	17	GLN	3.3
1	A	249	GLN	3.2
1	A	180	HIS	3.2
1	B	25	TYR	3.1
1	B	28	LEU	3.2
1	B	203	LYS	3.1
1	B	101	VAL	3.1
1	A	252	LEU	3.1
1	A	359	ARG	3.1
1	B	49	VAL	3.1
1	B	181	THR	3.1
1	B	12	GLU	3.0
1	B	345	GLU	3.0
1	B	102	TYR	3.0
1	B	264	LEU	3.0
1	B	96	GLU	2.9
1	B	356	PRO	2.9
1	B	59	PHE	2.9
1	A	267	LEU	2.9
1	B	329	PHE	2.8
1	B	97	GLN	2.8
1	A	179	ASP	2.8
1	A	13	MET	2.8
1	A	188	VAL	2.7
1	A	181	THR	2.7
1	B	247	PRO	2.6
1	A	329	PHE	2.6
1	B	90	ILE	2.5
1	B	98	MET	2.5
1	B	343	LEU	2.5
1	B	177	ASP	2.5
1	A	99	LYS	2.5
1	B	188	VAL	2.3
1	B	354	PHE	2.3
1	A	94	THR	2.3
1	B	48	LYS	2.2
1	B	357	GLY	2.2
1	A	262	ASN	2.2
1	B	178	HIS	2.2
1	B	179	ASP	2.2
1	B	69	LEU	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	335	LEU	2.2
1	B	89	ILE	2.1
1	B	351	THR	2.1
1	B	278	LEU	2.1
1	B	38	MET	2.1
1	B	88	ASP	2.1
1	A	59	PHE	2.1
1	A	95	ILE	2.1
1	A	255	ILE	2.1
1	A	268	PRO	2.1
1	A	336	ASP	2.0
1	A	260	ALA	2.0
1	A	16	GLY	2.0
1	A	204	GLY	2.0
1	A	278	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q < 0.9
1	NEP	B	125	14/15	0.97	0.18	-	53,62,65,66	0
1	NEP	A	125	14/15	0.97	0.20	-	55,59,66,66	0

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	41B	A	401	31/31	0.94	0.17	-0.20	55,66,98,104	0
2	41B	B	401	31/31	0.93	0.15	-0.73	59,74,94,101	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.