



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 11:30 PM GMT

PDB ID : 1XK3  
Title : NADPH- and Ascorbate-Supported Heme Oxygenase Reactions are Distinct.  
Regiospecificity of Heme Cleavage by the R183E Mutant  
Authors : Wang, J.; Lad, L.; Poulos, T.L.; Ortiz de montellano, P.R.  
Deposited on : 2004-09-26  
Resolution : 2.08 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.08 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	5101 (2.10-2.06)
Ramachandran outliers	100387	5048 (2.10-2.06)
Sidechain outliers	100360	5049 (2.10-2.06)
RSRZ outliers	91569	4556 (2.10-2.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	233	
1	B	233	



In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Heme oxygenase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	214	Total 1743	C 1120	N 296	O 322	S 5	0	0	0
1	B	214	Total 1743	C 1120	N 296	O 322	S 5	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	183	GLU	ARG	ENGINEERED	UNP P09601
B	183	GLU	ARG	ENGINEERED	UNP P09601

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is NITRIC OXIDE (three-letter code: NO) (formula: NO).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	N	O	0	0
			2	1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	150	Total	O	0	0
			150	150		
4	B	112	Total	O	0	0
			112	112		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Heme oxygenase 1



#### • Molecule 1: Heme oxygenase 1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	61.22Å 54.48Å 70.89Å 90.00° 98.75° 90.00°	Depositor
Resolution (Å)	50.00 – 2.08 30.25 – 2.09	Depositor EDS
% Data completeness (in resolution range)	(Not available) (50.00-2.08) 99.7 (30.25-2.09)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.37 (at 2.10Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, $R_{free}$	0.210 , 0.230 0.209 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	DCC
Wilson B-factor (Å <sup>2</sup> )	34.5	Xtriage
Anisotropy	0.563	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 58.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 27481 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3836	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.66% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, NO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.34	0/1784	0.54	0/2412
1	B	0.35	0/1784	0.56	0/2412
All	All	0.35	0/3568	0.55	0/4824

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1743	0	1731	59	0
1	B	1743	0	1731	41	0
2	A	43	0	30	1	0
2	B	43	0	30	3	0
3	A	2	0	0	0	0
4	A	150	0	0	8	0
4	B	112	0	0	6	0
All	All	3836	0	3522	96	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (96) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:113:ARG:HD2	1:A:212:GLN:HE22	1.22	1.02
1:A:113:ARG:HD2	1:A:212:GLN:NE2	1.82	0.93
1:B:185:ARG:HH11	1:B:185:ARG:HG2	1.32	0.92
1:A:179:LYS:O	1:A:183:GLU:HG3	1.76	0.86
1:A:157:LEU:HD23	1:A:157:LEU:H	1.43	0.80
1:A:71:SER:OG	1:A:73:VAL:HG22	1.89	0.73
1:B:76:PRO:O	1:B:185:ARG:HG2	1.88	0.73
1:A:211:ILE:O	1:A:215:GLU:HG3	1.89	0.72
1:B:36:ASN:HB3	1:B:41:GLN:HG3	1.74	0.69
1:A:91:GLN:HE22	1:B:85:ARG:HH22	1.41	0.69
1:A:177:LYS:HE2	1:A:180:GLN:OE1	1.94	0.68
1:A:85:ARG:HH22	1:B:91:GLN:HE22	1.42	0.67
1:A:47:PHE:O	1:A:50:VAL:HG12	1.96	0.66
1:B:185:ARG:NH1	1:B:185:ARG:HG2	2.04	0.65
1:B:10:PRO:HD2	1:B:16:ALA:HA	1.78	0.65
1:A:44:ARG:HG2	4:A:425:HOH:O	2.00	0.62
1:B:100:ARG:HD3	4:B:390:HOH:O	2.00	0.60
1:B:86:LYS:O	1:B:90:GLU:HG3	2.02	0.60
1:B:144:GLY:HA3	4:B:324:HOH:O	2.02	0.59
1:A:108:THR:O	1:A:112:GLN:HG3	2.03	0.58
1:A:145:GLN:CD	1:A:145:GLN:H	2.06	0.58
1:A:212:GLN:HG3	4:A:428:HOH:O	2.03	0.57
1:A:113:ARG:HH11	1:A:212:GLN:NE2	2.02	0.57
1:A:208:LEU:O	1:A:212:GLN:HG2	2.05	0.57
1:A:162:GLU:OE2	1:B:148:LYS:HE3	2.05	0.57
1:B:75:ALA:HB3	1:B:76:PRO:HD3	1.87	0.56
1:A:69:LYS:HG3	1:A:70:GLU:HG3	1.87	0.56
1:B:196:ARG:O	1:B:200:ILE:HG12	2.05	0.56
1:A:116:LYS:O	1:A:120:GLU:HG3	2.06	0.56
1:B:147:LEU:HD21	2:B:300:HEM:HBB2	1.87	0.56
1:B:71:SER:OG	1:B:73:VAL:HG22	2.06	0.55
1:B:55:TYR:O	1:B:59:VAL:HG23	2.06	0.55
1:A:125:GLU:N	1:A:126:PRO:HD3	2.22	0.54
1:A:55:TYR:O	1:A:59:VAL:HG23	2.06	0.54
1:A:145:GLN:H	1:A:145:GLN:NE2	2.06	0.53
1:B:84:HIS:HD2	4:B:356:HOH:O	1.90	0.53
1:A:34:MET:HE2	1:A:34:MET:HA	1.90	0.53
1:B:109:PRO:O	1:B:113:ARG:HG3	2.08	0.53
1:A:177:LYS:CE	1:A:177:LYS:HA	2.39	0.52
1:A:75:ALA:HB3	1:A:76:PRO:HD3	1.91	0.52

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:56:HIS:HD2	4:A:482:HOH:O	1.92	0.52
1:A:157:LEU:HB2	1:A:158:PRO:HD2	1.90	0.52
1:A:23:GLU:HB3	4:A:430:HOH:O	2.09	0.52
1:B:179:LYS:O	1:B:183:GLU:HG3	2.10	0.52
1:A:146:VAL:O	1:A:150:ILE:HG13	2.09	0.51
1:A:147:LEU:HD21	2:A:300:HEM:HBB2	1.93	0.51
1:B:96:TRP:CH2	1:B:157:LEU:HD21	2.46	0.51
1:B:56:HIS:HD2	4:B:379:HOH:O	1.92	0.50
1:B:18:LYS:NZ	2:B:300:HEM:CGD	2.74	0.50
1:A:84:HIS:HD2	4:A:490:HOH:O	1.94	0.49
1:B:17:LEU:HD12	1:B:186:MET:CE	2.42	0.49
1:A:153:LYS:HE3	4:A:546:HOH:O	2.11	0.49
1:A:123:ARG:NH1	4:A:518:HOH:O	2.45	0.49
1:A:65:ILE:HG23	1:A:74:PHE:CE2	2.48	0.48
1:B:146:VAL:HG23	4:B:394:HOH:O	2.12	0.48
1:A:177:LYS:O	1:A:180:GLN:HG2	2.13	0.48
1:A:10:PRO:HD2	1:A:16:ALA:HA	1.95	0.48
1:A:101:TRP:O	1:A:105:ILE:HG23	2.13	0.48
1:A:38:GLN:C	1:A:40:GLY:H	2.17	0.48
1:B:180:GLN:HB3	4:B:392:HOH:O	2.13	0.48
1:B:213:LEU:O	1:B:217:LEU:HG	2.13	0.48
1:A:174:SER:OG	1:A:177:LYS:HB2	2.14	0.47
1:B:23:GLU:HG2	1:B:24:VAL:N	2.29	0.47
1:A:149:LYS:HA	1:A:152:GLN:HE21	1.79	0.47
1:A:37:PHE:O	1:A:150:ILE:HG21	2.15	0.46
1:B:18:LYS:HZ1	2:B:300:HEM:CGD	2.29	0.46
1:B:23:GLU:N	1:B:23:GLU:OE1	2.49	0.46
1:B:163:GLY:O	1:B:164:LEU:HD23	2.17	0.45
1:B:192:THR:HB	1:B:193:PRO:HD2	1.99	0.45
1:B:116:LYS:HE2	1:B:120:GLU:OE2	2.17	0.45
1:A:58:TYR:O	1:A:62:GLU:HG3	2.17	0.45
1:A:177:LYS:HE2	1:A:180:GLN:CD	2.37	0.44
1:A:162:GLU:OE2	1:B:148:LYS:CE	2.65	0.44
1:A:86:LYS:HE3	4:A:488:HOH:O	2.17	0.44
1:B:58:TYR:CZ	1:B:136:ARG:HD3	2.52	0.44
1:A:157:LEU:CD2	1:A:157:LEU:H	2.22	0.44
1:B:53:SER:HB3	1:B:213:LEU:HD21	1.99	0.44
1:A:177:LYS:HE3	1:A:177:LYS:HA	2.00	0.44
1:B:186:MET:O	1:B:189:LEU:HB2	2.18	0.44
1:B:96:TRP:CH2	1:B:155:LEU:HD13	2.53	0.43
1:A:55:TYR:HB2	1:A:93:LEU:HD11	2.00	0.43

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:10:PRO:HD3	1:A:19:GLU:OE2	2.18	0.43
1:A:124:THR:C	1:A:126:PRO:HD3	2.40	0.42
1:B:44:ARG:O	1:B:48:LYS:HG3	2.19	0.42
1:A:55:TYR:HA	1:A:89:LEU:HD13	2.01	0.42
1:A:145:GLN:NE2	1:A:145:GLN:N	2.66	0.42
1:A:183:GLU:HA	1:A:186:MET:HE2	2.02	0.42
1:B:96:TRP:HH2	1:B:157:LEU:HD21	1.84	0.42
1:A:61:LEU:O	1:A:65:ILE:HG13	2.19	0.42
1:B:17:LEU:CD2	1:B:200:ILE:HD13	2.49	0.42
1:A:86:LYS:HE2	1:A:90:GLU:OE1	2.20	0.41
1:B:128:LEU:HD22	1:B:199:VAL:HA	2.02	0.41
1:A:47:PHE:CD2	1:A:155:LEU:HD21	2.56	0.41
1:A:197:GLN:O	1:A:201:GLU:HG2	2.20	0.41
1:A:36:ASN:HB3	1:A:41:GLN:O	2.21	0.40
1:A:46:GLY:HA2	1:A:221:LEU:HD11	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	212/233 (91%)	205 (97%)	7 (3%)	0	100	100
1	B	212/233 (91%)	206 (97%)	6 (3%)	0	100	100
All	All	424/466 (91%)	411 (97%)	13 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	184/202 (91%)	181 (98%)	3 (2%)	70	75
1	B	184/202 (91%)	181 (98%)	3 (2%)	70	75
All	All	368/404 (91%)	362 (98%)	6 (2%)	70	75

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	114	TYR
1	A	145	GLN
1	A	177	LYS
1	B	23	GLU
1	B	114	TYR
1	B	185	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	11	GLN
1	A	56	HIS
1	A	84	HIS
1	A	91	GLN
1	A	112	GLN
1	A	145	GLN
1	A	152	GLN
1	A	171	ASN
1	A	212	GLN
1	B	30	ASN
1	B	36	ASN
1	B	56	HIS
1	B	84	HIS
1	B	91	GLN
1	B	102	GLN
1	B	171	ASN
1	B	180	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	HEM	A	300	1,3	30,50,50	2.67	11 (36%)	24,82,82	2.09	8 (33%)
3	NO	A	400	2	0,1,1	0.00	-	0,0,0	0.00	-
2	HEM	B	300	1	30,50,50	2.78	12 (40%)	24,82,82	1.94	6 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	300	1,3	-	0/10/54/54	0/0/8/8
3	NO	A	400	2	-	0/0/0/0	0/0/0/0
2	HEM	B	300	1	-	0/10/54/54	0/0/8/8

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	300	HEM	C3B-CAB	-6.47	1.39	1.51
2	B	300	HEM	C3C-CAC	-6.27	1.39	1.51
2	B	300	HEM	C3B-CAB	-6.19	1.39	1.51
2	B	300	HEM	C2D-C3D	-6.18	1.36	1.54
2	A	300	HEM	C2D-C3D	-6.15	1.36	1.54
2	A	300	HEM	C3C-CAC	-6.01	1.40	1.51
2	B	300	HEM	C3B-C4B	-5.02	1.47	1.51
2	B	300	HEM	C3D-C4D	-4.24	1.46	1.51
2	A	300	HEM	C3B-C4B	-3.74	1.48	1.51
2	A	300	HEM	C3D-C4D	-3.46	1.47	1.51
2	B	300	HEM	C2C-C1C	-3.36	1.46	1.52
2	A	300	HEM	C2C-C1C	-3.07	1.46	1.52
2	B	300	HEM	C2B-C1B	-2.04	1.45	1.51
2	B	300	HEM	CHD-C4C	2.24	1.41	1.36
2	A	300	HEM	CHD-C4C	2.71	1.42	1.36
2	A	300	HEM	CBC-CAC	2.88	1.45	1.29
2	B	300	HEM	CBC-CAC	2.89	1.46	1.29
2	B	300	HEM	CBB-CAB	2.89	1.46	1.29
2	A	300	HEM	CBB-CAB	3.14	1.47	1.29
2	A	300	HEM	C4C-NC	3.51	1.40	1.36
2	B	300	HEM	C1C-NC	3.60	1.40	1.36
2	A	300	HEM	C1C-NC	3.65	1.40	1.36
2	B	300	HEM	C4C-NC	3.70	1.40	1.36

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	300	HEM	C3B-C4B-CHC	2.04	126.04	123.16
2	A	300	HEM	CMD-C2D-C3D	2.42	125.05	114.35
2	B	300	HEM	CMD-C2D-C3D	2.47	125.26	114.35
2	A	300	HEM	C2D-C3D-C4D	2.95	106.50	101.50
2	B	300	HEM	C2D-C3D-C4D	3.18	106.89	101.50
2	A	300	HEM	CMC-C2C-C3C	3.31	124.78	116.53
2	B	300	HEM	CAD-C3D-C4D	3.59	125.15	112.47
2	B	300	HEM	CMC-C2C-C3C	3.65	125.65	116.53
2	A	300	HEM	CMB-C2B-C3B	3.66	125.67	116.53
2	A	300	HEM	CBA-CAA-C2A	3.72	119.19	112.53
2	A	300	HEM	CAD-C3D-C4D	3.95	126.42	112.47
2	B	300	HEM	CMB-C2B-C3B	4.35	127.38	116.53
2	A	300	HEM	CAD-C3D-C2D	4.82	127.07	113.22
2	B	300	HEM	CAD-C3D-C2D	4.88	127.24	113.22

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	300	HEM	1	0
2	B	300	HEM	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	214/233 (91%)	0.39	15 (7%)	19 25	26, 40, 61, 65	10 (4%)
1	B	214/233 (91%)	0.53	19 (8%)	12 16	25, 38, 68, 71	11 (5%)
All	All	428/466 (91%)	0.46	34 (7%)	15 21	25, 39, 63, 71	21 (4%)

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	40	GLY	6.9
1	B	41	GLN	5.1
1	B	38	GLN	4.6
1	B	134	TYR	4.6
1	A	158	PRO	3.8
1	A	157	LEU	3.5
1	A	156	ASP	3.5
1	B	223	HIS	3.4
1	B	10	PRO	3.4
1	A	159	SER	3.3
1	A	11	GLN	3.2
1	A	10	PRO	3.2
1	B	35	ARG	3.2
1	B	149	LYS	2.9
1	B	150	ILE	2.8
1	A	153	LYS	2.7
1	A	155	LEU	2.7
1	B	36	ASN	2.6
1	B	54	LEU	2.6
1	B	37	PHE	2.5
1	B	146	VAL	2.5
1	A	134	TYR	2.5
1	A	40	GLY	2.4
1	A	22	LYS	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	32	GLU	2.4
1	B	27	GLN	2.3
1	A	180	GLN	2.3
1	B	39	LYS	2.3
1	B	137	TYR	2.3
1	B	133	ALA	2.3
1	A	154	ALA	2.2
1	A	41	GLN	2.2
1	A	42	VAL	2.2
1	B	222	THR	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	HEM	A	300	43/43	0.88	0.23	1.91	60,60,61,62	5
2	HEM	B	300	43/43	0.86	0.25	0.62	75,75,76,76	11
3	NO	A	400	2/2	0.94	0.09	-	60,60,60,60	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.