



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 10:26 PM GMT

PDB ID : 4XW6  
Title : X-ray structure of PKAc with ADP, free phosphate ion, CP20, magnesium ions  
Authors : Gerlits, O.; Tian, J.; Das, A.; Taylor, S.; Langan, P.; Heller, T.W.; Kovalevsky, A.  
Deposited on : 2015-01-28  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	4755 (1.90-1.90)
Clashscore	102246	5398 (1.90-1.90)
Ramachandran outliers	100387	5338 (1.90-1.90)
Sidechain outliers	100360	5339 (1.90-1.90)
RSRZ outliers	91569	4766 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	356	<div> <div>6%</div> <div>74%</div> <div>17%</div> <div>•• 5%</div> </div>
2	B	20	<div> <div>10%</div> <div>65%</div> <div>25%</div> <div>5% 5%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	PO4	A	404	-	-	X	-

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 3399 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called cAMP-dependent protein kinase catalytic subunit alpha.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	337	Total	C	N	O	P	S	0	3	0
			2809	1819	470	509	3	8			

There are 19 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	HIS	-	expression tag	UNP P05132
A	-4	HIS	-	expression tag	UNP P05132
A	-3	HIS	-	expression tag	UNP P05132
A	-2	HIS	-	expression tag	UNP P05132
A	-1	HIS	-	expression tag	UNP P05132
A	0	HIS	-	expression tag	UNP P05132
A	1	GLY	-	expression tag	UNP P05132
A	2	ASN	-	expression tag	UNP P05132
A	3	ALA	-	expression tag	UNP P05132
A	4	ALA	-	expression tag	UNP P05132
A	5	ALA	-	expression tag	UNP P05132
A	6	ALA	-	expression tag	UNP P05132
A	7	LYS	-	expression tag	UNP P05132
A	8	LYS	-	expression tag	UNP P05132
A	9	GLY	-	expression tag	UNP P05132
A	10	SER	-	expression tag	UNP P05132
A	11	GLU	-	expression tag	UNP P05132
A	12	GLN	-	expression tag	UNP P05132
A	13	GLU	-	expression tag	UNP P05132


- Molecule 2 is a protein called cAMP-dependent protein kinase inhibitor alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	20	Total	C	N	O	S	0	3	0
			179	108	36	33	2			

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Mg	0	0
			2	2		

- Molecule 4 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: C<sub>10</sub>H<sub>15</sub>N<sub>5</sub>O<sub>10</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 5 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	P	0	0
			5	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	346	Total	O	0	0
			346	346		
6	B	31	Total	O	0	0
			31	31		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: cAMP-dependent protein kinase catalytic subunit alpha



- Molecule 2: cAMP-dependent protein kinase inhibitor alpha



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	57.30Å 79.23Å 97.82Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.90 33.67 – 1.90	Depositor EDS
% Data completeness (in resolution range)	89.9 (20.00-1.90) 88.0 (33.67-1.90)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.29 (at 1.91Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, $R_{free}$	0.186 , 0.265 0.176 , 0.246	Depositor DCC
$R_{free}$ test set	1595 reflections (5.36%)	DCC
Wilson B-factor (Å <sup>2</sup> )	17.6	Xtriage
Anisotropy	0.091	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 76.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	1 of 32049 reflections (0.003%)	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3399	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.97% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, MG, PO4, ADP, SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.71	3/2850 (0.1%)	1.03	13/3834 (0.3%)
2	B	1.21	0/182	1.00	0/243
All	All	0.75	3/3032 (0.1%)	1.03	13/4077 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
2	B	0	5
All	All	0	6

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	101	PRO	N-CD	5.81	1.55	1.47
1	A	86	GLU	CD-OE2	-5.74	1.19	1.25
1	A	86	GLU	CD-OE1	-5.22	1.20	1.25

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	144	ARG	NE-CZ-NH2	9.50	125.05	120.30
1	A	144	ARG	CD-NE-CZ	9.06	136.29	123.60
1	A	321[A]	PRO	C-N-CA	8.19	139.49	122.30
1	A	321[B]	PRO	C-N-CA	8.19	139.49	122.30
1	A	144	ARG	NE-CZ-NH1	-7.73	116.44	120.30
1	A	166	ASP	CB-CG-OD1	7.07	124.67	118.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	241	ASP	CB-CG-OD1	6.88	124.50	118.30
1	A	217[A]	LYS	CA-C-N	-6.67	102.53	117.20
1	A	217[B]	LYS	CA-C-N	-6.67	102.53	117.20
1	A	270	ARG	NE-CZ-NH2	-6.16	117.22	120.30
1	A	217[A]	LYS	N-CA-C	6.15	127.62	111.00
1	A	217[B]	LYS	N-CA-C	6.15	127.62	111.00
1	A	270	ARG	NE-CZ-NH1	5.78	123.19	120.30

There are no chirality outliers.

All (6) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	284	LEU	Mainchain
2	B	620	ALA	Mainchain
2	B	623[A]	HIS	Mainchain,Peptide
2	B	623[B]	HIS	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2809	0	2798	68	0
2	B	179	0	166	23	0
3	A	2	0	0	0	0
4	A	27	0	12	2	0
5	A	5	0	0	4	0
6	A	346	0	0	22	0
6	B	31	0	0	2	0
All	All	3399	0	2976	82	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (82) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:404:PO4:O1	2:B:621[B]:CYS:SG	0.86	1.25
5:A:404:PO4:P	2:B:621[B]:CYS:SG	2.37	1.22
1:A:285[B]:LYS:HE3	1:A:285[B]:LYS:H	0.99	1.06
1:A:87:HIS:CE1	2:B:623[B]:HIS:CE1	2.43	1.05
2:B:622[B]:ILE:HG22	2:B:624:ASP:HB2	1.32	1.05
1:A:87:HIS:CE1	2:B:623[B]:HIS:HE1	1.75	1.05
1:A:87:HIS:HE1	2:B:623[B]:HIS:CE1	1.78	1.00
2:B:622[B]:ILE:CG2	2:B:624:ASP:HB2	1.92	0.99
1:A:285[B]:LYS:HE3	1:A:285[B]:LYS:N	1.82	0.94
5:A:404:PO4:O1	2:B:621[A]:CYS:HB3	1.74	0.88
1:A:253:GLY:O	6:A:818:HOH:O	1.98	0.81
1:A:336:ARG:HG2	6:A:766:HOH:O	1.81	0.79
1:A:336:ARG:CG	6:A:766:HOH:O	2.33	0.76
2:B:622[B]:ILE:HG22	2:B:623[B]:HIS:H	1.50	0.76
1:A:139:SEP:O2P	1:A:142:HIS:HB2	1.86	0.76
1:A:285[A]:LYS:N	1:A:285[A]:LYS:HD2	2.01	0.75
1:A:21:LYS:HG3	6:A:780:HOH:O	1.87	0.74
2:B:622[B]:ILE:HG22	2:B:623[B]:HIS:N	2.03	0.74
2:B:622[B]:ILE:CD1	6:B:726:HOH:O	2.37	0.72
1:A:336:ARG:NH1	6:A:766:HOH:O	2.23	0.72
1:A:303:ILE:O	1:A:307:GLN:HG3	1.91	0.71
1:A:163:ILE:HG12	1:A:217[A]:LYS:HG2	1.75	0.67
1:A:198:LEU:O	2:B:623[B]:HIS:HA	1.95	0.67
1:A:244:ILE:HG13	6:A:775:HOH:O	1.95	0.67
1:A:217[A]:LYS:HG3	6:A:811:HOH:O	1.93	0.67
1:A:336:ARG:CZ	6:A:766:HOH:O	2.43	0.66
1:A:52:GLY:HA3	4:A:403:ADP:O2B	1.95	0.66
1:A:192:LYS:HB2	6:A:507:HOH:O	1.98	0.64
1:A:14:SER:OG	1:A:17:GLU:HB2	1.98	0.64
2:B:622[B]:ILE:CG2	2:B:624:ASP:CB	2.74	0.62
1:A:285[A]:LYS:HD2	1:A:285[A]:LYS:H	1.66	0.61
1:A:336:ARG:HH12	1:A:338:SEP:HB3	1.68	0.58
2:B:622[B]:ILE:HD11	6:B:726:HOH:O	2.02	0.58
1:A:336:ARG:N	6:A:767:HOH:O	2.39	0.55
1:A:168:LYS:HZ3	5:A:404:PO4:P	2.30	0.54
1:A:333:GLU:HA	1:A:333:GLU:OE2	2.07	0.53
1:A:126:GLY:HA2	1:A:327:PHE:CZ	2.44	0.53
1:A:284:LEU:HB3	1:A:285[B]:LYS:CE	2.40	0.52
1:A:158:HIS:HD2	6:A:774:HOH:O	1.93	0.52
2:B:622[B]:ILE:HG22	2:B:624:ASP:CB	2.22	0.51
1:A:158:HIS:HE1	1:A:220:ASP:OD2	1.94	0.50
1:A:45:ARG:HA	1:A:60:VAL:HG12	1.91	0.50

Continued on next page...

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:335:ILE:HD13	1:A:335:ILE:O	2.10	0.50
2:B:622[B]:ILE:CG2	2:B:623[B]:HIS:H	2.16	0.50
1:A:278:THR:HG22	1:A:279:LYS:HG3	1.93	0.49
1:A:146:TYR:HB3	1:A:180:ILE:HD11	1.93	0.49
1:A:284:LEU:CB	1:A:285[A]:LYS:HD2	2.43	0.49
1:A:52:GLY:HA3	4:A:403:ADP:PB	2.53	0.48
1:A:158:HIS:HB3	1:A:217[B]:LYS:HE3	1.96	0.48
1:A:146:TYR:HB3	1:A:180:ILE:CD1	2.44	0.48
1:A:87:HIS:ND1	2:B:623[B]:HIS:HE1	2.08	0.48
1:A:194:ARG:HD2	1:A:214:GLY:HA3	1.96	0.47
1:A:99:ASN:HD22	1:A:105:LYS:NZ	2.13	0.46
1:A:270:ARG:HD3	6:A:842:HOH:O	2.16	0.46
1:A:186:GLY:HA3	6:A:713:HOH:O	2.15	0.46
1:A:286:ASN:O	1:A:289:ASN:HB2	2.16	0.46
1:A:284:LEU:HB2	1:A:285[A]:LYS:HD2	1.97	0.46
1:A:78:LYS:HE3	6:A:805:HOH:O	2.15	0.45
2:B:622[B]:ILE:HG21	2:B:624:ASP:HB2	1.93	0.45
1:A:78:LYS:HG2	6:A:769:HOH:O	2.16	0.45
1:A:21:LYS:HD2	1:A:21:LYS:C	2.37	0.45
1:A:161:ASP:HA	1:A:217[A]:LYS:HE2	2.00	0.44
1:A:284:LEU:HB3	1:A:285[B]:LYS:CD	2.47	0.44
1:A:95:LEU:HG	1:A:106:LEU:HB2	1.99	0.44
1:A:244:ILE:CD1	6:A:775:HOH:O	2.65	0.44
1:A:202:PRO:HD3	2:B:622[B]:ILE:HG12	1.99	0.44
1:A:339:ILE:HG12	6:A:501:HOH:O	2.17	0.43
1:A:315:ILE:HD11	6:A:747:HOH:O	2.18	0.43
1:A:317:LYS:HE2	6:A:783:HOH:O	2.18	0.43
1:A:84:GLN:HE22	2:B:623[B]:HIS:CE1	2.36	0.43
1:A:55:GLY:N	6:A:686:HOH:O	2.51	0.43
1:A:239:PHE:O	1:A:249:LYS:HE3	2.18	0.42
1:A:341:GLU:HG2	6:A:748:HOH:O	2.19	0.42
2:B:607:TYR:CZ	2:B:611:ILE:HG13	2.54	0.42
1:A:87:HIS:ND1	2:B:623[B]:HIS:CE1	2.85	0.41
1:A:284:LEU:HB3	1:A:285[B]:LYS:HD2	2.02	0.41
1:A:197:TPO:HG22	2:B:623[B]:HIS:NE2	2.36	0.41
1:A:215:TYR:CD2	1:A:219:VAL:HG11	2.55	0.41
1:A:21:LYS:O	1:A:21:LYS:HD2	2.21	0.40
1:A:244:ILE:CG1	6:A:775:HOH:O	2.63	0.40
1:A:292:LYS:HA	1:A:302:TRP:CZ2	2.57	0.40
1:A:107:GLU:N	1:A:119:VAL:O	2.50	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	335/356 (94%)	323 (96%)	10 (3%)	2 (1%)	30	17
2	B	21/20 (105%)	17 (81%)	0	4 (19%)	0	0
All	All	356/376 (95%)	340 (96%)	10 (3%)	6 (2%)	21	2

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	285[A]	LYS
1	A	285[B]	LYS
2	B	622[A]	ILE
2	B	622[B]	ILE
2	B	623[A]	HIS
2	B	623[B]	HIS

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	298/308 (97%)	283 (95%)	15 (5%)	30	18
2	B	18/15 (120%)	18 (100%)	0	100	100
All	All	316/323 (98%)	301 (95%)	15 (5%)	34	20

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	16	LYS
1	A	17	GLU
1	A	21	LYS
1	A	28	LYS
1	A	244	ILE
1	A	255	VAL
1	A	268	LEU
1	A	278	THR
1	A	284	LEU
1	A	285[A]	LYS
1	A	285[B]	LYS
1	A	331	GLU
1	A	333	GLU
1	A	335	ILE
1	A	345	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	36	ASN
1	A	68	HIS
1	A	84	GLN
1	A	99	ASN
1	A	113	ASN
1	A	158	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	SEP	A	139	1	8,9,10	1.61	2 (25%)	8,12,14	1.55	1 (12%)
1	TPO	A	197	1	8,10,11	1.26	1 (12%)	7,14,16	1.30	1 (14%)
1	SEP	A	338	1	8,9,10	1.26	2 (25%)	8,12,14	1.84	2 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SEP	A	139	1	-	0/6/8/10	0/0/0/0
1	TPO	A	197	1	-	0/8/11/13	0/0/0/0
1	SEP	A	338	1	-	0/6/8/10	0/0/0/0

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	338	SEP	P-O2P	2.03	1.62	1.54
1	A	338	SEP	P-O3P	2.03	1.62	1.54
1	A	139	SEP	P-O3P	2.13	1.62	1.54
1	A	197	TPO	P-O1P	2.84	1.60	1.51
1	A	139	SEP	P-O1P	3.53	1.62	1.51

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	338	SEP	O-C-CA	-2.04	120.17	125.49
1	A	197	TPO	C-CA-N	2.65	115.37	109.83
1	A	139	SEP	OG-CB-CA	2.93	110.77	108.27
1	A	338	SEP	OG-CB-CA	4.58	112.18	108.27

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	139	SEP	1	0
1	A	197	TPO	1	0
1	A	338	SEP	1	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	ADP	A	403	3	22,29,29	1.40	5 (22%)	27,45,45	1.59	3 (11%)
5	PO4	A	404	3	4,4,4	1.28	0	6,6,6	0.29	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ADP	A	403	3	-	0/12/32/32	0/3/3/3
5	PO4	A	404	3	-	0/0/0/0	0/0/0/0

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	403	ADP	PA-O1A	-2.14	1.43	1.51
4	A	403	ADP	PA-O2A	-2.10	1.46	1.54
4	A	403	ADP	PB-O3B	-2.09	1.47	1.54
4	A	403	ADP	C5-N7	-2.05	1.32	1.39
4	A	403	ADP	C5-C4	2.88	1.47	1.40

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	403	ADP	N3-C2-N1	-5.13	124.96	128.89
4	A	403	ADP	C4-C5-N7	-3.27	106.47	109.48

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	403	ADP	C2-N1-C6	2.29	122.85	118.77

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	403	ADP	2	0
5	A	404	PO4	4	0

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	334/356 (93%)	0.25	23 (6%) 20 22	12, 21, 55, 87	0
2	B	20/20 (100%)	0.48	2 (10%) 9 10	16, 20, 57, 113	0
All	All	354/376 (94%)	0.26	25 (7%) 19 21	12, 21, 56, 113	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	318	PHE	6.3
2	B	624	ASP	6.2
2	B	623[A]	HIS	5.8
1	A	14	SER	3.8
1	A	319	LYS	3.7
1	A	15	VAL	3.6
1	A	17	GLU	3.2
1	A	16	LYS	3.0
1	A	320	GLY	3.0
1	A	54	PHE	3.0
1	A	55	GLY	2.9
1	A	317	LYS	2.9
1	A	333	GLU	2.8
1	A	285[A]	LYS	2.7
1	A	316	PRO	2.5
1	A	253	GLY	2.5
1	A	321[A]	PRO	2.5
1	A	20	ALA	2.3
1	A	185	PHE	2.3
1	A	182	VAL	2.2
1	A	286	ASN	2.1
1	A	82	LEU	2.1
1	A	224	LEU	2.1
1	A	335	ILE	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	244	ILE	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
1	SEP	A	139	10/11	0.92	0.18	-	13,22,88,113	0
1	SEP	A	338	10/11	0.95	0.12	-	19,29,33,34	0
1	TPO	A	197	11/12	0.98	0.07	-	14,17,23,28	0

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
5	PO4	A	404	5/5	0.90	0.16	0.72	15,24,32,49	5
3	MG	A	401	1/1	0.86	0.13	-0.66	43,43,43,43	0
4	ADP	A	403	27/27	0.94	0.10	-0.67	15,19,44,52	0
3	MG	A	402	1/1	0.77	0.11	-	33,33,33,33	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.