



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 11:38 PM GMT

PDB ID : 1XXC
Title : C-TERMINAL DOMAIN OF ESCHERICHIA COLI ARGININE REPRES-
SOR
Authors : Van Duyne, G.D.; Ghosh, G.; Maas, W.K.; Sigler, P.B.
Deposited on : 1995-11-03
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	2827 (2.80-2.80)
Ramachandran outliers	100387	2782 (2.80-2.80)
Sidechain outliers	100360	2784 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	78	
1	B	78	
1	C	78	
1	D	78	
1	E	78	
1	F	78	

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 3234 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ARGININE REPRESSOR.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	72	Total	C	N	O	0	0	0
			539	349	84	106			
1	B	72	Total	C	N	O	0	0	0
			539	349	84	106			
1	C	72	Total	C	N	O	0	0	0
			539	349	84	106			
1	D	72	Total	C	N	O	0	0	0
			539	349	84	106			
1	E	72	Total	C	N	O	0	0	0
			539	349	84	106			
1	F	72	Total	C	N	O	0	0	0
			539	349	84	106			

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


• Molecule 1: ARGININE REPRESSOR




• Molecule 1: ARGININE REPRESSOR



• Molecule 1: ARGININE REPRESSOR



• Molecule 1: ARGININE REPRESSOR



L144
Y145
E146
A147
L148
L149
E150
L151
F152
ASP
GLN
GLU
LEU

● Molecule 1: ARGININE REPRESSOR



MET	SER	P81	L82	R83	R84	L85	V86	L87	D88	I89	D90	N91	N92	D93	A94	V95	V96	V97	I98	R99	T100	S101	P102	G103	A104	A105	Q106	L107	L111	L112	L115	G116	K117	G120	I121	L122	G123	T124	I125	D128	D129	T130	I131	F132	T133	T134	P135	A136	F139	T140	V141	K142	D143	L144
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------

Y145	E146	A147	L148	L149	E150	L151	F152	ASP	GLN	GLU	LEU
------	------	------	------	------	------	------	------	-----	-----	-----	-----

● Molecule 1: ARGININE REPRESSOR



MET	SER	P81	L82	R83	R84	L85	V86	L87	D88	I89	D90	N91	N92	D93	V96	V97	I98	R99	T100	S101	P102	A105	Q106	L107	I108	A109	R110	L111	L112	L115	G116	K117	A118	E119	G120	I121	L122	G123	T124	I125	D129	T130	I131	F132	T133	T134	P135	A136	F139	T140	V141	K142	D143	L144
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------

Y145	E146	L149	E150	L151	F152	ASP	GLN	GLU	LEU
------	------	------	------	------	------	-----	-----	-----	-----

4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	54.30 Å 86.70 Å 213.70 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 2.80	Depositor
% Data completeness (in resolution range)	(Not available) (8.00-2.80)	Depositor
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR 3.8	Depositor
R, R_{free}	0.210 , 0.370	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3234	wwPDB-VP
Average B, all atoms (Å ²)	67.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.72	0/547	0.99	0/746
1	B	0.75	0/547	1.03	2/746 (0.3%)
1	C	0.78	0/547	1.08	2/746 (0.3%)
1	D	0.67	0/547	0.98	0/746
1	E	0.74	0/547	0.97	1/746 (0.1%)
1	F	0.80	0/547	0.96	0/746
All	All	0.75	0/3282	1.00	5/4476 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	89	ILE	CB-CA-C	-6.04	99.52	111.60
1	B	96	VAL	N-CA-C	-5.67	95.68	111.00
1	E	89	ILE	N-CA-C	-5.66	95.72	111.00
1	C	101	SER	N-CA-C	-5.63	95.79	111.00
1	C	151	LEU	N-CA-C	-5.08	97.28	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	539	0	550	67	0
1	B	539	0	550	61	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	539	0	550	57	0
1	D	539	0	550	74	0
1	E	539	0	550	67	0
1	F	539	0	550	57	0
All	All	3234	0	3300	340	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 52.

The worst 5 of 340 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:115:LEU:HB2	1:E:121:ILE:HG13	1.39	1.04
1:B:115:LEU:HD11	1:B:151:LEU:HD22	1.41	1.02
1:D:105:ALA:HA	1:D:131:ILE:HD11	1.46	0.97
1:C:121:ILE:HG13	1:C:144:LEU:HD11	1.49	0.95
1:A:86:VAL:HA	1:A:100:THR:HG22	1.47	0.94

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	70/78 (90%)	53 (76%)	13 (19%)	4 (6%)	2	6
1	B	70/78 (90%)	58 (83%)	6 (9%)	6 (9%)	1	2
1	C	70/78 (90%)	51 (73%)	18 (26%)	1 (1%)	14	42
1	D	70/78 (90%)	51 (73%)	12 (17%)	7 (10%)	1	1
1	E	70/78 (90%)	58 (83%)	10 (14%)	2 (3%)	6	19
1	F	70/78 (90%)	55 (79%)	11 (16%)	4 (6%)	2	6

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	420/468 (90%)	326 (78%)	70 (17%)	24 (6%)	2 6

5 of 24 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	94	ALA
1	B	83	LYS
1	B	129	ASP
1	B	136	ALA
1	C	105	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	58/64 (91%)	43 (74%)	15 (26%)	0 2
1	B	58/64 (91%)	41 (71%)	17 (29%)	0 1
1	C	58/64 (91%)	44 (76%)	14 (24%)	1 2
1	D	58/64 (91%)	45 (78%)	13 (22%)	1 3
1	E	58/64 (91%)	47 (81%)	11 (19%)	2 5
1	F	58/64 (91%)	47 (81%)	11 (19%)	2 5
All	All	348/384 (91%)	267 (77%)	81 (23%)	1 3

5 of 81 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	93	ASP
1	C	150	GLU
1	F	93	ASP
1	C	107	LEU
1	C	114	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	99	HIS
1	B	106	GLN
1	F	84	ASN
1	F	99	HIS
1	F	106	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.