



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 06:41 AM GMT

PDB ID : 2XYW
Title : NOVEL SULFONYLTHIA DIAZOLES WITH AN UNUSUAL BINDING
MODE AS PARTIAL DUAL PEROXISOME PROLIFERATOR-ACTIV
ATED RECEPTOR (PPAR) GAMMA-DELTA AGONISTS WITH HIGH
POTENCY AND IN-VIVO EFFICACY
Authors : Marquette, J.-P.; Mathieu, M.
Deposited on : 2010-11-19
Resolution : 3.14 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.14 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1095 (3.18-3.10)
Clashscore	102246	1202 (3.18-3.10)
Ramachandran outliers	100387	1162 (3.18-3.10)
Sidechain outliers	100360	1162 (3.18-3.10)
RSRZ outliers	91569	1097 (3.18-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	288	<div> <div>5%</div> <div>73%</div> <div>17%</div> <div>9%</div> </div>
1	B	288	<div> <div>2%</div> <div>75%</div> <div>15%</div> <div>9%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	BOG	A	1442	-	-	-	X
3	BOG	B	1442	-	-	-	X

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4352 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PEROXISOME PROLIFERATOR-ACTIVATED RECEPTOR DELTA.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	263	Total	C	N	O	S	0	0	0
			2122	1376	357	379	10			
1	B	263	Total	C	N	O	S	0	0	0
			2122	1376	357	379	10			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	154	MET	-	EXPRESSION TAG	UNP Q03181
A	155	LYS	-	EXPRESSION TAG	UNP Q03181
A	156	LYS	-	EXPRESSION TAG	UNP Q03181
A	157	GLY	-	EXPRESSION TAG	UNP Q03181
A	158	HIS	-	EXPRESSION TAG	UNP Q03181
A	159	HIS	-	EXPRESSION TAG	UNP Q03181
A	160	HIS	-	EXPRESSION TAG	UNP Q03181
A	161	HIS	-	EXPRESSION TAG	UNP Q03181
A	162	HIS	-	EXPRESSION TAG	UNP Q03181
A	163	HIS	-	EXPRESSION TAG	UNP Q03181
A	164	GLY	-	EXPRESSION TAG	UNP Q03181
B	154	MET	-	EXPRESSION TAG	UNP Q03181
B	155	LYS	-	EXPRESSION TAG	UNP Q03181
B	156	LYS	-	EXPRESSION TAG	UNP Q03181
B	157	GLY	-	EXPRESSION TAG	UNP Q03181
B	158	HIS	-	EXPRESSION TAG	UNP Q03181
B	159	HIS	-	EXPRESSION TAG	UNP Q03181
B	160	HIS	-	EXPRESSION TAG	UNP Q03181
B	161	HIS	-	EXPRESSION TAG	UNP Q03181
B	162	HIS	-	EXPRESSION TAG	UNP Q03181
B	163	HIS	-	EXPRESSION TAG	UNP Q03181
B	164	GLY	-	EXPRESSION TAG	UNP Q03181


- Molecule 2 is 3-CHLORO-6-FLUORO-N-[2-[4-[(5-PROPAN-2-YL-1,3,4-THIADIAZOL

-2-YL)SULFAMOYL]PHENYL]ETHYL]-1-BENZOTHIOPHENE-2-CARBOXAMIDE
(three-letter code: 08S) (formula: $C_{22}H_{20}ClFN_4O_3S_3$).



Mol	Chain	Residues	Atoms							ZeroOcc	AltConf
2	A	1	Total	C	Cl	F	N	O	S	0	0
			34	22	1	1	4	3	3		
2	B	1	Total	C	Cl	F	N	O	S	0	0
			34	22	1	1	4	3	3		

- Molecule 3 is SUGAR (B-OCTYLGLUCOSIDE) (three-letter code: BOG) (formula: $C_{14}H_{28}O_6$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			20	14	6		
3	B	1	Total	C	O	0	0
			20	14	6		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: PEROXISOME PROLIFERATOR-ACTIVATED RECEPTOR DELTA



- Molecule 1: PEROXISOME PROLIFERATOR-ACTIVATED RECEPTOR DELTA



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	39.37Å 93.22Å 97.47Å 90.00° 97.31° 90.00°	Depositor
Resolution (Å)	14.99 – 3.14 14.99 – 3.14	Depositor EDS
% Data completeness (in resolution range)	99.7 (14.99-3.14) 99.8 (14.99-3.14)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.27 (at 3.12Å)	Xtriage
Refinement program	BUSTER 2.9.3	Depositor
R, R_{free}	0.188 , 0.246 0.203 , 0.257	Depositor DCC
R_{free} test set	581 reflections (5.03%)	DCC
Wilson B-factor (Å ²)	59.0	Xtriage
Anisotropy	0.097	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 71.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	2 of 12134 reflections (0.016%)	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	4352	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.47% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 08S, BOG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.48	0/2167	0.68	0/2927
1	B	0.48	0/2167	0.69	0/2927
All	All	0.48	0/4334	0.69	0/5854

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2122	0	2171	21	0
1	B	2122	0	2171	19	0
2	A	34	0	20	4	0
2	B	34	0	20	2	0
3	A	20	0	28	1	0
3	B	20	0	28	4	0
All	All	4352	0	4438	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:189:ASN:OD1	1:A:262:GLU:HB3	2.03	0.58
1:A:261:THR:HG21	3:A:1442:BOG:H1'2	1.87	0.57
1:B:294:LEU:HD12	1:B:297:ILE:HD11	1.89	0.54
1:A:294:LEU:HD12	1:A:297:ILE:HD11	1.90	0.53
1:B:282:LEU:O	1:B:286:VAL:HB	2.13	0.49
1:B:385:LEU:HD11	1:B:399:LEU:HD12	1.96	0.48
1:B:317:LEU:HD21	2:B:1441:08S:H9	1.95	0.48
1:A:254:VAL:HG21	1:A:430:HIS:CD2	2.49	0.47
1:B:258:ARG:HA	3:B:1442:BOG:H1'1	1.96	0.47
1:A:282:LEU:O	1:A:286:VAL:HB	2.14	0.47
1:A:245:VAL:HG13	2:A:1441:08S:H9	1.96	0.46
1:B:432:LEU:CD1	3:B:1442:BOG:H8'2	2.46	0.46
1:A:328:ILE:HD12	2:A:1441:08S:H5	1.98	0.46
1:A:396:PHE:HB3	1:A:397:PRO:HD3	1.98	0.46
1:A:343:LEU:HD11	1:A:399:LEU:HD21	1.98	0.46
1:A:201:LEU:HD22	1:A:299:ASN:HD22	1.79	0.46
1:B:201:LEU:HD22	1:B:299:ASN:HD22	1.81	0.45
1:A:385:LEU:HD11	1:A:399:LEU:HD12	1.98	0.45
2:A:1441:08S:CL	2:A:1441:08S:O27	2.71	0.45
1:B:242:SER:HA	1:B:320:LEU:HD22	1.99	0.44
1:B:396:PHE:HB3	1:B:397:PRO:HD3	1.99	0.44
1:B:235:LEU:HD22	1:B:247:TYR:HB3	2.00	0.44
1:A:245:VAL:HG13	2:A:1441:08S:C9	2.48	0.44
1:B:348:LEU:O	1:B:352:ILE:HG12	2.18	0.43
1:B:435:GLU:O	1:B:438:LYS:HB3	2.18	0.43
1:A:242:SER:HA	1:A:320:LEU:HD22	1.99	0.43
1:A:348:LEU:O	1:A:352:ILE:HG12	2.18	0.43
1:A:288:GLU:HG2	1:A:410:VAL:HG21	1.99	0.43
1:B:288:GLU:HG2	1:B:410:VAL:HG21	2.00	0.42
1:A:235:LEU:HD22	1:A:247:TYR:HB3	2.01	0.42
1:B:343:LEU:HD11	1:B:399:LEU:HD21	2.00	0.42
1:A:385:LEU:HD22	1:A:395:LEU:HD13	2.02	0.42
1:A:360:ASP:HB2	1:A:407:ARG:HH12	1.85	0.42
2:B:1441:08S:O27	2:B:1441:08S:CL	2.75	0.41
1:B:385:LEU:HD22	1:B:395:LEU:HD13	2.02	0.41
1:B:261:THR:HG21	3:B:1442:BOG:H1'2	2.02	0.41
1:B:258:ARG:HG3	3:B:1442:BOG:H1'1	2.01	0.41
1:B:184:ASN:O	1:B:188:LYS:HB2	2.21	0.41
1:A:185:ALA:O	1:A:189:ASN:HB2	2.20	0.40
1:B:320:LEU:O	1:B:325:SER:HB3	2.21	0.40
1:A:320:LEU:O	1:A:325:SER:HB3	2.22	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:253:THR:O	1:A:257:VAL:HG23	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	259/288 (90%)	247 (95%)	10 (4%)	2 (1%)	24	64
1	B	259/288 (90%)	247 (95%)	10 (4%)	2 (1%)	24	64
All	All	518/576 (90%)	494 (95%)	20 (4%)	4 (1%)	24	64

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	227	VAL
1	B	438	LYS
1	A	438	LYS
1	A	227	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	233/253 (92%)	211 (91%)	22 (9%)	11	39

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	233/253 (92%)	215 (92%)	18 (8%)	16	50
All	All	466/506 (92%)	426 (91%)	40 (9%)	13	45

All (40) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	198	ARG
1	A	199	SER
1	A	219	LEU
1	A	221	GLN
1	A	230	GLN
1	A	239	LYS
1	A	248	ARG
1	A	262	GLU
1	A	269	SER
1	A	271	SER
1	A	283	LYS
1	A	286	VAL
1	A	357	LEU
1	A	358	CYS
1	A	360	ASP
1	A	367	VAL
1	A	395	LEU
1	A	412	GLU
1	A	425	THR
1	A	429	LEU
1	A	434	GLN
1	A	440	MET
1	B	198	ARG
1	B	199	SER
1	B	219	LEU
1	B	221	GLN
1	B	230	GLN
1	B	239	LYS
1	B	248	ARG
1	B	262	GLU
1	B	269	SER
1	B	271	SER
1	B	286	VAL
1	B	357	LEU
1	B	360	ASP
1	B	367	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	395	LEU
1	B	412	GLU
1	B	425	THR
1	B	434	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	08S	A	1441	-	29,37,37	2.29	8 (27%)	32,54,54	3.87	21 (65%)
3	BOG	A	1442	-	20,20,20	2.33	8 (40%)	25,25,25	2.10	10 (40%)
2	08S	B	1441	-	29,37,37	2.54	10 (34%)	32,54,54	4.11	21 (65%)
3	BOG	B	1442	-	20,20,20	1.95	6 (30%)	25,25,25	1.87	9 (36%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	08S	A	1441	-	-	0/17/25/25	0/3/4/4
3	BOG	A	1442	-	-	0/11/31/31	0/1/1/1
2	08S	B	1441	-	-	0/17/25/25	0/3/4/4
3	BOG	B	1442	-	-	0/11/31/31	0/1/1/1

All (32) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1441	08S	C13-S33	-4.91	1.69	1.76
2	A	1441	08S	C13-S33	-3.96	1.70	1.76
2	B	1441	08S	S33-N26	-3.85	1.57	1.63
2	B	1441	08S	C17-CL	-3.32	1.60	1.73
2	A	1441	08S	F30-C14	-3.01	1.29	1.36
2	B	1441	08S	C19-C21	-2.94	1.45	1.50
2	A	1441	08S	C17-CL	-2.86	1.62	1.73
2	B	1441	08S	C19-S32	-2.56	1.65	1.73
2	B	1441	08S	F30-C14	-2.52	1.30	1.36
2	A	1441	08S	O29-S33	-2.28	1.41	1.43
2	A	1441	08S	C20-N26	2.20	1.43	1.40
3	A	1442	BOG	C1-C2	2.30	1.59	1.52
2	B	1441	08S	C4-C12	2.36	1.43	1.38
3	B	1442	BOG	O4-C4	2.43	1.48	1.43
3	A	1442	BOG	C4-C3	2.66	1.59	1.52
3	A	1442	BOG	O4-C4	2.93	1.50	1.43
3	A	1442	BOG	C4-C5	2.98	1.59	1.53
3	B	1442	BOG	C4-C3	3.03	1.60	1.52
3	A	1442	BOG	C3-C2	3.14	1.60	1.52
3	B	1442	BOG	O5-C5	3.21	1.52	1.44
3	B	1442	BOG	O5-C1	3.23	1.50	1.41
3	B	1442	BOG	C4-C5	3.64	1.60	1.53
3	B	1442	BOG	O1-C1	3.64	1.46	1.40
2	A	1441	08S	C3-C12	3.65	1.46	1.38
2	B	1441	08S	C9-C16	3.76	1.43	1.37
3	A	1442	BOG	O5-C5	3.83	1.53	1.44
2	B	1441	08S	C3-C12	3.95	1.47	1.38
2	A	1441	08S	C9-C16	4.00	1.44	1.37
3	A	1442	BOG	O5-C1	4.02	1.52	1.41
3	A	1442	BOG	O1-C1	5.14	1.49	1.40
2	A	1441	08S	C9-C14	7.16	1.48	1.35

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1441	08S	C9-C14	7.88	1.50	1.35

All (61) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1441	08S	F30-C14-C7	-9.55	102.58	118.52
2	A	1441	08S	O29-S33-O28	-9.03	107.56	119.54
2	A	1441	08S	F30-C14-C7	-8.67	104.06	118.52
2	B	1441	08S	O29-S33-O28	-8.21	108.65	119.54
2	B	1441	08S	C8-C15-C17	-7.30	119.40	135.20
2	A	1441	08S	C8-C15-C17	-6.24	121.71	135.20
2	B	1441	08S	C1-C21-C19	-5.34	95.43	112.01
2	B	1441	08S	C6-C13-S33	-5.08	113.90	119.78
2	B	1441	08S	O27-C22-N25	-4.36	113.41	122.58
2	B	1441	08S	C11-N25-C22	-4.25	112.64	122.15
2	A	1441	08S	O27-C22-N25	-4.14	113.88	122.58
2	B	1441	08S	O28-S33-C13	-4.04	102.84	107.96
2	B	1441	08S	O29-S33-C13	-3.70	103.28	107.96
2	A	1441	08S	C6-C13-S33	-3.64	115.58	119.78
2	A	1441	08S	C11-N25-C22	-3.60	114.09	122.15
2	A	1441	08S	C1-C21-C19	-3.59	100.87	112.01
3	B	1442	BOG	O5-C1-C2	-3.13	103.86	110.28
2	A	1441	08S	C11-C10-C12	-3.12	106.35	112.83
2	A	1441	08S	O28-S33-C13	-2.95	104.23	107.96
2	A	1441	08S	O29-S33-C13	-2.74	104.48	107.96
2	A	1441	08S	C8-C7-C14	-2.65	116.38	118.77
2	A	1441	08S	C6-C4-C12	-2.58	117.50	121.04
3	A	1442	BOG	O5-C1-C2	-2.54	105.06	110.28
2	B	1441	08S	C7-C8-C15	-2.35	118.04	121.13
3	A	1442	BOG	O6-C6-C5	2.06	118.14	111.33
2	A	1441	08S	C4-C12-C3	2.07	121.44	118.13
3	B	1442	BOG	O3-C3-C4	2.12	115.10	110.34
2	A	1441	08S	O29-S33-N26	2.12	112.08	106.69
3	A	1442	BOG	O1-C1'-C2'	2.22	118.72	109.88
3	A	1442	BOG	O5-C5-C4	2.23	113.87	109.68
2	B	1441	08S	C5-C13-S33	2.31	122.46	119.78
2	A	1441	08S	C6-C13-C5	2.35	123.56	120.42
2	B	1441	08S	C10-C11-N25	2.38	117.78	111.97
2	B	1441	08S	C6-C13-C5	2.40	123.63	120.42
3	A	1442	BOG	C1-C2-C3	2.43	114.77	109.97
3	B	1442	BOG	O4-C4-C3	2.45	115.86	110.34
2	B	1441	08S	O29-S33-N26	2.49	113.00	106.69

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1442	BOG	O6-C6-C5	2.58	119.85	111.33
3	B	1442	BOG	O5-C5-C6	2.61	112.94	106.36
3	A	1442	BOG	C1-O5-C5	2.61	118.81	113.75
3	A	1442	BOG	O2-C2-C3	2.65	116.30	110.34
3	B	1442	BOG	C1-O5-C5	2.67	118.94	113.75
3	B	1442	BOG	O2-C2-C1	2.71	115.97	110.02
3	B	1442	BOG	C4-C3-C2	2.72	115.86	110.79
3	A	1442	BOG	O4-C4-C3	3.06	117.23	110.34
3	B	1442	BOG	C3-C4-C5	3.25	115.86	110.20
2	B	1441	08S	C13-S33-N26	3.43	111.28	106.87
2	A	1441	08S	O27-C22-C18	3.56	125.69	121.02
2	A	1441	08S	C7-C14-C9	3.64	127.73	123.51
2	B	1441	08S	O27-C22-C18	3.66	125.84	121.02
3	A	1442	BOG	C4-C3-C2	3.85	117.97	110.79
2	B	1441	08S	O28-S33-N26	3.88	116.52	106.69
2	B	1441	08S	C7-C14-C9	4.13	128.30	123.51
2	B	1441	08S	C21-C19-N23	4.88	130.52	120.67
2	A	1441	08S	C21-C19-N23	4.92	130.59	120.67
3	A	1442	BOG	O5-C5-C6	5.03	119.06	106.36
2	A	1441	08S	F30-C14-C9	5.90	128.21	119.07
2	A	1441	08S	O28-S33-N26	6.08	122.11	106.69
2	A	1441	08S	C18-C22-N25	6.14	120.41	115.89
2	B	1441	08S	F30-C14-C9	6.48	129.10	119.07
2	B	1441	08S	C18-C22-N25	6.55	120.71	115.89

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1441	08S	4	0
3	A	1442	BOG	1	0
2	B	1441	08S	2	0
3	B	1442	BOG	4	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	263/288 (91%)	-0.08	13 (4%) 33 14	26, 57, 91, 109	0
1	B	263/288 (91%)	-0.28	6 (2%) 64 42	25, 49, 80, 113	0
All	All	526/576 (91%)	-0.18	19 (3%) 46 23	25, 54, 88, 113	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	229	LYS	5.1
1	B	393	GLN	4.5
1	B	233	ASN	4.3
1	B	204	LYS	3.1
1	A	394	TYR	3.0
1	A	227	VAL	2.9
1	A	393	GLN	2.8
1	A	383	PHE	2.7
1	A	233	ASN	2.6
1	A	229	LYS	2.6
1	A	230	GLN	2.6
1	A	231	LEU	2.5
1	A	268	PRO	2.4
1	A	204	LYS	2.2
1	A	391	ASP	2.2
1	A	439	ASP	2.1
1	B	268	PRO	2.1
1	A	180	LYS	2.0
1	B	203	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	BOG	B	1442	20/20	0.68	0.44	4.43	46,100,106,107	0
3	BOG	A	1442	20/20	0.73	0.39	4.05	40,85,89,90	0
2	08S	B	1441	34/34	0.96	0.16	-0.07	42,50,58,64	0
2	08S	A	1441	34/34	0.94	0.17	-0.18	52,58,68,72	0

6.5 Other polymers [i](#)

There are no such residues in this entry.