



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 06:43 AM GMT

PDB ID : 2Y3Y
Title : HOLO-NI(II) HPNIKR IS A SYMMETRIC TETRAMER CONTAINING
FOUR CANONIC SQUARE-PLANAR NI(II) IONS AT PHYSIOLOGICAL
PH
Authors : Benini, S.; Cianci, M.; Ciurli, S.
Deposited on : 2011-01-04
Resolution : 2.39 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.39 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	91	<div> <div>73%</div> <div>14%</div> <div>•</div> <div>9%</div> </div>
1	B	91	<div> <div>71%</div> <div>18%</div> <div>•</div> <div>9%</div> </div>
1	C	91	<div> <div>76%</div> <div>10%</div> <div>•</div> <div>11%</div> </div>
1	D	91	<div> <div>2%</div> <div>58%</div> <div>27%</div> <div>5%</div> <div>9%</div> </div>
2	Q	11	<div> <div>82%</div> <div>18%</div> </div>

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 2807 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PUTATIVE NICKEL-RESPONSIVE REGULATOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	83	Total	C	N	O	S	0	0	0
			661	412	123	122	4			
1	B	83	Total	C	N	O	S	0	0	0
			662	412	123	123	4			
1	C	81	Total	C	N	O	S	0	0	0
			648	405	120	119	4			
1	D	83	Total	C	N	O	S	0	0	0
			662	412	123	123	4			

- Molecule 2 is a protein called UNDECAPEPTIDE-GSSSGSASGAG.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	Q	11	Total	C	N	O	0	0	1
			53	28	10	15			

- Molecule 3 is NICKEL (II) ION (three-letter code: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Ni	0	0
			1	1		
3	A	1	Total	Ni	0	0
			1	1		
3	D	1	Total	Ni	0	0
			1	1		
3	C	1	Total	Ni	0	0
			1	1		

- Molecule 4 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (three-letter code: EPE) (formula: C₈H₁₈N₂O₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total	C	N	O	S	0	0
			15	8	2	4	1		
4	C	1	Total	C	N	O	S	0	0
			15	8	2	4	1		

- Molecule 5 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	C	1	Total	C	O	0	0
			7	4	3		
5	C	1	Total	C	O	0	0
			7	4	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	C	1	Total	C	O	0	0
			7	4	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	18	Total	O	0	0
			18	18		
6	B	17	Total	O	0	0
			17	17		
6	C	17	Total	O	0	0
			17	17		
6	D	13	Total	O	0	0
			13	13		
6	Q	1	Total	O	0	0
			1	1		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: PUTATIVE NICKEL-RESPONSIVE REGULATOR



- Molecule 1: PUTATIVE NICKEL-RESPONSIVE REGULATOR




- Molecule 1: PUTATIVE NICKEL-RESPONSIVE REGULATOR



- Molecule 1: PUTATIVE NICKEL-RESPONSIVE REGULATOR



- Molecule 2: UNDECAPEPTIDE-GSSSGSASGAG



4 Data and refinement statistics

Property	Value	Source
Space group	P 43	Depositor
Cell constants a, b, c, α , β , γ	72.97 Å 72.97 Å 116.73 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	72.98 – 2.39 19.94 – 2.39	Depositor EDS
% Data completeness (in resolution range)	94.8 (72.98-2.39) 95.0 (19.94-2.39)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.67 (at 2.38 Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.229 , 0.279 0.226 , 0.274	Depositor DCC
R_{free} test set	1177 reflections (5.41%)	DCC
Wilson B-factor (Å ²)	42.7	Xtriage
Anisotropy	0.103	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.26 , 12.9	EDS
Estimated twinning fraction	0.489 for h,-k,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 22943 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	2807	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, NI, EPE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.15	1/670 (0.1%)	1.03	1/902 (0.1%)
1	B	1.29	0/671	1.12	4/903 (0.4%)
1	C	1.19	2/657 (0.3%)	1.09	1/884 (0.1%)
1	D	1.16	4/671 (0.6%)	1.14	4/903 (0.4%)
2	Q	0.74	0/52	0.72	0/68
All	All	1.19	7/2721 (0.3%)	1.09	10/3660 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	119	GLU	CD-OE1	5.88	1.32	1.25
1	D	80	ASN	CB-CG	5.59	1.64	1.51
1	A	62	GLU	CB-CG	5.54	1.62	1.52
1	D	135	PHE	CE1-CZ	5.36	1.47	1.37
1	D	104	GLU	CD-OE2	5.33	1.31	1.25
1	C	66	ALA	CA-CB	5.05	1.63	1.52
1	D	109	GLU	CG-CD	-5.04	1.44	1.51

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	103	ASP	CB-CG-OD1	7.27	124.84	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	108	LEU	CA-CB-CG	7.23	131.93	115.30
1	D	125	LEU	CB-CG-CD1	-6.25	100.38	111.00
1	B	108	LEU	CA-CB-CG	6.06	129.24	115.30
1	D	103	ASP	CB-CG-OD1	5.86	123.58	118.30
1	D	123	LEU	CB-CG-CD2	5.85	120.95	111.00
1	C	122	ARG	NE-CZ-NH1	5.79	123.20	120.30
1	B	138	LEU	CB-CG-CD2	-5.57	101.54	111.00
1	B	108	LEU	CB-CG-CD1	5.21	119.86	111.00
1	A	73	ASP	CB-CG-OD1	5.08	122.88	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	60	ASN	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	661	0	660	18	0
1	B	662	0	663	16	0
1	C	648	0	652	10	0
1	D	662	0	663	16	0
2	Q	53	0	42	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	B	15	0	17	6	0
4	C	15	0	17	6	0
5	C	21	0	30	0	0
6	A	18	0	0	2	0
6	B	17	0	0	1	0
6	C	17	0	0	2	0
6	D	13	0	0	1	0
6	Q	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	2807	0	2744	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:C:1143:EPE:H52	4:C:1143:EPE:H102	1.33	1.05
1:A:60:ASN:HB3	1:A:62:GLU:OE1	1.72	0.90
1:A:116:ASN:ND2	1:A:118:PHE:H	1.70	0.88
1:A:116:ASN:HD22	1:A:118:PHE:H	1.21	0.86
1:B:110:THR:HG21	1:C:110:THR:HG21	1.57	0.86
4:B:1144:EPE:O8	4:B:1144:EPE:C5	2.22	0.85
4:C:1143:EPE:O3S	4:C:1143:EPE:H22	1.77	0.85
4:C:1143:EPE:S	4:C:1143:EPE:H22	2.21	0.80
1:D:92:THR:HG21	1:D:113:LEU:HD22	1.66	0.78
4:C:1143:EPE:H102	4:C:1143:EPE:C5	2.09	0.77
4:B:1144:EPE:N4	4:B:1144:EPE:H91	2.00	0.76
4:B:1144:EPE:O8	4:B:1144:EPE:H51	1.86	0.74
4:B:1144:EPE:N1	4:B:1144:EPE:O1S	2.18	0.74
1:B:86:ILE:HG22	1:B:123:LEU:CD1	2.22	0.70
1:B:110:THR:HG21	1:C:110:THR:CG2	2.24	0.67
1:C:61:ASP:N	6:C:2001:HOH:O	2.31	0.64
1:A:116:ASN:ND2	1:A:118:PHE:N	2.45	0.64
4:C:1143:EPE:C2	4:C:1143:EPE:O3S	2.46	0.63
1:D:86:ILE:HG22	1:D:123:LEU:HD11	1.80	0.62
4:B:1144:EPE:C9	4:B:1144:EPE:N4	2.62	0.61
1:C:110:THR:HG22	1:C:110:THR:O	2.01	0.61
1:A:116:ASN:HD22	1:A:118:PHE:N	1.96	0.59
1:A:116:ASN:C	1:A:116:ASN:HD22	2.05	0.59
1:B:86:ILE:HG22	1:B:123:LEU:HD11	1.84	0.58
1:B:76:GLN:NE2	1:B:131:ARG:HD2	2.19	0.57
1:B:131:ARG:HG3	6:B:2015:HOH:O	2.05	0.56
1:A:122:ARG:CZ	6:A:2015:HOH:O	2.54	0.56
1:C:99:HIS:CD2	1:C:101:HIS:CE1	2.96	0.55
1:D:65:ILE:H	1:D:142:SER:HB2	1.74	0.53
1:B:82:ARG:O	1:B:86:ILE:HG12	2.08	0.52
1:A:116:ASN:C	1:A:116:ASN:ND2	2.63	0.51
1:A:116:ASN:HD22	1:A:117:SER:N	2.08	0.51
4:C:1143:EPE:H52	4:C:1143:EPE:C10	2.25	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:1144:EPE:H52	4:B:1144:EPE:O8	2.08	0.51
1:D:82:ARG:HD2	6:D:2003:HOH:O	2.11	0.50
1:D:77:ARG:O	1:D:78:GLU:HB3	2.10	0.50
1:C:101:HIS:HE2	1:D:87:GLN:NE2	2.10	0.50
1:A:60:ASN:HB2	1:A:116:ASN:HB2	1.94	0.49
1:B:82:ARG:HH12	1:B:86:ILE:CD1	2.26	0.49
1:C:80:ASN:HB3	1:D:80:ASN:HD22	1.78	0.48
1:D:99:HIS:CE1	1:D:107:CYS:HB3	2.48	0.48
1:A:141:ALA:HB1	1:D:71:ILE:CD1	2.43	0.48
1:B:64:LYS:O	1:B:114:GLN:HA	2.13	0.47
1:B:65:ILE:HD13	1:B:65:ILE:N	2.29	0.47
1:B:76:GLN:HE22	1:B:131:ARG:HD2	1.78	0.47
1:C:110:THR:CG2	1:C:110:THR:O	2.63	0.47
1:A:141:ALA:HB1	1:D:71:ILE:HD11	1.97	0.47
1:D:123:LEU:HA	1:D:123:LEU:HD12	1.83	0.46
1:D:141:ALA:O	1:D:142:SER:HB3	2.15	0.46
1:B:82:ARG:HH12	1:B:86:ILE:HD11	1.81	0.46
1:D:106:ASN:HD21	1:D:134:LYS:NZ	2.14	0.45
1:A:122:ARG:NH1	6:A:2015:HOH:O	2.52	0.43
1:D:78:GLU:O	1:D:78:GLU:HG2	2.19	0.43
1:D:99:HIS:HA	1:D:108:LEU:O	2.19	0.43
1:A:80:ASN:HB3	1:B:80:ASN:HD22	1.84	0.42
1:A:97:THR:OG1	1:B:99:HIS:HB3	2.20	0.42
1:C:82:ARG:HG3	6:C:2006:HOH:O	2.21	0.41
1:B:65:ILE:HG13	1:C:102:MET:CE	2.51	0.41
1:A:99:HIS:CD2	1:A:101:HIS:CE1	3.09	0.41
1:A:86:ILE:HD13	1:A:126:GLU:HB3	2.02	0.41
1:A:88:HIS:CE1	1:B:107:CYS:SG	3.14	0.41
1:D:64:LYS:O	1:D:114:GLN:HA	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	81/91 (89%)	77 (95%)	4 (5%)	0	100	100
1	B	81/91 (89%)	76 (94%)	5 (6%)	0	100	100
1	C	79/91 (87%)	77 (98%)	2 (2%)	0	100	100
1	D	81/91 (89%)	74 (91%)	5 (6%)	2 (2%)	7	7
2	Q	9/11 (82%)	9 (100%)	0	0	100	100
All	All	331/375 (88%)	313 (95%)	16 (5%)	2 (1%)	30	43

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	61	ASP
1	D	135	PHE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	73/82 (89%)	69 (94%)	4 (6%)	27	42
1	B	74/82 (90%)	69 (93%)	5 (7%)	20	31
1	C	72/82 (88%)	67 (93%)	5 (7%)	19	30
1	D	74/82 (90%)	69 (93%)	5 (7%)	20	31
2	Q	5/5 (100%)	3 (60%)	2 (40%)	0	0
All	All	298/333 (90%)	277 (93%)	21 (7%)	19	29

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	116	ASN
1	A	122	ARG
1	A	131	ARG
1	A	140	LYS
1	B	65	ILE
1	B	85	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	92	THR
1	B	108	LEU
1	B	137	LYS
1	C	82	ARG
1	C	108	LEU
1	C	110	THR
1	C	122	ARG
1	C	125	LEU
1	D	62	GLU
1	D	63	SER
1	D	65	ILE
1	D	81	GLN
1	D	131	ARG
2	Q	2	SER
2	Q	6	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	60	ASN
1	A	81	GLN
1	A	106	ASN
1	A	116	ASN
1	B	74	HIS
1	B	76	GLN
1	B	114	GLN
1	C	106	ASN
1	D	80	ASN
1	D	87	GLN
1	D	106	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 9 ligands modelled in this entry, 4 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	EPE	B	1144	-	14,15,15	0.37	0	18,20,20	1.94	3 (16%)
4	EPE	C	1143	-	14,15,15	0.48	0	18,20,20	2.10	5 (27%)
5	PEG	C	1144	-	6,6,6	0.73	0	5,5,5	0.49	0
5	PEG	C	1145	-	6,6,6	0.61	0	5,5,5	0.35	0
5	PEG	C	1146	-	6,6,6	0.67	0	5,5,5	0.28	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EPE	B	1144	-	-	0/9/19/19	0/1/1/1
4	EPE	C	1143	-	-	0/9/19/19	0/1/1/1
5	PEG	C	1144	-	-	0/4/4/4	0/0/0/0
5	PEG	C	1145	-	-	0/4/4/4	0/0/0/0
5	PEG	C	1146	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	1143	EPE	C6-N1-C2	2.42	114.14	108.90
4	C	1143	EPE	C7-N4-C3	2.99	118.94	111.27
4	B	1144	EPE	C7-N4-C3	3.10	119.22	111.27

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
4	B	1144	EPE	C7-N4-C5	3.22	119.53	111.27
4	C	1143	EPE	C7-N4-C5	3.34	119.83	111.27
4	C	1143	EPE	C5-N4-C3	3.99	117.54	108.90
4	C	1143	EPE	O2S-S-C10	4.93	111.11	106.91
4	B	1144	EPE	C5-N4-C3	5.55	120.91	108.90

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	1144	EPE	6	0
4	C	1143	EPE	6	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	83/91 (91%)	-0.34	1 (1%) 81 81	24, 40, 65, 83	0
1	B	83/91 (91%)	-0.37	1 (1%) 81 81	23, 39, 59, 94	0
1	C	81/91 (89%)	-0.42	1 (1%) 81 81	23, 40, 57, 71	0
1	D	83/91 (91%)	-0.24	2 (2%) 62 61	22, 40, 64, 94	0
2	Q	11/11 (100%)	-0.19	0 100 100	79, 80, 83, 83	0
All	All	341/375 (90%)	-0.34	5 (1%) 76 75	22, 41, 79, 94	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	63	SER	2.7
1	C	61	ASP	2.5
1	D	61	ASP	2.5
1	B	60	ASN	2.3
1	A	60	ASN	2.2

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron

density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	NI	B	1143	1/1	1.00	0.11	0.13	29,29,29,29	0
3	NI	D	1143	1/1	1.00	0.10	0.06	31,31,31,31	0
3	NI	C	1142	1/1	0.99	0.08	-0.92	39,39,39,39	0
3	NI	A	1143	1/1	0.99	0.08	-1.14	41,41,41,41	0
5	PEG	C	1144	7/7	0.72	0.28	-	45,47,50,50	7
4	EPE	B	1144	15/15	0.80	0.20	-	73,80,84,84	15
5	PEG	C	1145	7/7	0.78	0.18	-	44,50,53,54	7
5	PEG	C	1146	7/7	0.69	0.28	-	35,43,50,51	7
4	EPE	C	1143	15/15	0.73	0.24	-	52,59,69,70	15

6.5 Other polymers

There are no such residues in this entry.