



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 06:51 AM GMT

PDB ID : 2Y69  
Title : Bovine heart cytochrome c oxidase re-refined with molecular oxygen  
Authors : Kaila, V.R.I.; Oksanen, E.; Goldman, A.; Verkhovsky, M.I.; Sundholm, D.; Wikstrom, M.  
Deposited on : 2011-01-20  
Resolution : 1.95 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.
























Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1833 (1.96-1.96)
Clashscore	102246	1953 (1.96-1.96)
Ramachandran outliers	100387	1936 (1.96-1.96)
Sidechain outliers	100360	1936 (1.96-1.96)
RSRZ outliers	91569	1835 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	514	<div> <div>3%</div> <div> <div></div> <div>86%</div> <div>13%</div> </div> </div>
1	N	514	<div> <div>3%</div> <div> <div></div> <div>82%</div> <div>18%</div> </div> </div>
2	B	227	<div> <div>2%</div> <div> <div></div> <div>79%</div> <div>18%</div> <div>•</div> </div> </div>
2	O	227	<div> <div>3%</div> <div> <div></div> <div>74%</div> <div>22%</div> <div>•</div> </div> </div>
3	C	261	<div> <div></div> <div> <div></div> <div>90%</div> <div>8%</div> <div>••</div> </div> </div>

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	P	261	
4	D	169	
4	Q	169	
5	E	152	
5	R	152	
6	F	129	
6	S	129	
7	G	97	
7	T	97	
8	H	86	
8	U	86	
9	I	74	
9	V	74	
10	J	80	
10	W	80	
11	K	80	
11	X	80	
12	L	63	
12	Y	63	
13	M	70	
13	Z	70	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
14	HEA	A	515	X	-	-	-
14	HEA	A	516	X	-	-	-

Continued on next page...

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
14	HEA	N	515	X	-	-	-
14	HEA	N	516	X	-	-	-
16	OXY	A	1515	-	-	-	X
23	DMU	M	1044	X	-	-	-
23	DMU	Z	1043	X	-	-	-

## 2 Entry composition

There are 24 unique types of molecules in this entry. The entry contains 30116 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CYTOCHROME C OXIDASE SUBUNIT 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	513	Total	C	N	O	S	0	0	0
			4017	2685	622	676	34			
1	N	513	Total	C	N	O	S	0	0	0
			4017	2685	622	676	34			

- Molecule 2 is a protein called CYTOCHROME C OXIDASE SUBUNIT 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	227	Total	C	N	O	S	0	0	0
			1822	1184	281	339	18			
2	O	226	Total	C	N	O	S	0	0	0
			1814	1179	280	338	17			

- Molecule 3 is a protein called CYTOCHROME C OXIDASE SUBUNIT 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	259	Total	C	N	O	S	0	0	0
			2110	1412	336	350	12			
3	P	259	Total	C	N	O	S	0	0	0
			2110	1412	336	350	12			

- Molecule 4 is a protein called CYTOCHROME C OXIDASE SUBUNIT 4 ISOFORM 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	144	Total	C	N	O	S	0	0	0
			1195	777	196	218	4			
4	Q	144	Total	C	N	O	S	0	0	0
			1195	777	196	218	4			

- Molecule 5 is a protein called CYTOCHROME C OXIDASE SUBUNIT 5A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	104	Total	C	N	O	S	0	0	0
			842	538	141	161	2			
5	R	104	Total	C	N	O	S	0	0	0
			842	538	141	161	2			

- Molecule 6 is a protein called CYTOCHROME C OXIDASE SUBUNIT 5B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	F	93	Total	C	N	O	S	0	0	0
			717	447	127	138	5			
6	S	93	Total	C	N	O	S	0	0	0
			717	447	127	138	5			

- Molecule 7 is a protein called CYTOCHROME C OXIDASE POLYPEPTIDE 6A2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
7	G	84	Total 671	C 428	N 128	O 113	P 1	S 1	0	0	0
7	T	84	Total 675	C 431	N 129	O 113	P 1	S 1	0	0	0

- Molecule 8 is a protein called CYTOCHROME C OXIDASE SUBUNIT VIB ISOFORM 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
8	H	75	Total	C	N	O	S	0	0	0
			628	395	114	114	5			
8	U	75	Total	C	N	O	S	0	0	0
			628	395	114	114	5			

- Molecule 9 is a protein called CYTOCHROME C OXIDASE POLYPEPTIDE VIC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
9	I	71	Total	C	N	O	S	0	0	0
			585	381	105	95	4			
9	V	71	Total	C	N	O	S	0	0	0
			585	381	105	95	4			

- Molecule 10 is a protein called CYTOCHROME C OXIDASE POLYPEPTIDE 7A1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	J	57	Total	C	N	O	S	0	0	0
			451	291	76	81	3			

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	W	57	Total	C	N	O	S	0	0	0
			451	291	76	81	3			

- Molecule 11 is a protein called CYTOCHROME C OXIDASE POLYPEPTIDE 7B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
11	K	49	Total	C	N	O	S	0	0	0
			384	250	65	67	2			
11	X	49	Total	C	N	O	S	0	0	0
			384	250	65	67	2			


- Molecule 12 is a protein called CYTOCHROME C OXIDASE SUBUNIT 7C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
12	L	46	Total	C	N	O	S	0	0	0
			380	254	64	60	2			
12	Y	46	Total	C	N	O	S	0	0	0
			380	254	64	60	2			

- Molecule 13 is a protein called CYTOCHROME C OXIDASE POLYPEPTIDE 8H.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
13	M	43	Total	C	N	O	0	0	0
			335	223	53	59			
13	Z	42	Total	C	N	O	0	0	0
			329	220	52	57			

- Molecule 14 is HEME-A (three-letter code: HEA) (formula: C<sub>49</sub>H<sub>56</sub>FeN<sub>4</sub>O<sub>6</sub>).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
14	A	1	Total	C	Fe	N	O	0	0
			60	49	1	4	6		
14	A	1	Total	C	Fe	N	O	0	0
			60	49	1	4	6		
14	N	1	Total	C	Fe	N	O	0	0
			60	49	1	4	6		
14	N	1	Total	C	Fe	N	O	0	0
			60	49	1	4	6		

- Molecule 15 is COPPER (II) ION (three-letter code: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
15	A	1	Total	Cu	0	0
			1	1		
15	N	1	Total	Cu	0	0
			1	1		

- Molecule 16 is OXYGEN MOLECULE (three-letter code: OXY) (formula: O<sub>2</sub>).






Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
16	A	1	Total O 2 2	0	0
16	N	1	Total O 2 2	0	0

- Molecule 17 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
17	A	1	Total Mg 1 1	0	0
17	N	1	Total Mg 1 1	0	0

- Molecule 18 is CHOLIC ACID (three-letter code: CHD) (formula: C<sub>24</sub>H<sub>40</sub>O<sub>5</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
18	A	1	Total 29	C 24	O 5	0	0
18	C	1	Total 29	C 24	O 5	0	0
18	G	1	Total 29	C 24	O 5	0	0
18	N	1	Total 29	C 24	O 5	0	0
18	P	1	Total 29	C 24	O 5	0	0
18	T	1	Total 29	C 24	O 5	0	0

- Molecule 19 is DINUCLEAR COPPER ION (three-letter code: CUA) (formula: Cu<sub>2</sub>).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
19	B	1	Total	Cu	0	0
			2	2		
19	O	1	Total	Cu	0	0
			2	2		

- Molecule 20 is (1S)-2-[[[(2-AMINOETHOXY)(HYDROXY)PHOSPHORYL]OXY}-1-[(STEAROYLOXY)METHYL]ETHYL (5E,8E,11E,14E)-ICOSA-5,8,11,14-TETRAENOATE (three-letter code: PEK) (formula: C<sub>43</sub>H<sub>78</sub>NO<sub>8</sub>P).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
20	C	1	Total	C	N	O	P	0	0
			53	43	1	8	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
20	P	1	Total	C	N	O	P	0	0
			53	43	1	8	1		

- Molecule 21 is (1R)-2-{{[(2S)-2,3-DIHYDROXYPROPYL]OXY}(HYDROXY)PHOSPHORYL]OXY}-1-[(PALMITOYLOXY)METHYL]ETHYL (11E)-OCTADEC-11-ENOATE (three-letter code: PGV) (formula: C<sub>40</sub>H<sub>77</sub>O<sub>10</sub>P).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
21	C	1	Total	C	O	P	0	0
			51	40	10	1		
21	C	1	Total	C	O	P	0	0
			51	40	10	1		
21	P	1	Total	C	O	P	0	0
			51	40	10	1		
21	P	1	Total	C	O	P	0	0
			51	40	10	1		

- Molecule 22 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
22	S	1	Total	Zn	0	0
			1	1		
22	F	1	Total	Zn	0	0
			1	1		

- Molecule 23 is SUGAR (DECYL-BETA-D-MALTOPYRANOSIDE) (three-letter code: DMU) (formula: C<sub>22</sub>H<sub>42</sub>O<sub>11</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
23	M	1	Total 33	C 22	O 11	0	0
23	Z	1	Total 33	C 22	O 11	0	0

- Molecule 24 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
24	A	114	Total O 114 114	0	0
24	B	102	Total O 102 102	0	0
24	C	85	Total O 85 85	0	0
24	D	43	Total O 43 43	0	0
24	F	45	Total O 45 45	0	0
24	G	41	Total O 41 41	0	0
24	H	49	Total O 49 49	0	0
24	I	20	Total O 20 20	0	0
24	J	26	Total O 26 26	0	0
24	K	10	Total O 10 10	0	0

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
24	L	11	Total 11	O 11	0	0
24	M	10	Total 10	O 10	0	0
24	N	118	Total 118	O 118	0	0
24	O	88	Total 88	O 88	0	0
24	P	17	Total 17	O 17	0	0
24	Q	54	Total 54	O 54	0	0
24	R	56	Total 56	O 56	0	0
24	S	46	Total 46	O 46	0	0
24	T	30	Total 30	O 30	0	0
24	U	32	Total 32	O 32	0	0
24	V	23	Total 23	O 23	0	0
24	X	12	Total 12	O 12	0	0
24	Y	6	Total 6	O 6	0	0
24	Z	10	Total 10	O 10	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: CYTOCHROME C OXIDASE SUBUNIT 1




#### • Molecule 1: CYTOCHROME C OXIDASE SUBUNIT 1




#### • Molecule 2: CYTOCHROME C OXIDASE SUBUNIT 2




#### • Molecule 2: CYTOCHROME C OXIDASE SUBUNIT 2




• Molecule 3: CYTOCHROME C OXIDASE SUBUNIT 3




• Molecule 3: CYTOCHROME C OXIDASE SUBUNIT 3



• Molecule 4: CYTOCHROME C OXIDASE SUBUNIT 4 ISOFORM 1




• Molecule 4: CYTOCHROME C OXIDASE SUBUNIT 4 ISOFORM 1




• Molecule 5: CYTOCHROME C OXIDASE SUBUNIT 5A









• Molecule 5: CYTOCHROME C OXIDASE SUBUNIT 5A




• Molecule 6: CYTOCHROME C OXIDASE SUBUNIT 5B



• Molecule 6: CYTOCHROME C OXIDASE SUBUNIT 5B




• Molecule 7: CYTOCHROME C OXIDASE POLYPEPTIDE 6A2




• Molecule 7: CYTOCHROME C OXIDASE POLYPEPTIDE 6A2








• Molecule 8: CYTOCHROME C OXIDASE SUBUNIT VIB ISOFORM 1




• Molecule 8: CYTOCHROME C OXIDASE SUBUNIT VIB ISOFORM 1




• Molecule 9: CYTOCHROME C OXIDASE POLYPEPTIDE VIC




• Molecule 9: CYTOCHROME C OXIDASE POLYPEPTIDE VIC



• Molecule 10: CYTOCHROME C OXIDASE POLYPEPTIDE 7A1



• Molecule 10: CYTOCHROME C OXIDASE POLYPEPTIDE 7A1






• Molecule 11: CYTOCHROME C OXIDASE POLYPEPTIDE 7B






• Molecule 11: CYTOCHROME C OXIDASE POLYPEPTIDE 7B



• Molecule 12: CYTOCHROME C OXIDASE SUBUNIT 7C



• Molecule 12: CYTOCHROME C OXIDASE SUBUNIT 7C



• Molecule 13: CYTOCHROME C OXIDASE POLYPEPTIDE 8H



• Molecule 13: CYTOCHROME C OXIDASE POLYPEPTIDE 8H



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	183.70Å 206.99Å 178.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	64.15 – 1.95 64.10 – 1.95	Depositor EDS
% Data completeness (in resolution range)	100.0 (64.15-1.95) 96.4 (64.10-1.95)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.54 (at 1.95Å)	Xtriage
Refinement program	REFMAC 5.5.0066	Depositor
R, $R_{free}$	0.211 , 0.245 0.210 , 0.240	Depositor DCC
$R_{free}$ test set	16433 reflections (3.61%)	DCC
Wilson B-factor (Å <sup>2</sup> )	34.0	Xtriage
Anisotropy	0.102	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 67.5	EDS
Estimated twinning fraction	0.007 for l,-k,h	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 471933 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	30116	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.25% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: MG, CHD, OXY, TPO, ZN, PGV, DMU, CUA, PEK, CU, HEA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.75	0/4156	0.72	1/5678 (0.0%)
1	N	0.60	0/4156	0.65	1/5678 (0.0%)
2	B	0.67	1/1868 (0.1%)	0.74	0/2544
2	O	0.50	0/1860	0.65	0/2534
3	C	0.59	0/2197	0.61	0/3005
3	P	0.65	3/2197 (0.1%)	0.63	1/3005 (0.0%)
4	D	0.61	0/1229	0.67	2/1658 (0.1%)
4	Q	0.39	0/1229	0.53	0/1658
5	E	0.56	0/860	0.62	0/1167
5	R	0.45	0/860	0.58	0/1167
6	F	0.58	0/733	0.71	0/996
6	S	0.53	0/733	0.66	0/996
7	G	0.48	0/686	0.60	0/933
7	T	0.46	0/690	0.63	0/937
8	H	0.52	0/648	0.57	0/877
8	U	0.44	0/648	0.57	0/877
9	I	0.53	0/598	0.57	0/792
9	V	0.43	0/598	0.50	0/792
10	J	0.44	0/462	0.54	0/625
10	W	0.42	0/462	0.56	0/625
11	K	0.55	0/398	0.59	0/546
11	X	0.37	0/398	0.50	0/546
12	L	0.60	0/393	0.59	0/526
12	Y	0.44	0/393	0.51	0/526
13	M	0.54	0/345	0.60	0/470
13	Z	0.40	0/339	0.52	0/462
All	All	0.58	4/29136 (0.0%)	0.64	5/39620 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms

of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
6	S	0	1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	P	236	GLU	CB-CG	9.64	1.70	1.52
3	P	236	GLU	CD-OE2	6.80	1.33	1.25
2	B	200	CYS	CB-SG	5.69	1.92	1.82
3	P	236	GLU	CG-CD	5.24	1.59	1.51

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	35	LEU	CA-CB-CG	-6.75	99.77	115.30
3	P	236	GLU	CB-CA-C	5.79	121.97	110.40
4	D	20	ARG	NE-CZ-NH1	5.65	123.13	120.30
1	N	278	MET	CG-SD-CE	-5.61	91.23	100.20
4	D	20	ARG	NE-CZ-NH2	-5.47	117.57	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
6	S	93	PRO	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4017	0	3990	78	0
1	N	4017	0	3990	91	0
2	B	1822	0	1834	51	0
2	O	1814	0	1822	48	0
3	C	2110	0	2027	19	0
3	P	2110	0	2027	36	0
4	D	1195	0	1183	19	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	Q	1195	0	1183	109	0
5	E	842	0	838	4	0
5	R	842	0	838	29	0
6	F	717	0	700	33	0
6	S	717	0	700	44	0
7	G	671	0	633	54	0
7	T	675	0	644	46	0
8	H	628	0	578	20	0
8	U	628	0	580	32	0
9	I	585	0	597	36	0
9	V	585	0	597	57	0
10	J	451	0	446	11	0
10	W	451	0	446	7	1
11	K	384	0	366	4	0
11	X	384	0	366	35	1
12	L	380	0	380	15	0
12	Y	380	0	380	33	0
13	M	335	0	351	19	0
13	Z	329	0	347	32	0
14	A	120	0	108	22	0
14	N	120	0	108	23	0
15	A	1	0	0	0	0
15	N	1	0	0	0	0
16	A	2	0	0	0	0
16	N	2	0	0	0	0
17	A	1	0	0	0	0
17	N	1	0	0	0	0
18	A	29	0	39	1	0
18	C	29	0	39	2	0
18	G	29	0	39	1	0
18	N	29	0	39	2	0
18	P	29	0	39	1	0
18	T	29	0	39	1	0
19	B	2	0	0	0	0
19	O	2	0	0	0	0
20	C	53	0	77	10	0
20	P	53	0	77	13	0
21	C	102	0	152	6	0
21	P	102	0	152	11	0
22	F	1	0	0	0	0
22	S	1	0	0	0	0
23	M	33	0	40	2	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
23	Z	33	0	40	8	0
24	A	114	0	0	11	0
24	B	102	0	0	12	0
24	C	85	0	0	2	0
24	D	43	0	0	4	0
24	F	45	0	0	15	0
24	G	41	0	0	7	0
24	H	49	0	0	11	0
24	I	20	0	0	3	0
24	J	26	0	0	5	0
24	K	10	0	0	0	0
24	L	11	0	0	1	0
24	M	10	0	0	1	0
24	N	118	0	0	1	0
24	O	88	0	0	3	0
24	P	17	0	0	0	0
24	Q	54	0	0	28	0
24	R	56	0	0	16	0
24	S	46	0	0	6	0
24	T	30	0	0	1	0
24	U	32	0	0	4	0
24	V	23	0	0	8	0
24	X	12	0	0	3	0
24	Y	6	0	0	5	0
24	Z	10	0	0	4	0
All	All	30116	0	28831	900	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 900 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
9:V:18:ARG:CB	9:V:18:ARG:HH11	1.09	1.59
13:M:42:LYS:CE	13:M:42:LYS:O	1.66	1.40
9:V:18:ARG:NH1	9:V:18:ARG:HB3	1.07	1.38
7:G:37:LEU:HD23	7:G:38:HIS:CE1	1.59	1.38
4:Q:4:SER:HB2	24:Q:2001:HOH:O	1.18	1.36

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
10:W:15:ASP:OD2	11:X:52:GLU:OE1[2_685]	2.16	0.04

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	511/514 (99%)	497 (97%)	13 (2%)	1 (0%)	52	43
1	N	511/514 (99%)	496 (97%)	15 (3%)	0	100	100
2	B	225/227 (99%)	213 (95%)	11 (5%)	1 (0%)	39	27
2	O	224/227 (99%)	209 (93%)	13 (6%)	2 (1%)	21	9
3	C	257/261 (98%)	253 (98%)	4 (2%)	0	100	100
3	P	257/261 (98%)	253 (98%)	4 (2%)	0	100	100
4	D	142/169 (84%)	138 (97%)	4 (3%)	0	100	100
4	Q	142/169 (84%)	126 (89%)	13 (9%)	3 (2%)	9	2
5	E	102/152 (67%)	101 (99%)	1 (1%)	0	100	100
5	R	102/152 (67%)	101 (99%)	1 (1%)	0	100	100
6	F	91/129 (70%)	87 (96%)	2 (2%)	2 (2%)	8	2
6	S	91/129 (70%)	87 (96%)	2 (2%)	2 (2%)	8	2
7	G	81/97 (84%)	69 (85%)	7 (9%)	5 (6%)	2	0
7	T	81/97 (84%)	65 (80%)	6 (7%)	10 (12%)	0	0
8	H	73/86 (85%)	68 (93%)	0	5 (7%)	1	0
8	U	73/86 (85%)	66 (90%)	4 (6%)	3 (4%)	3	0
9	I	69/74 (93%)	66 (96%)	3 (4%)	0	100	100
9	V	69/74 (93%)	64 (93%)	5 (7%)	0	100	100
10	J	55/80 (69%)	55 (100%)	0	0	100	100
10	W	55/80 (69%)	55 (100%)	0	0	100	100
11	K	47/80 (59%)	46 (98%)	1 (2%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
11	X	47/80 (59%)	42 (89%)	4 (8%)	1 (2%)	9	2
12	L	44/63 (70%)	41 (93%)	3 (7%)	0	100	100
12	Y	44/63 (70%)	42 (96%)	2 (4%)	0	100	100
13	M	41/70 (59%)	39 (95%)	1 (2%)	1 (2%)	7	1
13	Z	40/70 (57%)	35 (88%)	4 (10%)	1 (2%)	7	1
All	All	3474/4004 (87%)	3314 (95%)	123 (4%)	37 (1%)	17	6

5 of 37 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	59	GLN
6	F	95	GLN
7	G	3	ALA
7	G	8	HIS
8	H	43	MET

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	426/427 (100%)	418 (98%)	8 (2%)	65	58
1	N	426/427 (100%)	417 (98%)	9 (2%)	61	53
2	B	211/211 (100%)	200 (95%)	11 (5%)	29	13
2	O	210/211 (100%)	200 (95%)	10 (5%)	31	15
3	C	224/226 (99%)	219 (98%)	5 (2%)	60	51
3	P	224/226 (99%)	221 (99%)	3 (1%)	76	72
4	D	128/148 (86%)	127 (99%)	1 (1%)	86	85
4	Q	128/148 (86%)	124 (97%)	4 (3%)	47	34
5	E	91/123 (74%)	90 (99%)	1 (1%)	80	77
5	R	91/123 (74%)	89 (98%)	2 (2%)	60	51
6	F	79/103 (77%)	76 (96%)	3 (4%)	40	25

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
6	S	79/103 (77%)	74 (94%)	5 (6%)	22	8
7	G	66/78 (85%)	58 (88%)	8 (12%)	6	1
7	T	67/78 (86%)	60 (90%)	7 (10%)	9	2
8	H	67/76 (88%)	65 (97%)	2 (3%)	48	36
8	U	67/76 (88%)	62 (92%)	5 (8%)	17	5
9	I	56/59 (95%)	53 (95%)	3 (5%)	27	12
9	V	56/59 (95%)	52 (93%)	4 (7%)	18	6
10	J	48/68 (71%)	47 (98%)	1 (2%)	61	53
10	W	48/68 (71%)	47 (98%)	1 (2%)	61	53
11	K	39/66 (59%)	37 (95%)	2 (5%)	29	13
11	X	39/66 (59%)	37 (95%)	2 (5%)	29	13
12	L	39/55 (71%)	37 (95%)	2 (5%)	29	13
12	Y	39/55 (71%)	37 (95%)	2 (5%)	29	13
13	M	37/57 (65%)	33 (89%)	4 (11%)	8	2
13	Z	36/57 (63%)	34 (94%)	2 (6%)	26	11
All	All	3021/3394 (89%)	2914 (96%)	107 (4%)	43	29

5 of 107 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
13	M	34	LEU
1	N	485	VAL
9	V	41	GLU
13	M	39	ASN
1	N	138	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 42 such sidechains are listed below:

Mol	Chain	Res	Type
1	N	178	GLN
2	O	181	GLN
7	T	76	ASN
1	N	180	GLN
2	O	10	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
7	TPO	G	11	7	8,10,11	0.62	0	7,14,16	1.48	0
7	TPO	T	11	7	8,10,11	0.62	0	7,14,16	1.39	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	TPO	G	11	7	-	1/8/11/13	0/0/0/0
7	TPO	T	11	7	-	1/8/11/13	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	G	11	TPO	P-OG1-CB-CA
7	T	11	TPO	P-OG1-CB-CA

There are no ring outliers.

2 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	G	11	TPO	5	0
7	T	11	TPO	5	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 28 ligands modelled in this entry, 6 are monoatomic - leaving 22 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
16	OXY	A	1515	15,14	1,1,1	0.82	0	0,0,0	0.00	-
18	CHD	A	1517	-	29,32,32	0.75	1 (3%)	48,51,51	1.21	6 (12%)
14	HEA	A	515	1	40,67,67	1.14	4 (10%)	41,103,103	2.81	17 (41%)
14	HEA	A	516	1,16	40,67,67	1.12	6 (15%)	41,103,103	2.38	10 (24%)
19	CUA	B	228	2	0,1,1	0.00	-	0,0,0	0.00	-
20	PEK	C	1262	-	51,52,52	0.82	2 (3%)	52,57,57	1.02	3 (5%)
21	PGV	C	1263	-	50,50,50	0.87	2 (4%)	51,56,56	1.08	3 (5%)
21	PGV	C	1264	-	50,50,50	0.88	2 (4%)	51,56,56	0.96	4 (7%)
18	CHD	C	1265	-	29,32,32	0.71	0	48,51,51	1.24	6 (12%)
18	CHD	G	1085	-	29,32,32	0.70	0	48,51,51	1.24	5 (10%)
23	DMU	M	1044	-	34,34,34	1.34	1 (2%)	45,45,45	2.13	11 (24%)
16	OXY	N	1515	15,14	1,1,1	0.72	0	0,0,0	0.00	-
18	CHD	N	1517	-	29,32,32	0.69	1 (3%)	48,51,51	1.05	4 (8%)
14	HEA	N	515	1	40,67,67	1.13	4 (10%)	41,103,103	2.62	18 (43%)
14	HEA	N	516	1,16	40,67,67	1.00	3 (7%)	41,103,103	2.67	16 (39%)
19	CUA	O	228	2	0,1,1	0.00	-	0,0,0	0.00	-
20	PEK	P	1262	-	51,52,52	0.85	2 (3%)	52,57,57	1.07	4 (7%)
21	PGV	P	1263	-	50,50,50	0.87	2 (4%)	51,56,56	1.23	4 (7%)
21	PGV	P	1264	-	50,50,50	0.85	3 (6%)	51,56,56	0.98	4 (7%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
18	CHD	P	1265	-	29,32,32	0.69	0	48,51,51	1.37	11 (22%)
18	CHD	T	1085	-	29,32,32	0.61	0	48,51,51	1.41	6 (12%)
23	DMU	Z	1043	-	34,34,34	1.34	1 (2%)	45,45,45	1.90	7 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
16	OXY	A	1515	15,14	-	0/0/0/0	0/0/0/0
18	CHD	A	1517	-	-	0/7/74/74	0/4/4/4
14	HEA	A	515	1	3/3/7/16	0/24/76/76	0/0/8/8
14	HEA	A	516	1,16	3/3/7/16	0/24/76/76	0/0/8/8
19	CUA	B	228	2	-	0/0/0/0	0/0/0/0
20	PEK	C	1262	-	-	0/56/56/56	0/0/0/0
21	PGV	C	1263	-	-	0/55/55/55	0/0/0/0
21	PGV	C	1264	-	-	0/55/55/55	0/0/0/0
18	CHD	C	1265	-	-	0/7/74/74	0/4/4/4
18	CHD	G	1085	-	-	0/7/74/74	0/4/4/4
23	DMU	M	1044	-	4/4/10/10	0/19/59/59	0/2/2/2
16	OXY	N	1515	15,14	-	0/0/0/0	0/0/0/0
18	CHD	N	1517	-	-	0/7/74/74	0/4/4/4
14	HEA	N	515	1	3/3/7/16	1/24/76/76	0/0/8/8
14	HEA	N	516	1,16	3/3/7/16	0/24/76/76	0/0/8/8
19	CUA	O	228	2	-	0/0/0/0	0/0/0/0
20	PEK	P	1262	-	-	0/56/56/56	0/0/0/0
21	PGV	P	1263	-	-	0/55/55/55	0/0/0/0
21	PGV	P	1264	-	-	0/55/55/55	0/0/0/0
18	CHD	P	1265	-	-	0/7/74/74	0/4/4/4
18	CHD	T	1085	-	-	0/7/74/74	0/4/4/4
23	DMU	Z	1043	-	4/4/10/10	0/19/59/59	0/2/2/2

The worst 5 of 34 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
23	M	1044	DMU	C3-C4	-7.43	1.32	1.52
23	Z	1043	DMU	C3-C4	-7.40	1.32	1.52
18	A	1517	CHD	C10-C5	-2.41	1.51	1.55
21	P	1264	PGV	O01-C02	-2.07	1.41	1.46

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
18	N	1517	CHD	C13-C14	-2.06	1.51	1.55

The worst 5 of 139 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
14	N	515	HEA	C1A-C2A-C3A	-7.59	99.48	107.07
14	A	516	HEA	C1A-C2A-C3A	-7.49	99.57	107.07
14	A	515	HEA	C1A-C2A-C3A	-6.91	100.15	107.07
14	N	516	HEA	C1A-C2A-C3A	-6.48	100.59	107.07
14	A	515	HEA	CAA-CBA-CGA	-5.92	101.89	112.75

5 of 20 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
14	N	516	HEA	ND
14	N	516	HEA	NA
14	N	516	HEA	NB
23	M	1044	DMU	C2
23	M	1044	DMU	C3

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
14	N	515	HEA	C19-C18-C17-C16

There are no ring outliers.

18 monomers are involved in 103 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
18	A	1517	CHD	1	0
14	A	515	HEA	16	0
14	A	516	HEA	6	0
20	C	1262	PEK	10	0
21	C	1263	PGV	1	0
21	C	1264	PGV	5	0
18	C	1265	CHD	2	0
18	G	1085	CHD	1	0
23	M	1044	DMU	2	0
18	N	1517	CHD	2	0
14	N	515	HEA	17	0
14	N	516	HEA	6	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
20	P	1262	PEK	13	0
21	P	1263	PGV	4	0
21	P	1264	PGV	7	0
18	P	1265	CHD	1	0
18	T	1085	CHD	1	0
23	Z	1043	DMU	8	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	513/514 (99%)	0.46	13 (2%) 61 71	24, 29, 36, 56	0
1	N	513/514 (99%)	0.39	16 (3%) 52 62	27, 37, 50, 106	0
2	B	227/227 (100%)	0.11	4 (1%) 71 80	24, 35, 54, 67	0
2	O	226/227 (99%)	0.13	6 (2%) 58 68	32, 46, 82, 136	0
3	C	259/261 (99%)	-0.06	1 (0%) 93 95	27, 35, 50, 114	0
3	P	259/261 (99%)	-0.05	3 (1%) 81 87	29, 37, 55, 122	0
4	D	144/169 (85%)	-0.15	2 (1%) 78 85	30, 38, 64, 99	0
4	Q	144/169 (85%)	0.88	20 (13%) 4 6	40, 59, 91, 210	0
5	E	104/152 (68%)	0.28	8 (7%) 16 25	31, 39, 67, 110	0
5	R	104/152 (68%)	0.60	4 (3%) 44 56	36, 49, 79, 109	0
6	F	93/129 (72%)	0.42	4 (4%) 39 50	29, 40, 83, 243	0
6	S	93/129 (72%)	0.33	4 (4%) 39 50	32, 43, 89, 168	0
7	G	83/97 (85%)	0.95	15 (18%) 2 2	30, 42, 138, 172	0
7	T	83/97 (85%)	1.05	18 (21%) 1 1	30, 47, 146, 256	0
8	H	75/86 (87%)	0.73	12 (16%) 3 4	31, 44, 88, 166	0
8	U	75/86 (87%)	0.78	11 (14%) 3 5	38, 51, 99, 141	0
9	I	71/74 (95%)	0.89	9 (12%) 5 8	31, 47, 90, 126	0
9	V	71/74 (95%)	1.02	6 (8%) 13 21	37, 59, 90, 106	0
10	J	57/80 (71%)	0.55	7 (12%) 5 9	34, 44, 81, 95	0
10	W	57/80 (71%)	0.68	10 (17%) 2 3	38, 50, 100, 127	0
11	K	49/80 (61%)	0.06	0 100 100	33, 43, 70, 118	0
11	X	49/80 (61%)	1.36	15 (30%) 1 0	49, 58, 84, 125	0
12	L	46/63 (73%)	-0.07	1 (2%) 65 74	27, 37, 58, 137	0
12	Y	46/63 (73%)	0.15	2 (4%) 39 50	37, 50, 89, 114	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
13	M	43/70 (61%)	0.71	5 (11%) 6 10	28, 37, 122, 212	0
13	Z	42/70 (60%)	0.79	8 (19%) 2 2	44, 54, 120, 171	0
All	All	3526/4004 (88%)	0.38	204 (5%) 26 37	24, 39, 81, 256	0

The worst 5 of 204 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	Q	6	VAL	13.2
13	M	43	SER	12.6
6	F	95	GLN	11.1
6	F	96	LEU	11.0
4	Q	4	SER	9.9

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
7	TPO	G	11	11/12	0.58	0.34	-	73,140,486,493	0
7	TPO	T	11	11/12	0.68	0.39	-	65,143,410,489	0

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
16	OXY	A	1515	2/2	0.90	0.20	2.21	32,32,32,41	0
21	PGV	P	1264	51/51	0.95	0.14	1.95	28,43,86,192	0
23	DMU	Z	1043	33/33	0.84	0.20	1.58	42,72,127,190	0
23	DMU	M	1044	33/33	0.91	0.14	1.48	35,51,76,92	0
19	CUA	O	228	2/2	0.96	0.13	1.02	39,39,39,40	0
19	CUA	B	228	2/2	0.99	0.14	0.96	28,28,28,30	0
20	PEK	P	1262	53/53	0.94	0.14	0.92	35,52,149,483	0
20	PEK	C	1262	53/53	0.95	0.14	0.92	30,48,94,136	0
21	PGV	C	1263	51/51	0.97	0.14	0.91	26,46,86,132	0
14	HEA	N	516	60/60	0.97	0.17	0.70	21,31,40,48	0
18	CHD	A	1517	29/29	0.96	0.13	0.69	25,32,44,46	0
14	HEA	N	515	60/60	0.97	0.17	0.69	25,38,64,319	0
21	PGV	P	1263	51/51	0.97	0.12	0.46	30,48,80,146	0
18	CHD	C	1265	29/29	0.96	0.15	0.42	37,51,65,68	0
18	CHD	N	1517	29/29	0.97	0.13	0.37	27,37,42,44	0
21	PGV	C	1264	51/51	0.96	0.10	0.28	24,37,97,273	0
22	ZN	F	1097	1/1	1.00	0.10	0.25	35,35,35,35	0
14	HEA	A	515	60/60	0.98	0.16	0.13	19,26,49,53	0
14	HEA	A	516	60/60	0.98	0.13	-0.25	21,26,34,35	0
18	CHD	P	1265	29/29	0.94	0.15	-0.27	38,57,66,74	0
22	ZN	S	1097	1/1	0.99	0.09	-0.37	39,39,39,39	0
18	CHD	G	1085	29/29	0.98	0.09	-0.42	27,35,48,63	0
18	CHD	T	1085	29/29	0.97	0.08	-0.88	26,31,46,51	0
17	MG	N	1516	1/1	0.94	0.09	-1.01	35,35,35,35	0
17	MG	A	1516	1/1	0.99	0.08	-2.55	26,26,26,26	0
16	OXY	N	1515	2/2	0.95	0.09	-3.71	31,31,31,33	0
15	CU	N	517	1/1	0.99	0.16	-	32,32,32,32	0
15	CU	A	517	1/1	0.99	0.15	-	28,28,28,28	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.