



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 11:40 PM GMT

PDB ID : 1Y8O
Title : Crystal structure of the PDK3-L2 complex
Authors : Kato, M.; Chuang, J.L.; Wynn, R.M.; Chuang, D.T.
Deposited on : 2004-12-13
Resolution : 2.48 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.48 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4309 (2.50-2.46)
Clashscore	102246	5050 (2.50-2.46)
Ramachandran outliers	100387	4961 (2.50-2.46)
Sidechain outliers	100360	4963 (2.50-2.46)
RSRZ outliers	91569	4319 (2.50-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	419	
2	B	128	

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 3910 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called [Pyruvate dehydrogenase [lipoamide]] kinase isozyme 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	374	Total	C	N	O	S	0	1	0
			3066	1977	510	566	13			

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-12	GLY	-	CLONING ARTIFACT	UNP Q15120
A	-11	GLY	-	CLONING ARTIFACT	UNP Q15120
A	-10	SER	-	CLONING ARTIFACT	UNP Q15120
A	-9	HIS	-	EXPRESSION TAG	UNP Q15120
A	-8	HIS	-	EXPRESSION TAG	UNP Q15120
A	-7	HIS	-	EXPRESSION TAG	UNP Q15120
A	-6	HIS	-	EXPRESSION TAG	UNP Q15120
A	-5	HIS	-	EXPRESSION TAG	UNP Q15120
A	-4	HIS	-	EXPRESSION TAG	UNP Q15120
A	-3	GLY	-	CLONING ARTIFACT	UNP Q15120
A	-2	MET	-	CLONING ARTIFACT	UNP Q15120
A	-1	ALA	-	CLONING ARTIFACT	UNP Q15120
A	0	ARG	-	CLONING ARTIFACT	UNP Q15120
A	1	LEU	-	CLONING ARTIFACT	UNP Q15120
A	2	GLU	-	CLONING ARTIFACT	UNP Q15120
A	3	ASN	-	CLONING ARTIFACT	UNP Q15120
A	4	LEU	-	CLONING ARTIFACT	UNP Q15120
A	5	TYR	-	CLONING ARTIFACT	UNP Q15120
A	6	PHE	-	CLONING ARTIFACT	UNP Q15120
A	7	GLN	-	CLONING ARTIFACT	UNP Q15120
A	8	GLY	-	CLONING ARTIFACT	UNP Q15120

- Molecule 2 is a protein called Dihydrolipoyllysine-residue acetyltransferase component of pyruvate dehydrogenase complex.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	97	Total	C	N	O	S	0	0	0
			736	471	115	146	4			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	106	GLY	-	CLONING ARTIFACT	UNP P10515
B	107	GLY	-	CLONING ARTIFACT	UNP P10515
B	108	SER	-	CLONING ARTIFACT	UNP P10515
B	109	HIS	-	EXPRESSION TAG	UNP P10515
B	110	HIS	-	EXPRESSION TAG	UNP P10515
B	111	HIS	-	EXPRESSION TAG	UNP P10515
B	112	HIS	-	EXPRESSION TAG	UNP P10515
B	113	HIS	-	EXPRESSION TAG	UNP P10515
B	114	HIS	-	EXPRESSION TAG	UNP P10515
B	115	GLY	-	CLONING ARTIFACT	UNP P10515
B	116	MET	-	CLONING ARTIFACT	UNP P10515
B	117	ALA	-	CLONING ARTIFACT	UNP P10515
B	118	ARG	-	CLONING ARTIFACT	UNP P10515
B	119	LEU	-	CLONING ARTIFACT	UNP P10515
B	120	GLU	-	CLONING ARTIFACT	UNP P10515
B	121	ASN	-	CLONING ARTIFACT	UNP P10515
B	122	LEU	-	CLONING ARTIFACT	UNP P10515
B	123	TYR	-	CLONING ARTIFACT	UNP P10515
B	124	PHE	-	CLONING ARTIFACT	UNP P10515
B	125	GLN	-	CLONING ARTIFACT	UNP P10515


- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		

- Molecule 4 is POTASSIUM ION (three-letter code: K) (formula: K).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	K	0	0
			2	2		

- Molecule 5 is DIHYDROLIPOIC ACID (three-letter code: RED) (formula: C₈H₁₆O₂S₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	O	S	0	0
			11	8	1	2		

- Molecule 6 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $\text{C}_{10}\text{H}_{15}\text{N}_5\text{O}_{10}\text{P}_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	63	Total 63	O 63	0	0
7	B	4	Total 4	O 4	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	120.75Å 120.75Å 239.15Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 2.48 48.13 – 2.48	Depositor EDS
% Data completeness (in resolution range)	99.7 (50.00-2.48) 99.7 (48.13-2.48)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.95 (at 2.48Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.210 , 0.234 0.263 , 0.279	Depositor DCC
R_{free} test set	1492 reflections (4.18%)	DCC
Wilson B-factor (Å ²)	64.3	Xtriage
Anisotropy	0.027	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 43.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	1 of 37232 reflections (0.003%)	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3910	wwPDB-VP
Average B, all atoms (Å ²)	64.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.03% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, K, ADP, RED

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.84	3/3145 (0.1%)	0.85	10/4257 (0.2%)
2	B	1.27	5/749 (0.7%)	0.83	1/1017 (0.1%)
All	All	0.94	8/3894 (0.2%)	0.85	11/5274 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	216	ALA	C-N	13.97	1.66	1.34
2	B	215	SER	CB-OG	13.31	1.59	1.42
2	B	220	TYR	C-O	11.16	1.44	1.23
2	B	216	ALA	C-O	-10.40	1.03	1.23
2	B	211	GLU	CD-OE1	6.59	1.32	1.25
1	A	18	ARG	CZ-NH1	6.27	1.41	1.33
1	A	101	ASP	CG-OD1	5.16	1.37	1.25
1	A	102	PRO	C-O	5.08	1.33	1.23

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	336	ARG	NE-CZ-NH1	9.49	125.05	120.30
1	A	336	ARG	NE-CZ-NH2	-8.21	116.20	120.30
1	A	388	ASP	N-CA-CB	-7.08	97.86	110.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	18	ARG	NE-CZ-NH2	-6.25	117.18	120.30
1	A	199	ASP	CB-CG-OD1	6.09	123.78	118.30
1	A	101	ASP	CB-CG-OD2	-5.95	112.94	118.30
1	A	199	ASP	CB-CG-OD2	-5.76	113.11	118.30
1	A	332	ARG	NE-CZ-NH2	-5.61	117.49	120.30
1	A	387	ASP	C-N-CA	5.44	135.30	121.70
1	A	297	ILE	CG1-CB-CG2	-5.44	99.44	111.40
2	B	172	ASP	N-CA-CB	-5.38	100.91	110.60

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	171	THR	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3066	0	3029	39	0
2	B	736	0	727	15	0
3	A	1	0	0	0	0
4	A	2	0	0	0	0
5	B	11	0	15	3	0
6	A	27	0	12	0	0
7	A	63	0	0	3	0
7	B	4	0	0	0	0
All	All	3910	0	3783	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:173:LYS:HZ3	5:B:373:RED:C1	1.14	1.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:173:LYS:NZ	5:B:373:RED:C1	1.78	1.43
1:A:247:GLU:HG3	7:A:560:HOH:O	1.35	1.21
2:B:173:LYS:HZ1	5:B:373:RED:C1	1.82	0.88
1:A:28:SER:H	1:A:31:GLN:HE21	1.26	0.82
2:B:208:VAL:HG11	2:B:213:ASP:HB2	1.65	0.76
2:B:187:ALA:HB2	2:B:208:VAL:HG23	1.74	0.69
1:A:199:ASP:O	1:A:203:THR:HG23	1.95	0.66
1:A:332:ARG:HD2	7:A:540:HOH:O	1.97	0.64
1:A:320:PRO:O	1:A:321:LEU:HB2	2.03	0.58
1:A:163:MET:HE2	7:A:538:HOH:O	2.03	0.58
2:B:208:VAL:HG11	2:B:213:ASP:CB	2.34	0.57
1:A:276:LEU:HD13	1:A:281:LEU:HD23	1.86	0.57
1:A:69:ASP:OD2	1:A:69:ASP:N	2.32	0.57
1:A:99:PRO:C	1:A:101:ASP:H	2.08	0.56
1:A:57:LEU:O	1:A:61:MET:HG3	2.06	0.56
2:B:206:ILE:HD11	2:B:220:TYR:CE2	2.41	0.55
1:A:67:LEU:HG	1:A:71:LEU:HD23	1.88	0.54
1:A:378:ARG:O	1:A:382:THR:HG23	2.09	0.53
1:A:332:ARG:O	1:A:336:ARG:HG2	2.09	0.52
1:A:87:SER:OG	1:A:119:HIS:HE1	1.93	0.52
1:A:158:ARG:HH12	1:A:162:ARG:NH1	2.08	0.51
1:A:374:LYS:HG3	2:B:163:GLY:HA3	1.93	0.51
1:A:158:ARG:HH22	1:A:162:ARG:NH1	2.09	0.51
1:A:123:VAL:HB	1:A:124:PRO:HD3	1.92	0.50
1:A:293:PRO:HG2	1:A:296:LYS:HD3	1.94	0.50
2:B:151:ARG:HD3	2:B:153:GLU:OE2	2.11	0.50
1:A:74:ARG:HD3	1:A:132:GLU:HB3	1.95	0.49
1:A:16:ILE:HD11	1:A:85:MET:SD	2.55	0.47
1:A:57:LEU:HG	1:A:161:PHE:CE2	2.51	0.46
1:A:64:VAL:HA	1:A:67:LEU:HD22	1.98	0.46
1:A:260:TYR:HA	1:A:263:ARG:HG3	1.97	0.46
1:A:51:LYS:O	1:A:54:PRO:HD2	2.16	0.46
2:B:217:PHE:C	2:B:219:ASP:H	2.17	0.46
1:A:98:SER:HA	1:A:99:PRO:HD2	1.84	0.45
1:A:343:LYS:HD3	1:A:345[A]:TYR:OH	2.17	0.45
1:A:109:LEU:HD21	1:A:173:GLY:HA2	1.98	0.44
2:B:208:VAL:HG12	2:B:209:GLU:N	2.33	0.44
1:A:263:ARG:HB3	1:A:265:GLU:HG2	1.99	0.44
1:A:87:SER:OG	1:A:119:HIS:CE1	2.71	0.44
2:B:209:GLU:OE1	2:B:210:LYS:HE3	2.18	0.44
1:A:140:ASP:HA	1:A:141:PRO:HD2	1.64	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:173:LYS:HB3	2:B:173:LYS:HE3	1.77	0.42
2:B:152:TRP:HD1	2:B:193:GLU:HG3	1.83	0.42
1:A:101:ASP:HA	1:A:102:PRO:HD2	1.96	0.42
1:A:336:ARG:HD3	1:A:341:ASP:OD1	2.20	0.41
1:A:61:MET:HA	1:A:64:VAL:HG13	2.02	0.41
1:A:259:LEU:HD23	1:A:260:TYR:CE2	2.55	0.41
1:A:383:THR:OG1	1:A:384:PRO:HD2	2.21	0.41
1:A:305:TYR:O	1:A:306:SER:HB2	2.20	0.40
1:A:111:VAL:O	1:A:115:VAL:HG13	2.21	0.40
1:A:377:TRP:CZ2	1:A:381:LYS:HG3	2.56	0.40
2:B:190:LEU:HD23	2:B:203:PRO:HB2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	369/419 (88%)	361 (98%)	6 (2%)	2 (0%)	34	53
2	B	93/128 (73%)	91 (98%)	1 (1%)	1 (1%)	17	29
All	All	462/547 (84%)	452 (98%)	7 (2%)	3 (1%)	26	48

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	172	ASP
1	A	100	GLU
1	A	141	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	341/374 (91%)	315 (92%)	26 (8%)	16	29
2	B	79/109 (72%)	73 (92%)	6 (8%)	16	29
All	All	420/483 (87%)	388 (92%)	32 (8%)	16	29

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	13	PRO
1	A	23	SER
1	A	27	LEU
1	A	39	ASN
1	A	64	VAL
1	A	66	LEU
1	A	67	LEU
1	A	69	ASP
1	A	72	LEU
1	A	86	GLN
1	A	105	LEU
1	A	116	ARG
1	A	119	HIS
1	A	128	GLN
1	A	158	ARG
1	A	162	ARG
1	A	177	ASN
1	A	203	THR
1	A	216	PRO
1	A	265	GLU
1	A	328	LEU
1	A	332	ARG
1	A	336	ARG
1	A	346	SER
1	A	400	SER
1	A	401	LYS
2	B	128	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	133	MET
2	B	140	LEU
2	B	141	SER
2	B	186	LEU
2	B	190	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	31	GLN
1	A	39	ASN
1	A	70	ASN
1	A	73	ASN
1	A	86	GLN
1	A	119	HIS
1	A	128	GLN
1	A	177	ASN
1	A	240	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length

(or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	ADP	A	504	3,4	22,29,29	1.36	2 (9%)	27,45,45	1.74	5 (18%)
5	RED	B	373	-	8,10,11	0.41	0	4,10,12	1.48	1 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	ADP	A	504	3,4	-	0/12/32/32	0/3/3/3
5	RED	B	373	-	-	0/6/9/10	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	504	ADP	C5-C4	3.67	1.48	1.40
6	A	504	ADP	O4'-C1'	3.94	1.46	1.41

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	504	ADP	N3-C2-N1	-6.42	123.98	128.89
6	A	504	ADP	C4-C5-N7	-2.66	107.03	109.48
6	A	504	ADP	PA-O3A-PB	-2.10	125.61	132.67
5	B	373	RED	C3-C2-C1	-2.06	107.46	113.36
6	A	504	ADP	O2B-PB-O1B	2.29	117.97	110.58
6	A	504	ADP	C2-N1-C6	2.37	123.00	118.77

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	373	RED	3	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	374/419 (89%)	0.74	40 (10%) 8 8	46, 62, 81, 121	0
2	B	97/128 (75%)	0.93	18 (18%) 2 1	53, 63, 86, 121	0
All	All	471/547 (86%)	0.78	58 (12%) 5 5	46, 62, 82, 121	0

All (58) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	176	THR	7.6
1	A	139	PHE	7.0
2	B	220	TYR	5.7
1	A	179	VAL	5.2
1	A	304	MET	5.1
1	A	38	ASP	4.9
1	A	399	ALA	4.6
1	A	177	ASN	4.4
1	A	161	PHE	4.4
1	A	400	SER	4.3
1	A	174	GLY	4.2
2	B	187	ALA	4.2
1	A	401	LYS	4.1
2	B	218	ALA	4.0
1	A	175	ASP	3.8
1	A	324	PHE	3.8
2	B	212	ALA	3.8
2	B	219	ASP	3.8
1	A	142	PHE	3.8
1	A	305	TYR	3.7
2	B	129	TYR	3.7
2	B	226	THR	3.6
1	A	180	HIS	3.4
2	B	217	PHE	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	209	GLU	2.9
1	A	136	LYS	2.9
2	B	214	ILE	2.8
1	A	158	ARG	2.7
2	B	131	PRO	2.6
1	A	104	VAL	2.6
2	B	128	SER	2.6
1	A	156	THR	2.6
1	A	137	PHE	2.4
1	A	103	GLN	2.4
1	A	226	LYS	2.4
1	A	230	LYS	2.4
1	A	370	PRO	2.4
2	B	213	ASP	2.4
1	A	178	PRO	2.3
1	A	40	ALA	2.3
1	A	60	THR	2.3
1	A	182	LYS	2.3
1	A	99	PRO	2.3
2	B	211	GLU	2.2
2	B	192	PRO	2.2
1	A	159	ILE	2.2
1	A	398	ASP	2.2
2	B	155	LYS	2.2
1	A	333	LEU	2.2
1	A	157	ASN	2.2
2	B	210	LYS	2.1
2	B	215	SER	2.1
1	A	345[A]	TYR	2.1
1	A	43	LYS	2.1
1	A	397	ARG	2.1
1	A	389	TRP	2.0
1	A	151	LEU	2.0
1	A	321	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	RED	B	373	11/12	0.93	0.19	-0.06	46,50,57,62	0
6	ADP	A	504	27/27	0.89	0.17	-0.67	59,76,84,85	0
3	MG	A	501	1/1	0.76	0.18	-1.56	96,96,96,96	0
4	K	A	502	1/1	0.90	0.08	-5.38	88,88,88,88	0
4	K	A	503	1/1	0.82	0.11	-5.51	65,65,65,65	0

6.5 Other polymers [i](#)

There are no such residues in this entry.