



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 06:45 AM GMT

PDB ID : 2Y8U
Title : A. nidulans chitin deacetylase
Authors : Penman, G.; Gay, L.M.; Van Aalten, D.M.F.
Deposited on : 2011-02-10
Resolution : 1.99 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	230	 77% 13% 8%
1	B	230	 79% 13% 8%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PO4	A	1238	-	-	-	X
2	PO4	B	1238	-	-	-	X
3	CO	B	1239	-	-	-	X
4	NA	A	1240	-	-	-	X
5	CL	A	1242	-	-	-	X
5	CL	B	1240	-	-	-	X

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 3376 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CHITIN DEACETYLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	211	Total	C	N	O	S	0	0	0
			1619	1022	276	314	7			
1	B	212	Total	C	N	O	S	0	0	0
			1622	1023	276	316	7			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	8	MET	-	EXPRESSION TAG	UNP Q5AQQ0
A	9	ALA	-	EXPRESSION TAG	UNP Q5AQQ0
A	10	HIS	-	EXPRESSION TAG	UNP Q5AQQ0
A	11	HIS	-	EXPRESSION TAG	UNP Q5AQQ0
A	12	HIS	-	EXPRESSION TAG	UNP Q5AQQ0
A	13	HIS	-	EXPRESSION TAG	UNP Q5AQQ0
A	14	HIS	-	EXPRESSION TAG	UNP Q5AQQ0
A	15	HIS	-	EXPRESSION TAG	UNP Q5AQQ0
A	16	HIS	-	EXPRESSION TAG	UNP Q5AQQ0
A	17	ARG	-	EXPRESSION TAG	UNP Q5AQQ0
A	18	SER	-	EXPRESSION TAG	UNP Q5AQQ0
B	8	MET	-	EXPRESSION TAG	UNP Q5AQQ0
B	9	ALA	-	EXPRESSION TAG	UNP Q5AQQ0
B	10	HIS	-	EXPRESSION TAG	UNP Q5AQQ0
B	11	HIS	-	EXPRESSION TAG	UNP Q5AQQ0
B	12	HIS	-	EXPRESSION TAG	UNP Q5AQQ0
B	13	HIS	-	EXPRESSION TAG	UNP Q5AQQ0
B	14	HIS	-	EXPRESSION TAG	UNP Q5AQQ0
B	15	HIS	-	EXPRESSION TAG	UNP Q5AQQ0
B	16	HIS	-	EXPRESSION TAG	UNP Q5AQQ0
B	17	ARG	-	EXPRESSION TAG	UNP Q5AQQ0
B	18	SER	-	EXPRESSION TAG	UNP Q5AQQ0

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is COBALT (II) ION (three-letter code: CO) (formula: Co).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Co	0	0
			1	1		
3	A	1	Total	Co	0	0
			1	1		

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	3	Total	Na	0	0
			3	3		

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Cl	0	0
			1	1		
5	A	1	Total	Cl	0	0
			1	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	60	Total 60	O 60	0	0
6	B	58	Total 58	O 58	0	0

- Molecule 1: CHITIN DEACETYLASE

• Molecule 1: CHITIN DEACETYLASE

E208	R209	Q212	G218	C227	D230	I233	H237																												
MET	ALA	HIS	HIS	HIS	HIS	HIS	HIS	HIS	ARG	R26	C36	D48	R67	V72	L83	M87	R88	D100	H101	S102	S103	L104	P105	A113	T117	R118	V161	D162	N168	Q169	D170	A173	Q185	H196	H199

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	35.64Å 64.52Å 86.54Å 90.00° 101.16° 90.00°	Depositor
Resolution (Å)	84.90 – 1.99 51.37 – 1.99	Depositor EDS
% Data completeness (in resolution range)	99.5 (84.90-1.99) 99.5 (51.37-1.99)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.13 (at 1.98Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.157 , 0.205 0.163 , 0.210	Depositor DCC
R_{free} test set	1335 reflections (5.33%)	DCC
Wilson B-factor (Å ²)	20.3	Xtriage
Anisotropy	0.791	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 49.5	EDS
Estimated twinning fraction	0.037 for h,-k,-h-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 26418 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3376	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, PO4, CO, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.13	2/1653 (0.1%)	1.02	5/2259 (0.2%)
1	B	1.07	1/1656 (0.1%)	0.99	3/2264 (0.1%)
All	All	1.10	3/3309 (0.1%)	1.00	8/4523 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	218	GLY	N-CA	5.38	1.54	1.46
1	B	72	VAL	CB-CG2	5.06	1.63	1.52
1	A	63	ARG	CB-CG	-5.00	1.39	1.52

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	217	ARG	C-N-CA	-9.98	101.35	122.30
1	B	48	ASP	CB-CG-OD1	8.03	125.53	118.30
1	A	197	ASP	CB-CG-OD1	6.53	124.17	118.30
1	B	209	ARG	NE-CZ-NH2	-6.35	117.13	120.30
1	A	184	ASP	CB-CG-OD1	5.76	123.48	118.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	218	GLY	Peptide
1	B	218	GLY	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1619	0	1563	23	0
1	B	1622	0	1559	19	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	3	0	0	1	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	60	0	0	5	0
6	B	58	0	0	2	0
All	All	3376	0	3122	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 40 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:208:GLU:O	1:B:212:GLN:HG2	1.67	0.94
1:B:162:ASP:H	1:B:185:GLN:HE22	1.23	0.82
4:A:1241:NA:NA	4:A:1244:NA:NA	1.20	0.69
1:B:162:ASP:H	1:B:185:GLN:NE2	1.90	0.68
1:B:161:VAL:HA	1:B:185:GLN:HE21	1.60	0.66

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	209/230 (91%)	202 (97%)	7 (3%)	0	100	100
1	B	210/230 (91%)	201 (96%)	9 (4%)	0	100	100
All	All	419/460 (91%)	403 (96%)	16 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	170/193 (88%)	167 (98%)	3 (2%)	66	69
1	B	170/193 (88%)	168 (99%)	2 (1%)	78	81
All	All	340/386 (88%)	335 (98%)	5 (2%)	72	75

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	63	ARG
1	A	170	ASP
1	A	230	ASP
1	B	230	ASP
1	B	237	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	199	HIS
1	B	40	ASN
1	B	168	ASN
1	A	169	GLN
1	B	147	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 7 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PO4	A	1238	3	4,4,4	1.21	0	6,6,6	0.34	0
2	PO4	B	1238	3	4,4,4	1.02	0	6,6,6	0.33	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PO4	A	1238	3	-	0/0/0/0	0/0/0/0
2	PO4	B	1238	3	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	211/230 (91%)	-0.38	0 100 100	12, 20, 30, 40	0
1	B	212/230 (92%)	-0.33	0 100 100	13, 23, 33, 42	0
All	All	423/460 (91%)	-0.35	0 100 100	12, 21, 33, 42	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	CL	A	1242	1/1	0.79	0.48	14.41	86,86,86,86	0
4	NA	A	1240	1/1	0.89	0.25	6.91	40,40,40,40	0
2	PO4	A	1238	5/5	0.98	0.12	4.02	35,39,42,45	0
2	PO4	B	1238	5/5	0.99	0.12	3.56	36,38,41,42	0
3	CO	B	1239	1/1	1.00	0.11	3.36	19,19,19,19	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	CL	B	1240	1/1	0.89	0.15	2.08	57,57,57,57	0
3	CO	A	1239	1/1	1.00	0.10	1.55	21,21,21,21	0
4	NA	A	1244	1/1	0.98	0.06	-2.32	43,43,43,43	0
4	NA	A	1241	1/1	0.98	0.06	-	44,44,44,44	0

6.5 Other polymers [i](#)

There are no such residues in this entry.