



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 06:46 AM GMT

PDB ID : 2YB7  
Title : CBM62 in complex with 6-alpha-D-Galactosyl-mannotriose  
Authors : Montanier, C.Y.; Correia, M.A.S.; Flint, J.E.; Zhu, Y.; Mckee, L.S.; Prates, J.A.M.; Polizzi, S.J.; Coutinho, P.M.; Henrissat, B.; Fontes, C.M.G.A.; Gilbert, H.J.  
Deposited on : 2011-03-02  
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3190 (1.70-1.70)
Clashscore	102246	3585 (1.70-1.70)
Ramachandran outliers	100387	3527 (1.70-1.70)
Sidechain outliers	100360	3527 (1.70-1.70)
RSRZ outliers	91569	3200 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	155	 2% 77% 12% • 10%
1	B	155	 2% 79% 10% • 10%

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 2703 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CARBOHYDRATE BINDING FAMILY 6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	140	Total	C	N	O	S	0	2	0
			1073	678	182	210	3			
1	B	140	Total	C	N	O	S	0	2	0
			1076	680	183	210	3			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	737	MET	-	EXPRESSION TAG	UNP D1NNT8
A	738	ALA	-	EXPRESSION TAG	UNP D1NNT8
A	739	SER	-	EXPRESSION TAG	UNP D1NNT8
A	884	LEU	-	EXPRESSION TAG	UNP D1NNT8
A	885	GLU	-	EXPRESSION TAG	UNP D1NNT8
A	886	HIS	-	EXPRESSION TAG	UNP D1NNT8
A	887	HIS	-	EXPRESSION TAG	UNP D1NNT8
A	888	HIS	-	EXPRESSION TAG	UNP D1NNT8
A	889	HIS	-	EXPRESSION TAG	UNP D1NNT8
A	890	HIS	-	EXPRESSION TAG	UNP D1NNT8
A	891	HIS	-	EXPRESSION TAG	UNP D1NNT8
B	737	MET	-	EXPRESSION TAG	UNP D1NNT8
B	738	ALA	-	EXPRESSION TAG	UNP D1NNT8
B	739	SER	-	EXPRESSION TAG	UNP D1NNT8
B	884	LEU	-	EXPRESSION TAG	UNP D1NNT8
B	885	GLU	-	EXPRESSION TAG	UNP D1NNT8
B	886	HIS	-	EXPRESSION TAG	UNP D1NNT8
B	887	HIS	-	EXPRESSION TAG	UNP D1NNT8
B	888	HIS	-	EXPRESSION TAG	UNP D1NNT8
B	889	HIS	-	EXPRESSION TAG	UNP D1NNT8
B	890	HIS	-	EXPRESSION TAG	UNP D1NNT8
B	891	HIS	-	EXPRESSION TAG	UNP D1NNT8

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Ca	0	0
			1	1		
3	A	1	Total	Ca	0	0
			1	1		

- Molecule 4 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	2	Total	C	O	0	0
			23	12	11		

- Molecule 5 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	2	Total	C	O	0	0
			23	12	11		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	251	Total 251	O 251	0	0
6	B	243	Total 243	O 243	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: CARBOHYDRATE BINDING FAMILY 6



#### • Molecule 1: CARBOHYDRATE BINDING FAMILY 6



## 4 Data and refinement statistics

Property	Value	Source
Space group	F 2 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	191.35Å 191.35Å 191.35Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	110.43 – 1.70 39.06 – 1.70	Depositor EDS
% Data completeness (in resolution range)	100.0 (110.43-1.70) 100.0 (39.06-1.70)	Depositor EDS
$R_{merge}$	0.15	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.74 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.165 , 0.186 0.164 , 0.184	Depositor DCC
$R_{free}$ test set	3220 reflections (5.34%)	DCC
Wilson B-factor (Å <sup>2</sup> )	17.2	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 42.1	EDS
Estimated twinning fraction	0.478 for k,h,-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 63488 reflections	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	2703	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.28% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CA, BMA, GLA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.53	10/1102 (0.9%)	1.31	5/1496 (0.3%)
1	B	1.50	4/1105 (0.4%)	1.32	2/1500 (0.1%)
All	All	1.52	14/2207 (0.6%)	1.32	7/2996 (0.2%)

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	754	TRP	CE3-CZ3	7.12	1.50	1.38
1	B	754	TRP	CE3-CZ3	6.46	1.49	1.38
1	B	767	GLY	N-CA	5.95	1.54	1.46
1	A	852	PHE	CE2-CZ	5.79	1.48	1.37
1	A	767	GLY	N-CA	5.75	1.54	1.46
1	A	797	GLN	CB-CG	-5.67	1.37	1.52
1	A	808	GLN	CG-CD	5.55	1.63	1.51
1	A	808	GLN	CD-OE1	5.35	1.35	1.24
1	B	797	GLN	CG-CD	5.24	1.63	1.51
1	A	868	ALA	CA-CB	5.05	1.63	1.52
1	A	782	TRP	CE3-CZ3	5.05	1.47	1.38
1	B	834	SER	CB-OG	-5.04	1.35	1.42
1	A	773	PHE	CD2-CE2	5.04	1.49	1.39
1	A	834	SER	CB-OG	-5.01	1.35	1.42

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	845	ASP	CB-CG-OD2	-7.45	111.59	118.30
1	A	845	ASP	CB-CG-OD1	5.91	123.62	118.30
1	B	872	PHE	CB-CG-CD1	-5.73	116.79	120.80
1	A	834	SER	CB-CA-C	-5.46	99.72	110.10
1	B	822	ASP	CB-CG-OD1	5.23	123.00	118.30

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	766	ASP	CB-CG-OD1	5.22	123.00	118.30
1	A	765	PHE	CB-CG-CD2	-5.06	117.26	120.80

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1073	0	1019	10	0
1	B	1076	0	1022	13	0
2	A	6	0	8	0	0
2	B	6	0	8	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	23	0	21	0	0
5	B	23	0	21	1	0
6	A	251	0	0	4	0
6	B	243	0	0	5	1
All	All	2703	0	2099	24	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:792[A]:ARG:NH2	6:A:2102:HOH:O	2.20	0.74
1:B:848:ASN:HD22	1:B:850:THR:H	1.39	0.71
1:A:848:ASN:HD22	1:A:850:THR:H	1.42	0.67
1:B:808[B]:GLN:H	1:B:808[B]:GLN:NE2	1.98	0.61
1:A:792[A]:ARG:NH2	6:A:2098:HOH:O	2.34	0.61
5:B:1881:BMA:O4	6:B:2234:HOH:O	2.16	0.60
1:B:848:ASN:ND2	1:B:850:THR:H	1.99	0.60
1:A:792[A]:ARG:CZ	6:A:2102:HOH:O	2.51	0.58

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:848:ASN:ND2	1:A:850:THR:H	2.00	0.58
1:A:848:ASN:C	1:A:848:ASN:HD22	2.09	0.56
1:B:792[A]:ARG:NE	6:B:2090:HOH:O	2.39	0.56
1:A:792[A]:ARG:NE	6:A:2102:HOH:O	2.38	0.55
1:B:792[A]:ARG:NH2	6:B:2090:HOH:O	2.39	0.54
1:B:792[A]:ARG:NH2	6:B:2091:HOH:O	2.43	0.52
1:B:848:ASN:C	1:B:848:ASN:HD22	2.14	0.49
1:B:808[B]:GLN:CD	1:B:808[B]:GLN:H	2.13	0.48
1:A:783:LEU:HD23	1:A:783:LEU:N	2.30	0.47
1:B:859:SER:CB	1:B:865:GLY:HA2	2.45	0.47
1:B:772:PHE:HA	1:B:868:ALA:HA	1.96	0.47
1:B:783:LEU:HD23	1:B:783:LEU:N	2.31	0.46
1:A:859:SER:CB	1:A:865:GLY:HA2	2.46	0.46
1:A:772:PHE:HA	1:A:868:ALA:HA	1.99	0.44
1:B:792[A]:ARG:CZ	6:B:2090:HOH:O	2.67	0.42
1:B:799:LYS:HA	1:B:842:THR:O	2.20	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:B:2204:HOH:O	6:B:2204:HOH:O[27_555]	1.62	0.58

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	140/155 (90%)	138 (99%)	2 (1%)	0	100	100
1	B	140/155 (90%)	139 (99%)	1 (1%)	0	100	100
All	All	280/310 (90%)	277 (99%)	3 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	115/127 (91%)	111 (96%)	4 (4%)	43	20
1	B	115/127 (91%)	112 (97%)	3 (3%)	54	32
All	All	230/254 (91%)	223 (97%)	7 (3%)	47	26

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	739	SER
1	A	783	LEU
1	A	838	SER
1	A	848	ASN
1	B	783	LEU
1	B	838	SER
1	B	848	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	797	GLN
1	A	848	ASN
1	B	797	GLN
1	B	848	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

4 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
4	GLA	A	1881	4	11,11,12	1.08	1 (9%)	14,15,17	1.68	4 (28%)
4	BMA	A	1882	4	12,12,12	1.03	0	17,17,17	1.94	4 (23%)
5	BMA	B	1881	5	12,12,12	1.33	2 (16%)	17,17,17	1.86	4 (23%)
5	GLA	B	1882	5	11,11,12	1.28	1 (9%)	14,15,17	1.88	4 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GLA	A	1881	4	-	0/2/19/22	0/1/1/1
4	BMA	A	1882	4	-	0/2/22/22	0/1/1/1
5	BMA	B	1881	5	-	0/2/22/22	0/1/1/1
5	GLA	B	1882	5	-	0/2/19/22	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1881	GLA	O5-C1	-2.22	1.40	1.43
5	B	1882	GLA	C2-C3	2.50	1.55	1.52
5	B	1881	BMA	O5-C1	2.57	1.47	1.43
5	B	1881	BMA	C1-C2	2.71	1.58	1.52

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1882	BMA	O4-C4-C3	-4.31	100.64	110.34
5	B	1882	GLA	C1-C2-C3	-3.23	105.72	109.54
4	A	1881	GLA	C3-C4-C5	-3.09	104.81	110.20
5	B	1881	BMA	C4-C3-C2	-2.55	106.03	110.79
5	B	1882	GLA	O6-C6-C5	-2.38	103.46	111.33
4	A	1881	GLA	O6-C6-C5	-2.34	103.60	111.33
4	A	1881	GLA	O3-C3-C2	-2.31	105.83	110.00
5	B	1882	GLA	C3-C4-C5	-2.18	106.39	110.20
4	A	1881	GLA	O5-C5-C6	2.50	112.75	107.35
4	A	1882	BMA	C4-C3-C2	2.80	116.01	110.79
5	B	1881	BMA	O5-C1-C2	3.08	114.72	109.80
5	B	1881	BMA	O2-C2-C1	3.15	116.76	109.82
4	A	1882	BMA	C6-C5-C4	3.27	121.08	113.02
5	B	1882	GLA	C1-O5-C5	3.32	116.46	112.25
5	B	1881	BMA	C1-O5-C5	3.47	119.90	113.47
4	A	1882	BMA	C3-C4-C5	3.81	116.83	110.20

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	1881	BMA	1	0

## 5.6 Ligand geometry

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	GOL	A	1879	-	5,5,5	0.40	0	5,5,5	1.01	0
2	GOL	B	1880	-	5,5,5	0.42	0	5,5,5	0.85	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	1879	-	-	0/4/4/4	0/0/0/0
2	GOL	B	1880	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	140/155 (90%)	-0.12	3 (2%) 67 71	10, 14, 28, 55	0
1	B	140/155 (90%)	-0.13	3 (2%) 67 71	10, 14, 27, 57	0
All	All	280/310 (90%)	-0.13	6 (2%) 67 71	10, 14, 27, 57	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	878	GLY	7.8
1	B	878	GLY	6.7
1	B	740	TYR	6.0
1	A	740	TYR	5.8
1	B	739	SER	4.9
1	A	739	SER	4.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
5	GLA	B	1882	11/12	0.96	0.09	0.80	14,16,19,20	0
4	GLA	A	1881	11/12	0.96	0.08	0.74	14,17,20,21	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
4	BMA	A	1882	12/12	0.79	0.33	-	24,65,71,73	0
5	BMA	B	1881	12/12	0.79	0.34	-	25,64,68,68	0

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	CA	B	1879	1/1	1.00	0.07	0.47	12,12,12,12	0
2	GOL	A	1879	6/6	0.96	0.07	-0.07	27,29,32,32	0
2	GOL	B	1880	6/6	0.96	0.07	-0.29	27,30,32,32	0
3	CA	A	1880	1/1	1.00	0.07	-0.48	12,12,12,12	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.