



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 22, 2016 – 08:05 AM EDT

PDB ID : 4YB7
Title : Adenosine triphosphate phosphoribosyltransferase from *Campylobacter jejuni*
in complex with ATP
Authors : Mittelstaedt, G.; Moggre, G.-J.; Parker, E.J.
Deposited on : 2015-02-18
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027107
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0122
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027107

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3774 (2.20-2.20)
Clashscore	102246	4477 (2.20-2.20)
Ramachandran outliers	100387	4404 (2.20-2.20)
Sidechain outliers	100360	4405 (2.20-2.20)
RSRZ outliers	91569	3781 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	300	<div><div></div><div>92%6% .</div></div>
1	B	300	<div><div></div><div>89%10% .</div></div>
1	C	300	<div><div>%</div><div>86%12% ..</div></div>
1	D	300	<div><div>%</div><div>91%7% .</div></div>
1	E	300	<div><div>%</div><div>90%8% .</div></div>
1	F	300	<div><div></div><div>89%7% ..</div></div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	300	 90% 6% ..
1	H	300	 91% 8% .
1	I	300	 88% 9% ..
1	J	300	 90% 8% .
1	K	300	 89% 8% ..
1	L	300	 87% 9% ..

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	PO4	J	302	-	-	-	X

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 26987 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ATP phosphoribosyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	296	Total	C	N	O	S	0	0	0
			2228	1418	381	416	13			
1	C	294	Total	C	N	O	S	0	1	0
			2198	1399	376	409	14			
1	D	294	Total	C	N	O	S	0	0	0
			2168	1381	370	404	13			
1	E	295	Total	C	N	O	S	0	0	0
			2203	1395	377	418	13			
1	I	292	Total	C	N	O	S	0	0	0
			2169	1377	376	403	13			
1	K	294	Total	C	N	O	S	0	1	0
			2168	1380	369	405	14			
1	A	296	Total	C	N	O	S	0	0	0
			2202	1404	379	406	13			
1	F	293	Total	C	N	O	S	0	1	0
			2191	1389	375	413	14			
1	G	292	Total	C	N	O	S	0	0	0
			2149	1364	365	407	13			
1	H	296	Total	C	N	O	S	5	0	0
			2162	1372	372	405	13			
1	J	296	Total	C	N	O	S	0	0	0
			2218	1409	379	417	13			
1	L	293	Total	C	N	O	S	0	1	0
			2148	1360	368	406	14			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	GLY	-	expression tag	UNP Q5HSJ4
C	0	GLY	-	expression tag	UNP Q5HSJ4
D	0	GLY	-	expression tag	UNP Q5HSJ4
E	0	GLY	-	expression tag	UNP Q5HSJ4
I	0	GLY	-	expression tag	UNP Q5HSJ4

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
K	0	GLY	-	expression tag	UNP Q5HSJ4
A	0	GLY	-	expression tag	UNP Q5HSJ4
F	0	GLY	-	expression tag	UNP Q5HSJ4
G	0	GLY	-	expression tag	UNP Q5HSJ4
H	0	GLY	-	expression tag	UNP Q5HSJ4
J	0	GLY	-	expression tag	UNP Q5HSJ4
L	0	GLY	-	expression tag	UNP Q5HSJ4

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	G	1	Total Mg 1 1	0	0
2	J	1	Total Mg 1 1	0	0
2	D	1	Total Mg 1 1	0	0
2	K	1	Total Mg 1 1	0	0
2	E	1	Total Mg 1 1	0	0
2	H	1	Total Mg 1 1	0	0
2	B	1	Total Mg 1 1	0	0
2	I	1	Total Mg 1 1	0	0
2	C	1	Total Mg 1 1	0	0
2	A	1	Total Mg 1 1	0	0
2	L	1	Total Mg 1 1	0	0
2	F	1	Total Mg 1 1	0	0

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	O	P	0	0
			5	4	1		
3	C	1	Total	O	P	0	0
			5	4	1		
3	C	1	Total	O	P	0	0
			5	4	1		
3	E	1	Total	O	P	0	0
			5	4	1		
3	I	1	Total	O	P	0	0
			5	4	1		
3	J	1	Total	O	P	0	0
			5	4	1		

- Molecule 4 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: $C_{10}H_{16}N_5O_{13}P_3$).



Id	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total 31	C 10	N 5	O 13	P 3	0	0
4	C	1	Total 31	C 10	N 5	O 13	P 3	0	0
4	D	1	Total 31	C 10	N 5	O 13	P 3	0	0
4	E	1	Total 31	C 10	N 5	O 13	P 3	0	0
4	I	1	Total 31	C 10	N 5	O 13	P 3	0	0
4	K	1	Total 31	C 10	N 5	O 13	P 3	0	0
4	A	1	Total 31	C 10	N 5	O 13	P 3	0	0
4	F	1	Total 31	C 10	N 5	O 13	P 3	0	0
4	G	1	Total 31	C 10	N 5	O 13	P 3	0	0
4	H	1	Total 31	C 10	N 5	O 13	P 3	0	0
4	J	1	Total 31	C 10	N 5	O 13	P 3	0	0
4	L	1	Total 31	C 10	N 5	O 13	P 3	0	0

- Molecule 5 is ACETIC ACID (three-letter code: ACY) (formula: $\text{C}_2\text{H}_4\text{O}_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	J	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	52	Total	O	0	0
			52	52		
6	C	44	Total	O	0	0
			44	44		
6	D	31	Total	O	0	0
			31	31		
6	E	19	Total	O	0	0
			19	19		
6	I	22	Total	O	0	0
			22	22		
6	K	15	Total	O	0	0
			15	15		
6	A	68	Total	O	0	0
			68	68		
6	F	29	Total	O	0	0
			29	29		
6	G	34	Total	O	0	0
			34	34		
6	H	12	Total	O	0	0
			12	12		
6	J	26	Total	O	0	0
			26	26		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	L	13	Total	O	0	0
			13	13		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ATP phosphoribosyltransferase




- Molecule 1: ATP phosphoribosyltransferase




- Molecule 1: ATP phosphoribosyltransferase




- Molecule 1: ATP phosphoribosyltransferase




- Molecule 1: ATP phosphoribosyltransferase








- Molecule 1: ATP phosphoribosyltransferase





- Molecule 1: ATP phosphoribosyltransferase




- Molecule 1: ATP phosphoribosyltransferase



- Molecule 1: ATP phosphoribosyltransferase



- Molecule 1: ATP phosphoribosyltransferase




● Molecule 1: ATP phosphoribosyltransferase

Chain J:

90%

8%




● Molecule 1: ATP phosphoribosyltransferase

Chain L:

87%

9%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	91.67Å 91.83Å 154.90Å 101.11° 95.21° 118.14°	Depositor
Resolution (Å)	50.00 – 2.20 48.06 – 2.20	Depositor EDS
% Data completeness (in resolution range)	98.1 (50.00-2.20) 88.2 (48.06-2.20)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.83 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.231 , 0.254 0.235 , 0.258	Depositor DCC
R_{free} test set	10624 reflections (5.25%)	DCC
Wilson B-factor (Å ²)	36.8	Xtriage
Anisotropy	0.028	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 30.2	EDS
Estimated twinning fraction	0.053 for k,h,-h-k-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 212950 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	26987	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.43% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, ACY, MG, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.49	0/2229	0.79	4/3020 (0.1%)
1	B	0.49	0/2257	0.75	6/3055 (0.2%)
1	C	0.48	0/2225	0.85	10/3011 (0.3%)
1	D	0.45	0/2195	0.78	6/2977 (0.2%)
1	E	0.44	0/2231	0.82	8/3026 (0.3%)
1	F	0.44	0/2216	0.78	8/2997 (0.3%)
1	G	0.44	0/2174	0.85	9/2950 (0.3%)
1	H	0.41	0/2190	0.80	9/2974 (0.3%)
1	I	0.47	0/2197	0.82	7/2976 (0.2%)
1	J	0.44	0/2247	0.80	8/3042 (0.3%)
1	K	0.44	0/2194	0.82	9/2975 (0.3%)
1	L	0.46	0/2174	0.83	10/2952 (0.3%)
All	All	0.45	0/26529	0.81	94/35955 (0.3%)

There are no bond length outliers.

The worst 5 of 94 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	192	ARG	NE-CZ-NH2	-14.79	112.91	120.30
1	I	192	ARG	NE-CZ-NH2	-14.55	113.03	120.30
1	J	192	ARG	NE-CZ-NH2	-14.34	113.13	120.30
1	H	192	ARG	NE-CZ-NH2	-14.30	113.15	120.30
1	A	192	ARG	NE-CZ-NH1	13.90	127.25	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2202	0	2219	16	0
1	B	2228	0	2247	24	0
1	C	2198	0	2200	30	0
1	D	2168	0	2142	28	0
1	E	2203	0	2185	11	1
1	F	2191	0	2190	15	1
1	G	2149	0	2110	21	0
1	H	2162	0	2109	17	0
1	I	2169	0	2159	21	0
1	J	2218	0	2210	23	1
1	K	2168	0	2131	16	1
1	L	2148	0	2087	34	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
2	G	1	0	0	0	0
2	H	1	0	0	0	0
2	I	1	0	0	0	0
2	J	1	0	0	0	0
2	K	1	0	0	0	0
2	L	1	0	0	0	0
3	B	5	0	0	0	0
3	C	10	0	0	0	0
3	E	5	0	0	0	0
3	I	5	0	0	0	0
3	J	5	0	0	0	0
4	A	31	0	12	2	0
4	B	31	0	12	2	0
4	C	31	0	12	0	0
4	D	31	0	12	2	0
4	E	31	0	12	0	0
4	F	31	0	12	0	0
4	G	31	0	12	1	0
4	H	31	0	12	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	I	31	0	12	1	0
4	J	31	0	12	2	0
4	K	31	0	12	1	0
4	L	31	0	12	2	0
5	J	4	0	3	0	0
6	A	68	0	0	2	0
6	B	52	0	0	1	0
6	C	44	0	0	2	0
6	D	31	0	0	3	0
6	E	19	0	0	0	0
6	F	29	0	0	1	0
6	G	34	0	0	0	0
6	H	12	0	0	0	0
6	I	22	0	0	0	0
6	J	26	0	0	2	0
6	K	15	0	0	0	0
6	L	13	0	0	0	0
All	All	26987	0	26136	207	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 207 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:138:PHE:HD2	1:D:139:MET:HE2	1.09	1.18
1:K:139:MET:O	1:K:142:ASN:O	1.68	1.12
1:E:156:GLU:OE1	1:E:176:THR:HG21	1.50	1.11
1:B:245:LEU:HD21	1:B:273:LEU:HD11	1.28	1.10
1:D:138:PHE:CD2	1:D:139:MET:CE	2.38	1.07

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:118:ASN:ND2	1:J:235:LYS:O[1_445]	1.72	0.48
1:E:235:LYS:NZ	1:K:141:GLU:OE2[1_655]	2.09	0.11

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	294/300 (98%)	291 (99%)	3 (1%)	0	100	100
1	B	294/300 (98%)	291 (99%)	3 (1%)	0	100	100
1	C	291/300 (97%)	288 (99%)	3 (1%)	0	100	100
1	D	290/300 (97%)	287 (99%)	3 (1%)	0	100	100
1	E	293/300 (98%)	290 (99%)	3 (1%)	0	100	100
1	F	290/300 (97%)	287 (99%)	3 (1%)	0	100	100
1	G	288/300 (96%)	285 (99%)	3 (1%)	0	100	100
1	H	294/300 (98%)	291 (99%)	3 (1%)	0	100	100
1	I	288/300 (96%)	284 (99%)	4 (1%)	0	100	100
1	J	294/300 (98%)	291 (99%)	3 (1%)	0	100	100
1	K	291/300 (97%)	288 (99%)	3 (1%)	0	100	100
1	L	290/300 (97%)	287 (99%)	3 (1%)	0	100	100
All	All	3497/3600 (97%)	3460 (99%)	37 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	227/263 (86%)	225 (99%)	2 (1%)	84	92
1	B	234/263 (89%)	231 (99%)	3 (1%)	76	87

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	228/263 (87%)	225 (99%)	3 (1%)	76	87
1	D	219/263 (83%)	218 (100%)	1 (0%)	92	96
1	E	229/263 (87%)	225 (98%)	4 (2%)	68	81
1	F	228/263 (87%)	223 (98%)	5 (2%)	60	72
1	G	218/263 (83%)	216 (99%)	2 (1%)	84	92
1	H	215/263 (82%)	213 (99%)	2 (1%)	84	92
1	I	223/263 (85%)	220 (99%)	3 (1%)	76	87
1	J	230/263 (88%)	227 (99%)	3 (1%)	76	87
1	K	218/263 (83%)	215 (99%)	3 (1%)	74	85
1	L	215/263 (82%)	212 (99%)	3 (1%)	74	85
All	All	2684/3156 (85%)	2650 (99%)	34 (1%)	76	87

5 of 34 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	K	265	LEU
1	F	114	ASN
1	L	185	VAL
1	A	170	LEU
1	D	265	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:

Mol	Chain	Res	Type
1	I	12	GLN
1	I	85	GLN
1	H	148	ASN
1	E	118	ASN
1	E	148	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 31 ligands modelled in this entry, 12 are monoatomic - leaving 19 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	ATP	A	302	2	26,33,33	1.11	2 (7%)	26,52,52	1.81	2 (7%)
3	PO4	B	302	-	4,4,4	0.61	0	6,6,6	0.24	0
4	ATP	B	303	2	26,33,33	1.00	2 (7%)	26,52,52	1.94	3 (11%)
3	PO4	C	302	-	4,4,4	0.70	0	6,6,6	0.23	0
3	PO4	C	303	-	4,4,4	0.60	0	6,6,6	0.23	0
4	ATP	C	304	2	26,33,33	1.08	2 (7%)	26,52,52	1.85	2 (7%)
4	ATP	D	302	2	26,33,33	0.98	1 (3%)	26,52,52	2.11	3 (11%)
3	PO4	E	302	-	4,4,4	0.54	0	6,6,6	0.24	0
4	ATP	E	303	2	26,33,33	1.03	2 (7%)	26,52,52	1.84	4 (15%)
4	ATP	F	302	2	26,33,33	1.18	3 (11%)	26,52,52	1.99	5 (19%)
4	ATP	G	302	2	26,33,33	1.05	2 (7%)	26,52,52	1.82	2 (7%)
4	ATP	H	302	2	26,33,33	0.98	1 (3%)	26,52,52	1.93	2 (7%)
3	PO4	I	302	-	4,4,4	0.56	0	6,6,6	0.23	0
4	ATP	I	303	2	26,33,33	0.97	2 (7%)	26,52,52	2.00	3 (11%)
3	PO4	J	302	-	4,4,4	0.60	0	6,6,6	0.24	0
4	ATP	J	303	2	26,33,33	0.96	2 (7%)	26,52,52	1.85	3 (11%)
5	ACY	J	304	-	0,3,3	0.00	-	0,3,3	0.00	-
4	ATP	K	302	2	26,33,33	1.04	2 (7%)	26,52,52	1.86	1 (3%)
4	ATP	L	302	2	26,33,33	1.07	2 (7%)	26,52,52	2.02	5 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ATP	A	302	2	-	0/18/38/38	0/3/3/3
3	PO4	B	302	-	-	0/0/0/0	0/0/0/0
4	ATP	B	303	2	-	0/18/38/38	0/3/3/3
3	PO4	C	302	-	-	0/0/0/0	0/0/0/0
3	PO4	C	303	-	-	0/0/0/0	0/0/0/0
4	ATP	C	304	2	-	0/18/38/38	0/3/3/3
4	ATP	D	302	2	-	0/18/38/38	0/3/3/3
3	PO4	E	302	-	-	0/0/0/0	0/0/0/0
4	ATP	E	303	2	-	0/18/38/38	0/3/3/3
4	ATP	F	302	2	-	0/18/38/38	0/3/3/3
4	ATP	G	302	2	-	0/18/38/38	0/3/3/3
4	ATP	H	302	2	-	0/18/38/38	0/3/3/3
3	PO4	I	302	-	-	0/0/0/0	0/0/0/0
4	ATP	I	303	2	-	0/18/38/38	0/3/3/3
3	PO4	J	302	-	-	0/0/0/0	0/0/0/0
4	ATP	J	303	2	-	0/18/38/38	0/3/3/3
5	ACY	J	304	-	-	0/0/0/0	0/0/0/0
4	ATP	K	302	2	-	0/18/38/38	0/3/3/3
4	ATP	L	302	2	-	0/18/38/38	0/3/3/3

The worst 5 of 23 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	K	302	ATP	C2-N3	2.02	1.35	1.32
4	I	303	ATP	C2-N3	2.11	1.35	1.32
4	J	303	ATP	C2-N3	2.18	1.36	1.32
4	A	302	ATP	C2-N3	2.23	1.36	1.32
4	E	303	ATP	C2-N3	2.29	1.36	1.32

The worst 5 of 35 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	302	ATP	N3-C2-N1	-8.69	122.04	128.87
4	H	302	ATP	N3-C2-N1	-8.36	122.30	128.87
4	I	303	ATP	N3-C2-N1	-8.16	122.46	128.87
4	L	302	ATP	N3-C2-N1	-8.15	122.47	128.87
4	K	302	ATP	N3-C2-N1	-8.15	122.47	128.87

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

8 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	302	ATP	2	0
4	B	303	ATP	2	0
4	D	302	ATP	2	0
4	G	302	ATP	1	0
4	I	303	ATP	1	0
4	J	303	ATP	2	0
4	K	302	ATP	1	0
4	L	302	ATP	2	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	296/300 (98%)	-0.17	0 100 100	18, 35, 64, 92	0
1	B	296/300 (98%)	-0.15	1 (0%) 94 94	18, 38, 72, 92	0
1	C	294/300 (98%)	-0.09	2 (0%) 89 88	19, 39, 77, 94	0
1	D	294/300 (98%)	-0.09	2 (0%) 89 88	23, 43, 79, 99	0
1	E	295/300 (98%)	-0.10	3 (1%) 84 83	23, 49, 79, 100	0
1	F	293/300 (97%)	-0.20	1 (0%) 94 94	21, 42, 69, 85	0
1	G	292/300 (97%)	-0.05	1 (0%) 94 94	23, 44, 77, 99	0
1	H	295/300 (98%)	0.03	4 (1%) 78 77	33, 55, 85, 102	0
1	I	292/300 (97%)	-0.03	2 (0%) 89 88	23, 48, 82, 92	0
1	J	296/300 (98%)	-0.23	1 (0%) 94 94	23, 42, 69, 82	0
1	K	294/300 (98%)	-0.00	5 (1%) 73 72	30, 52, 81, 102	0
1	L	293/300 (97%)	0.09	7 (2%) 62 61	28, 52, 82, 91	0
All	All	3530/3600 (98%)	-0.08	29 (0%) 87 87	18, 45, 78, 102	0

The worst 5 of 29 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	257	ALA	3.7
1	K	175	ALA	3.6
1	E	175	ALA	3.4
1	L	181	ASN	3.3
1	B	236	GLU	3.3

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	PO4	J	302	5/5	0.55	0.47	8.20	128,130,132,136	0
3	PO4	I	302	5/5	0.72	0.23	1.99	93,96,97,98	0
3	PO4	C	302	5/5	0.85	0.27	1.57	92,104,106,111	0
3	PO4	E	302	5/5	0.56	0.23	0.73	103,108,109,114	0
4	ATP	D	302	31/31	0.93	0.16	0.69	42,56,70,71	0
3	PO4	B	302	5/5	0.88	0.16	0.58	92,95,99,99	0
4	ATP	A	302	31/31	0.94	0.14	0.36	27,45,61,62	0
4	ATP	C	304	31/31	0.94	0.13	-0.03	28,40,54,56	0
4	ATP	E	303	31/31	0.93	0.13	-0.07	44,54,70,74	0
4	ATP	L	302	31/31	0.90	0.14	-0.17	38,55,69,70	0
4	ATP	I	303	31/31	0.94	0.11	-0.20	37,48,68,77	0
4	ATP	H	302	31/31	0.93	0.12	-0.23	45,56,68,72	0
4	ATP	J	303	31/31	0.95	0.12	-0.31	30,44,60,66	0
4	ATP	B	303	31/31	0.94	0.12	-0.34	36,43,53,55	0
4	ATP	G	302	31/31	0.93	0.12	-0.39	37,52,62,68	0
4	ATP	K	302	31/31	0.92	0.13	-0.53	39,59,71,76	0
4	ATP	F	302	31/31	0.94	0.11	-0.86	28,42,58,59	0
2	MG	H	301	1/1	0.97	0.05	-	55,55,55,55	0
2	MG	I	301	1/1	0.94	0.11	-	43,43,43,43	0
2	MG	D	301	1/1	0.89	0.07	-	49,49,49,49	0
2	MG	J	301	1/1	0.95	0.06	-	42,42,42,42	0
2	MG	E	301	1/1	0.94	0.09	-	59,59,59,59	0
3	PO4	C	303	5/5	0.66	0.19	-	98,101,108,110	0
2	MG	G	301	1/1	0.94	0.05	-	51,51,51,51	0
2	MG	B	301	1/1	0.94	0.04	-	39,39,39,39	0
2	MG	F	301	1/1	0.95	0.07	-	39,39,39,39	0
2	MG	C	301	1/1	0.95	0.07	-	45,45,45,45	0
2	MG	A	301	1/1	0.99	0.07	-	44,44,44,44	0
2	MG	K	301	1/1	0.88	0.12	-	70,70,70,70	0
5	ACY	J	304	4/4	0.58	0.19	-	56,60,62,65	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	MG	L	301	1/1	0.84	0.19	-	62,62,62,62	0

6.5 Other polymers [i](#)

There are no such residues in this entry.