



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 10:37 PM GMT

PDB ID : 4YDU
Title : Crystal structure of E. coli YgjD-YeaZ heterodimer in complex with ADP
Authors : Zhang, W.; Collinet, B.; Perrochia, L.; Durand, D.; Van Tilbeurgh, H.
Deposited on : 2015-02-23
Resolution : 2.33 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.33 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1406 (2.36-2.32)
Clashscore	102246	1509 (2.36-2.32)
Ramachandran outliers	100387	1490 (2.36-2.32)
Sidechain outliers	100360	1491 (2.36-2.32)
RSRZ outliers	91569	1412 (2.36-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	343	<div> <div></div> <div>89% 9% .</div> </div>
1	B	343	<div> <div></div> <div>89% 9% ..</div> </div>
2	C	237	<div> <div></div> <div>89% 7% ..</div> </div>
2	D	237	<div> <div></div> <div>84% 10% ..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	ACT	A	407	-	-	X	X
6	ACT	C	303	-	-	-	X
6	ACT	D	302	-	-	X	-
6	ACT	D	303	-	-	-	X
6	ACT	D	305	-	-	-	X
6	ACT	D	306	-	-	X	X

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 9012 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called tRNA N6-adenosine threonylcarbamoyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	336	Total	C	N	O	S	0	0	0
			2470	1573	423	460	14			
1	B	338	Total	C	N	O	S	0	0	0
			2498	1587	429	468	14			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	338	HIS	-	expression tag	UNP P05852
A	339	HIS	-	expression tag	UNP P05852
A	340	HIS	-	expression tag	UNP P05852
A	341	HIS	-	expression tag	UNP P05852
A	342	HIS	-	expression tag	UNP P05852
A	343	HIS	-	expression tag	UNP P05852
B	338	HIS	-	expression tag	UNP P05852
B	339	HIS	-	expression tag	UNP P05852
B	340	HIS	-	expression tag	UNP P05852
B	341	HIS	-	expression tag	UNP P05852
B	342	HIS	-	expression tag	UNP P05852
B	343	HIS	-	expression tag	UNP P05852


- Molecule 2 is a protein called tRNA threonylcarbamoyladenine biosynthesis protein Tsab.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	230	Total	C	N	O	S	0	0	0
			1718	1091	294	321	12			
2	D	229	Total	C	N	O	S	0	0	0
			1724	1095	299	318	12			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	232	HIS	-	expression tag	UNP P76256
C	233	HIS	-	expression tag	UNP P76256
C	234	HIS	-	expression tag	UNP P76256
C	235	HIS	-	expression tag	UNP P76256
C	236	HIS	-	expression tag	UNP P76256
C	237	HIS	-	expression tag	UNP P76256
D	232	HIS	-	expression tag	UNP P76256
D	233	HIS	-	expression tag	UNP P76256
D	234	HIS	-	expression tag	UNP P76256
D	235	HIS	-	expression tag	UNP P76256
D	236	HIS	-	expression tag	UNP P76256
D	237	HIS	-	expression tag	UNP P76256

- Molecule 3 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
3	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
3	C	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
3	D	1	Total	C	N	O	P	0	0
			27	10	5	10	2		


- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total Mg 1 1	0	0
4	A	1	Total Mg 1 1	0	0

- Molecule 5 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total Fe 1 1	0	0
5	A	1	Total Fe 1 1	0	0

- Molecule 6 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	B	1	Total C O 4 2 2	0	0
6	B	1	Total C O 4 2 2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	C	1	Total C O 4 2 2	0	0
6	C	1	Total C O 4 2 2	0	0
6	D	1	Total C O 4 2 2	0	0
6	D	1	Total C O 4 2 2	0	0
6	D	1	Total C O 4 2 2	0	0
6	D	1	Total C O 4 2 2	0	0
6	D	1	Total C O 4 2 2	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	105	Total O 105 105	0	0
7	B	124	Total O 124 124	0	0
7	C	103	Total O 103 103	0	0
7	D	106	Total O 106 106	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: tRNA N6-adenosine threonylcarbamoyltransferase




- Molecule 1: tRNA N6-adenosine threonylcarbamoyltransferase



- Molecule 2: tRNA threonylcarbamoyladenosine biosynthesis protein Tsab



- Molecule 2: tRNA threonylcarbamoyladenosine biosynthesis protein Tsab



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	63.62Å 68.48Å 87.07Å 109.38° 92.66° 117.66°	Depositor
Resolution (Å)	49.38 – 2.33 49.38 – 2.25	Depositor EDS
% Data completeness (in resolution range)	97.9 (49.38-2.33) 82.8 (49.38-2.25)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.75 (at 2.25Å)	Xtriage
Refinement program	REFMAC 5.8.0071	Depositor
R, R_{free}	0.202 , 0.253 0.173 , 0.221	Depositor DCC
R_{free} test set	2503 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	28.6	Xtriage
Anisotropy	0.179	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 34.1	EDS
Estimated twinning fraction	0.045 for h,-h-k,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 55653 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9012	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.41% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, FE, ADP, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.68	0/2519	0.80	2/3423 (0.1%)
1	B	0.78	2/2548 (0.1%)	0.90	5/3462 (0.1%)
2	C	0.78	1/1755 (0.1%)	0.81	4/2394 (0.2%)
2	D	1.08	6/1761 (0.3%)	0.99	10/2398 (0.4%)
All	All	0.82	9/8583 (0.1%)	0.87	21/11677 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	C	0	1
2	D	0	1
All	All	0	2

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	117	ALA	C-N	-24.97	0.76	1.34
2	C	224	TRP	C-N	15.13	1.68	1.34
2	D	104	ARG	C-N	-14.44	1.00	1.34
1	B	291	PHE	C-N	-11.68	1.07	1.34
2	D	50	GLY	C-N	-10.90	1.08	1.34

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	291	PHE	O-C-N	-14.12	100.10	122.70
1	B	291	PHE	C-N-CA	13.82	156.25	121.70

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	D	117	ALA	C-N-CA	13.49	155.43	121.70
2	D	104	ARG	O-C-N	-10.62	105.71	122.70
1	B	291	PHE	CA-C-N	9.89	138.97	117.20

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	C	223	ALA	Mainchain
2	D	118	ARG	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2470	0	2428	26	0
1	B	2498	0	2458	20	0
2	C	1718	0	1658	13	0
2	D	1724	0	1690	20	0
3	A	27	0	12	0	0
3	B	27	0	12	0	0
3	C	27	0	12	0	0
3	D	27	0	12	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	16	0	12	2	0
6	B	8	0	6	1	0
6	C	8	0	6	0	0
6	D	20	0	15	5	0
7	A	105	0	0	2	0
7	B	124	0	0	2	1
7	C	103	0	0	0	0
7	D	106	0	0	2	1
All	All	9012	0	8321	77	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 77 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:224:TRP:C	2:C:225:LYS:N	1.68	1.42
2:D:117:ALA:O	2:D:118:ARG:N	1.59	1.33
2:D:117:ALA:CA	2:D:118:ARG:N	2.10	1.14
2:D:117:ALA:C	2:D:118:ARG:CA	2.19	1.10
2:D:79:GLN:HE22	6:D:306:ACT:H1	1.43	0.82

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:B:517:HOH:O	7:D:403:HOH:O[1_554]	2.10	0.10

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	334/343 (97%)	326 (98%)	8 (2%)	0	100	100
1	B	336/343 (98%)	326 (97%)	9 (3%)	1 (0%)	46	54
2	C	228/237 (96%)	221 (97%)	7 (3%)	0	100	100
2	D	227/237 (96%)	219 (96%)	7 (3%)	1 (0%)	39	45
All	All	1125/1160 (97%)	1092 (97%)	31 (3%)	2 (0%)	52	61

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	292	TYR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	D	228	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	239/264 (90%)	237 (99%)	2 (1%)	86	93
1	B	245/264 (93%)	244 (100%)	1 (0%)	93	97
2	C	165/187 (88%)	164 (99%)	1 (1%)	90	96
2	D	170/187 (91%)	170 (100%)	0	100	100
All	All	819/902 (91%)	815 (100%)	4 (0%)	92	96

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	50	ASP
1	A	292	TYR
1	B	50	ASP
2	C	168	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	D	79	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 21 ligands modelled in this entry, 4 are monoatomic - leaving 17 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ADP	A	401	-	22,29,29	0.96	0	27,45,45	2.36	11 (40%)
6	ACT	A	404	-	1,3,3	2.07	1 (100%)	0,3,3	0.00	-
6	ACT	A	405	-	1,3,3	1.80	0	0,3,3	0.00	-
6	ACT	A	406	-	1,3,3	2.02	1 (100%)	0,3,3	0.00	-
6	ACT	A	407	-	1,3,3	2.58	1 (100%)	0,3,3	0.00	-
3	ADP	B	401	5	22,29,29	0.92	1 (4%)	27,45,45	2.10	5 (18%)
6	ACT	B	404	-	1,3,3	2.98	1 (100%)	0,3,3	0.00	-
6	ACT	B	405	-	1,3,3	2.16	1 (100%)	0,3,3	0.00	-
3	ADP	C	301	-	22,29,29	0.88	1 (4%)	27,45,45	2.70	5 (18%)
6	ACT	C	302	-	1,3,3	1.76	0	0,3,3	0.00	-
6	ACT	C	303	-	1,3,3	2.55	1 (100%)	0,3,3	0.00	-
3	ADP	D	301	-	22,29,29	1.02	3 (13%)	27,45,45	2.51	6 (22%)
6	ACT	D	302	-	1,3,3	1.03	0	0,3,3	0.00	-
6	ACT	D	303	-	1,3,3	0.48	0	0,3,3	0.00	-
6	ACT	D	304	-	1,3,3	3.17	1 (100%)	0,3,3	0.00	-
6	ACT	D	305	-	1,3,3	1.28	0	0,3,3	0.00	-
6	ACT	D	306	-	1,3,3	0.51	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ADP	A	401	-	-	0/12/32/32	0/3/3/3
6	ACT	A	404	-	-	0/0/0/0	0/0/0/0
6	ACT	A	405	-	-	0/0/0/0	0/0/0/0
6	ACT	A	406	-	-	0/0/0/0	0/0/0/0
6	ACT	A	407	-	-	0/0/0/0	0/0/0/0
3	ADP	B	401	5	-	0/12/32/32	0/3/3/3
6	ACT	B	404	-	-	0/0/0/0	0/0/0/0
6	ACT	B	405	-	-	0/0/0/0	0/0/0/0
3	ADP	C	301	-	-	0/12/32/32	0/3/3/3
6	ACT	C	302	-	-	0/0/0/0	0/0/0/0
6	ACT	C	303	-	-	0/0/0/0	0/0/0/0
3	ADP	D	301	-	-	0/12/32/32	0/3/3/3
6	ACT	D	302	-	-	0/0/0/0	0/0/0/0
6	ACT	D	303	-	-	0/0/0/0	0/0/0/0
6	ACT	D	304	-	-	0/0/0/0	0/0/0/0
6	ACT	D	305	-	-	0/0/0/0	0/0/0/0
6	ACT	D	306	-	-	0/0/0/0	0/0/0/0

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	406	ACT	CH3-C	2.02	1.51	1.48
3	D	301	ADP	C2-N3	2.05	1.35	1.32
6	A	404	ACT	CH3-C	2.07	1.51	1.48
3	D	301	ADP	O4'-C1'	2.07	1.43	1.41
6	B	405	ACT	CH3-C	2.16	1.51	1.48

The worst 5 of 27 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	301	ADP	C2'-C1'-N9	-9.24	100.17	114.29
3	D	301	ADP	N3-C2-N1	-9.10	121.92	128.89
3	C	301	ADP	N3-C2-N1	-8.45	122.43	128.89
3	A	401	ADP	N3-C2-N1	-8.06	122.73	128.89
3	B	401	ADP	N3-C2-N1	-7.88	122.86	128.89

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	407	ACT	2	0
6	B	404	ACT	1	0
6	D	302	ACT	2	0
6	D	305	ACT	1	0
6	D	306	ACT	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
2	D	4
1	B	1
2	C	1

The worst 5 of 6 chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	C	224:TRP	C	225:LYS	N	1.68
1	D	103:TRP	C	104:ARG	N	1.10
1	D	50:GLY	C	51:THR	N	1.09
1	B	291:PHE	C	292:TYR	N	1.07
1	D	104:ARG	C	105:LYS	N	1.00

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	336/343 (97%)	-0.39	2 (0%) 90 94	19, 37, 68, 86	0
1	B	338/343 (98%)	-0.53	3 (0%) 85 91	19, 30, 54, 73	0
2	C	230/237 (97%)	-0.52	3 (1%) 79 86	20, 32, 62, 84	0
2	D	229/237 (96%)	-0.60	1 (0%) 93 97	18, 30, 56, 75	0
All	All	1133/1160 (97%)	-0.50	9 (0%) 87 92	18, 32, 61, 86	0

The worst 5 of 9 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	229	GLY	2.9
1	A	226	GLY	2.5
2	C	224	TRP	2.5
1	B	338	HIS	2.3
2	C	223	ALA	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
6	ACT	A	407	4/4	0.82	0.19	8.12	40,41,48,51	0
6	ACT	D	306	4/4	0.87	0.20	6.19	32,47,49,53	0
6	ACT	C	303	4/4	0.88	0.25	4.64	42,45,47,47	0
6	ACT	D	303	4/4	0.94	0.12	2.65	51,52,52,54	0
6	ACT	D	305	4/4	0.91	0.16	2.39	37,37,45,53	0
6	ACT	A	405	4/4	0.90	0.15	1.23	39,41,42,44	0
5	FE	B	403	1/1	1.00	0.09	0.63	23,23,23,23	0
3	ADP	B	401	27/27	0.98	0.09	0.18	19,23,32,38	0
3	ADP	A	401	27/27	0.98	0.08	-0.59	23,25,36,38	0
3	ADP	C	301	27/27	0.95	0.08	-1.13	30,37,61,70	0
3	ADP	D	301	27/27	0.96	0.07	-1.15	24,29,51,60	0
6	ACT	A	404	4/4	0.94	0.07	-1.66	39,41,42,49	0
6	ACT	B	405	4/4	0.80	0.24	-	49,50,56,60	0
4	MG	A	402	1/1	0.95	0.04	-	33,33,33,33	0
6	ACT	C	302	4/4	0.89	0.13	-	46,48,50,51	0
4	MG	B	402	1/1	0.95	0.07	-	35,35,35,35	0
6	ACT	D	302	4/4	0.95	0.19	-	41,44,45,47	0
5	FE	A	403	1/1	1.00	0.07	-	25,25,25,25	0
6	ACT	A	406	4/4	0.96	0.09	-	45,48,56,61	0
6	ACT	B	404	4/4	0.71	0.19	-	41,51,53,59	0
6	ACT	D	304	4/4	0.86	0.14	-	48,50,54,61	0

6.5 Other polymers [i](#)

There are no such residues in this entry.