



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2016 – 08:28 PM GMT

PDB ID : 4YHJ
Title : Structure and Function of the Hypertension Variant A486V of G Protein-coupled Receptor Kinase 4 (GRK4)
Authors : Allen, S.J.; Parthasarathy, G.; Soisson, S.; Munshi, S.
Deposited on : 2015-02-27
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026982
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20026982


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2328 (2.60-2.60)
Clashscore	102246	2679 (2.60-2.60)
Ramachandran outliers	100387	2635 (2.60-2.60)
Sidechain outliers	100360	2635 (2.60-2.60)
RSRZ outliers	91569	2334 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	584	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; width: 100%; height: 1px; background-color: red;"></div> <div style="position: absolute; bottom: 0; left: 0; width: 100%; height: 10px;"> <div style="width: 72%; height: 10px; background-color: green;"></div> <div style="width: 13%; height: 10px; background-color: yellow;"></div> <div style="width: 1%; height: 10px; background-color: orange;"></div> <div style="width: 14%; height: 10px; background-color: grey;"></div> </div> </div> <div>72% 13% • 14%</div> </div>
1	B	584	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; width: 100%; height: 1px; background-color: red;"></div> <div style="position: absolute; bottom: 0; left: 0; width: 100%; height: 10px;"> <div style="width: 70%; height: 10px; background-color: green;"></div> <div style="width: 13%; height: 10px; background-color: yellow;"></div> <div style="width: 1%; height: 10px; background-color: orange;"></div> <div style="width: 16%; height: 10px; background-color: grey;"></div> </div> </div> <div>70% 13% • 16%</div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 8222 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called G protein-coupled receptor kinase 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	501	Total	C	N	O	S	0	0	0
			4067	2580	708	752	27			
1	B	490	Total	C	N	O	S	0	0	0
			3978	2521	696	734	27			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	563	SER	CYS	engineered mutation	UNP P32298
A	578	SER	CYS	engineered mutation	UNP P32298
A	579	GLU	-	expression tag	UNP P32298
A	580	ASN	-	expression tag	UNP P32298
A	581	LEU	-	expression tag	UNP P32298
A	582	TYR	-	expression tag	UNP P32298
A	583	PHE	-	expression tag	UNP P32298
A	584	GLN	-	expression tag	UNP P32298
B	563	SER	CYS	engineered mutation	UNP P32298
B	578	SER	CYS	engineered mutation	UNP P32298
B	579	GLU	-	expression tag	UNP P32298
B	580	ASN	-	expression tag	UNP P32298
B	581	LEU	-	expression tag	UNP P32298
B	582	TYR	-	expression tag	UNP P32298
B	583	PHE	-	expression tag	UNP P32298
B	584	GLN	-	expression tag	UNP P32298

- Molecule 2 is AMP PHOSPHORAMIDATE (three-letter code: AN2) (formula: C₁₀H₁₆N₆O₉P₂).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			27	10	6	9	2		
2	B	1	Total	C	N	O	P	0	0
			27	10	6	9	2		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	62	Total	O	0	0
			62	62		
3	B	61	Total	O	0	0
			61	61		

- Molecule 1: G protein-coupled receptor kinase 4



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	104.84Å 104.84Å 221.84Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	47.00 – 2.60 47.33 – 2.49	Depositor EDS
% Data completeness (in resolution range)	99.5 (47.00-2.60) 99.3 (47.33-2.49)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.11 (at 2.48Å)	Xtriage
Refinement program	BUSTER-TNT	Depositor
R, R_{free}	0.197 , 0.248 0.200 , 0.252	Depositor DCC
R_{free} test set	2254 reflections (5.39%)	DCC
Wilson B-factor (Å ²)	46.6	Xtriage
Anisotropy	0.138	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 38.8	EDS
Estimated twinning fraction	0.027 for -h,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 49678 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8222	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: AN2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.51	0/4158	0.76	2/5599 (0.0%)
1	B	0.51	0/4067	0.73	0/5474
All	All	0.51	0/8225	0.75	2/11073 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	477	ASP	C-N-CA	6.51	137.97	121.70
1	A	482	GLU	C-N-CA	5.71	135.98	121.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4067	0	4025	35	0
1	B	3978	0	3932	26	0
2	A	27	0	14	0	0
2	B	27	0	14	0	0
3	A	62	0	0	2	0
3	B	61	0	0	0	0
All	All	8222	0	7985	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:60:GLN:HE21	1:A:513:GLN:HE22	1.24	0.84
1:B:60:GLN:HE21	1:B:513:GLN:HE22	1.25	0.79
1:B:344:ARG:HH21	1:B:360:ASN:HD21	1.36	0.73
3:A:708:HOH:O	1:B:155:ARG:HD2	1.91	0.69
1:B:204:CYS:SG	1:B:215:CYS:HB2	2.32	0.69
1:A:487:VAL:HA	1:A:490:ILE:HD12	1.77	0.66
1:A:344:ARG:HD2	1:A:360:ASN:HD21	1.60	0.65
1:A:388:LYS:HE3	1:A:388:LYS:H	1.61	0.65
1:A:345:GLY:H	1:A:360:ASN:HD22	1.47	0.63
1:A:211:LYS:HD2	1:A:515:GLU:HG3	1.84	0.59
1:B:43:SER:O	1:B:47:HIS:HD2	1.86	0.58
1:B:253:ALA:O	1:B:506:GLY:HA2	2.03	0.58
1:A:256:THR:HG22	1:A:258:ASP:H	1.68	0.58
1:A:345:GLY:H	1:A:360:ASN:ND2	2.02	0.57
1:A:446:HIS:ND1	1:A:448:VAL:HG22	2.20	0.57
1:A:356:GLU:HG2	1:A:429:PRO:HG3	1.90	0.54
1:B:79:ARG:HD3	1:B:117:PRO:HB3	1.91	0.53
1:B:522:PHE:O	1:B:526:ASN:HB3	2.09	0.52
1:A:479:LEU:CB	1:A:481:ILE:HD12	2.40	0.52
1:A:204:CYS:SG	1:A:215:CYS:HB2	2.50	0.51
1:A:79:ARG:HD3	1:A:117:PRO:HB3	1.93	0.51
1:A:346:ARG:HG3	1:A:358:VAL:HA	1.92	0.51
1:A:252:TYR:HB2	1:A:263:VAL:HB	1.95	0.49
1:A:221:LYS:HD3	1:A:487:VAL:HB	1.95	0.48
1:A:356:GLU:CG	1:A:429:PRO:HG3	2.43	0.48
1:B:346:ARG:HG3	1:B:358:VAL:HA	1.96	0.48
1:A:479:LEU:HB3	1:A:481:ILE:HD12	1.96	0.47
1:B:311:ARG:HH21	1:B:335:THR:HG21	1.81	0.46
1:A:194:GLY:HA2	1:A:478:VAL:HG21	1.98	0.46
1:B:438:GLU:HB2	1:B:442:GLY:HA3	1.98	0.45
1:A:393:VAL:HG21	1:A:401:ARG:HH12	1.80	0.45
1:B:198:PHE:HE1	1:B:226:ARG:HB2	1.81	0.45
1:A:250:LEU:HA	1:A:264:LEU:HD23	1.99	0.45
1:B:356:GLU:HG2	1:B:429:PRO:HG3	1.98	0.45
1:B:218:LEU:HB3	1:B:223:ILE:HD11	1.99	0.45
1:B:242:VAL:HG21	1:B:303:LEU:HD23	1.98	0.45
1:B:396:GLU:HG2	1:B:396:GLU:H	1.46	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:383:HIS:CD2	1:B:388:LYS:HA	2.52	0.44
1:B:433:LEU:HG	1:B:443:VAL:HG22	2.00	0.44
1:A:107:ASP:HA	1:A:111:ASN:ND2	2.33	0.44
1:B:276:ILE:HD11	1:B:379:MET:HG2	1.99	0.44
1:A:375:LEU:HD12	1:A:375:LEU:HA	1.86	0.43
1:A:482:GLU:HB3	1:A:483:GLN:H	1.29	0.43
1:A:151:HIS:O	1:A:155:ARG:HB2	2.19	0.43
1:A:477:ASP:HA	1:A:478:VAL:HB	2.01	0.42
1:B:234:ASN:O	1:B:238:ILE:HG12	2.19	0.42
1:B:449:PHE:HA	1:B:452:ILE:HD12	2.01	0.42
1:B:446:HIS:ND1	1:B:448:VAL:HG22	2.35	0.42
1:A:30:TRP:HA	1:A:33:ILE:HD12	2.02	0.42
1:A:311:ARG:HD2	3:A:734:HOH:O	2.19	0.42
1:A:479:LEU:HB2	1:A:481:ILE:HD12	2.02	0.41
1:A:428:ASN:HA	1:A:429:PRO:HD2	1.94	0.41
1:A:46:ARG:HB2	1:A:172:LEU:HD13	2.02	0.41
1:A:481:ILE:HG12	1:A:482:GLU:HG3	2.03	0.41
1:B:133:LEU:HD11	1:B:143:PHE:CZ	2.56	0.41
1:A:433:LEU:HG	1:A:443:VAL:HG22	2.02	0.41
1:B:90:TYR:CE1	1:B:98:ARG:HG2	2.56	0.41
1:B:355:PRO:HG2	1:B:427:LYS:HA	2.02	0.41
1:A:388:LYS:HZ1	1:A:401:ARG:HH22	1.68	0.40
1:B:453:ASN:CG	1:B:456:ARG:HG3	2.42	0.40
1:A:122:PRO:HA	1:A:123:PRO:HD3	1.97	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	499/584 (85%)	472 (95%)	23 (5%)	4 (1%)	24 46

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	B	486/584 (83%)	466 (96%)	19 (4%)	1 (0%)	52 77
All	All	985/1168 (84%)	938 (95%)	42 (4%)	5 (0%)	34 60

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	483	GLN
1	A	478	VAL
1	A	137	ASN
1	B	391	GLU
1	A	481	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	441/515 (86%)	405 (92%)	36 (8%)	14 27
1	B	430/515 (84%)	397 (92%)	33 (8%)	16 31
All	All	871/1030 (85%)	802 (92%)	69 (8%)	15 30

All (69) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	28	LYS
1	A	51	LYS
1	A	59	LYS
1	A	65	ARG
1	A	103	LEU
1	A	106	LEU
1	A	113	LYS
1	A	114	LEU
1	A	131	LEU
1	A	135	GLU
1	A	157	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	177	LEU
1	A	191	ARG
1	A	217	LYS
1	A	219	GLN
1	A	222	ARG
1	A	233	LEU
1	A	236	LYS
1	A	249	SER
1	A	257	LYS
1	A	285	ASP
1	A	286	GLU
1	A	296	LEU
1	A	313	LEU
1	A	322	ASP
1	A	375	LEU
1	A	388	LYS
1	A	397	GLU
1	A	406	THR
1	A	411	GLU
1	A	431	LYS
1	A	448	VAL
1	A	463	GLU
1	A	476	LYS
1	A	481	ILE
1	A	483	GLN
1	B	28	LYS
1	B	32	GLU
1	B	41	GLN
1	B	49	ILE
1	B	66	LEU
1	B	99	SER
1	B	103	LEU
1	B	106	LEU
1	B	113	LYS
1	B	124	ASP
1	B	136	GLU
1	B	175	LYS
1	B	177	LEU
1	B	227	LYS
1	B	236	LYS
1	B	257	LYS
1	B	268	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	287	GLN
1	B	311	ARG
1	B	335	THR
1	B	356	GLU
1	B	393	VAL
1	B	396	GLU
1	B	405	ASP
1	B	408	GLU
1	B	411	GLU
1	B	415	GLU
1	B	438	GLU
1	B	451	ASP
1	B	455	ARG
1	B	475	CYS
1	B	498	ASP
1	B	507	CYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	60	GLN
1	A	69	GLN
1	A	163	GLN
1	A	360	ASN
1	A	383	HIS
1	B	47	HIS
1	B	60	GLN
1	B	69	GLN
1	B	170	GLN
1	B	189	HIS
1	B	268	ASN
1	B	278	ASN
1	B	360	ASN
1	B	383	HIS
1	B	428	ASN
1	B	514	ASN
1	B	526	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	AN2	A	601	-	24,29,29	1.47	4 (16%)	22,45,45	2.55	3 (13%)
2	AN2	B	601	-	24,29,29	3.41	4 (16%)	22,45,45	2.30	4 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	AN2	A	601	-	-	0/9/32/32	0/3/3/3
2	AN2	B	601	-	-	0/9/32/32	0/3/3/3

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	601	AN2	PB-O2B	-3.09	1.48	1.56
2	A	601	AN2	C8-N7	-2.40	1.30	1.34
2	A	601	AN2	PB-O1B	2.76	1.49	1.46
2	A	601	AN2	PB-O3A	3.29	1.63	1.59
2	B	601	AN2	O4'-C1'	3.31	1.46	1.41
2	A	601	AN2	O4'-C1'	3.58	1.46	1.41

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	601	AN2	PB-O3A	3.88	1.64	1.59
2	B	601	AN2	PB-O1B	15.16	1.62	1.46

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	AN2	N3-C2-N1	-9.45	121.45	128.87
2	B	601	AN2	N3-C2-N1	-8.59	122.13	128.87
2	A	601	AN2	PA-O3A-PB	-5.47	115.56	132.78
2	B	601	AN2	PA-O3A-PB	-4.51	118.60	132.78
2	B	601	AN2	O2A-PA-O3A	2.00	113.85	105.27
2	B	601	AN2	O4'-C1'-N9	2.91	113.61	108.11
2	A	601	AN2	O4'-C1'-N9	3.32	114.38	108.11

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	501/584 (85%)	-0.43	4 (0%) 87 85	26, 45, 79, 109	0
1	B	490/584 (83%)	-0.35	9 (1%) 71 66	27, 47, 83, 135	0
All	All	991/1168 (84%)	-0.39	13 (1%) 79 75	26, 46, 81, 135	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	474	TYR	3.6
1	B	491	TYR	3.2
1	B	490	ILE	3.2
1	B	136	GLU	2.9
1	B	32	GLU	2.6
1	A	471	HIS	2.5
1	B	475	CYS	2.5
1	A	390	LYS	2.4
1	A	483	GLN	2.3
1	B	28	LYS	2.3
1	A	137	ASN	2.2
1	B	27	SER	2.2
1	B	137	ASN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	AN2	B	601	27/27	0.92	0.13	-0.53	35,66,103,105	0
2	AN2	A	601	27/27	0.93	0.13	-0.67	48,69,88,89	0

6.5 Other polymers [i](#)

There are no such residues in this entry.