



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 06:51 AM GMT

PDB ID : 2YLS
Title : SNAPSHOTS OF ENZYMATIC BAEYER-VILLIGER CATALYSIS: OXY-
GEN ACTIVATION AND INTERMEDIATE STABILIZATION: REDUCED
ENZYME BOUND TO NADP
Authors : Orru, R.; Dudek, H.M.; Martinoli, C.; Torres Pazmino, D.E.; Royant, A.;
Weik, M.; Fraaije, M.W.; Mattevi, A.
Deposited on : 2011-06-06
Resolution : 2.26 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.26 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1640 (2.28-2.24)
Clashscore	102246	1095 (2.26-2.26)
Ramachandran outliers	100387	1063 (2.26-2.26)
Sidechain outliers	100360	1063 (2.26-2.26)
RSRZ outliers	91569	1647 (2.28-2.24)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	542	 85% 11% • •

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4569 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PHENYLACETONE MONOOXYGENASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	532	Total	C	N	O	S	0	0	0
			4230	2699	728	794	9			

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula: $C_{21}H_{28}N_7O_{17}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	238	Total	O	0	0
			238	238		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: PHENYLACETONE MONOOXYGENASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	107.92Å 107.92Å 106.83Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	42.81 – 2.26 42.81 – 2.26	Depositor EDS
% Data completeness (in resolution range)	99.2 (42.81-2.26) 97.8 (42.81-2.26)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.09 (at 2.27Å)	Xtriage
Refinement program	REFMAC 5.5.0110	Depositor
R, R_{free}	0.187 , 0.235 0.187 , 0.232	Depositor DCC
R_{free} test set	1640 reflections (5.17%)	DCC
Wilson B-factor (Å ²)	25.5	Xtriage
Anisotropy	0.106	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 38.3	EDS
Estimated twinning fraction	0.035 for -h,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 33527 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4569	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAP, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.05	3/4343 (0.1%)	0.97	15/5911 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	530	GLU	CB-CG	5.62	1.62	1.52
1	A	163	GLY	N-CA	5.55	1.54	1.46
1	A	79	GLU	CG-CD	5.53	1.60	1.51

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	246	ARG	NE-CZ-NH2	-16.52	112.04	120.30
1	A	246	ARG	NE-CZ-NH1	12.48	126.54	120.30
1	A	158	LEU	CA-CB-CG	9.83	137.91	115.30
1	A	525	ARG	NE-CZ-NH2	-6.84	116.88	120.30
1	A	162	PRO	C-N-CA	-6.78	108.07	122.30
1	A	392	LEU	CA-CB-CG	6.56	130.38	115.30
1	A	297	ARG	NE-CZ-NH2	-6.50	117.05	120.30
1	A	525	ARG	NE-CZ-NH1	5.96	123.28	120.30
1	A	297	ARG	CG-CD-NE	-5.93	99.35	111.80
1	A	402	ARG	NE-CZ-NH2	-5.72	117.44	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	158	LEU	CB-CG-CD2	-5.55	101.56	111.00
1	A	398	LYS	CD-CE-NZ	5.24	123.75	111.70
1	A	11	ARG	C-N-CA	5.13	134.53	121.70
1	A	513	VAL	CB-CA-C	-5.08	101.74	111.40
1	A	246	ARG	CD-NE-CZ	5.06	130.69	123.60

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	163	GLY	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4230	0	4071	51	0
2	A	53	0	31	1	0
3	A	48	0	25	0	0
4	A	238	0	0	7	0
All	All	4569	0	4127	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:159:PRO:HG2	1:A:161:PHE:CE2	1.95	1.01
1:A:87:THR:CG2	1:A:95:GLU:OE2	2.22	0.88
1:A:506:ASN:HD22	1:A:506:ASN:H	1.24	0.84
1:A:402:ARG:HD2	1:A:406:ASN:ND2	1.93	0.84
1:A:163:GLY:HA3	1:A:366:ILE:O	1.77	0.83
1:A:402:ARG:HD2	1:A:406:ASN:HD21	1.42	0.82
1:A:159:PRO:HG2	1:A:161:PHE:HE2	1.48	0.77

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:160:ASN:CB	4:A:2083:HOH:O	2.33	0.77
1:A:335:THR:OG1	1:A:498:THR:HG21	1.88	0.74
1:A:87:THR:HG23	1:A:95:GLU:OE2	1.91	0.70
1:A:87:THR:HG22	1:A:95:GLU:OE2	1.95	0.67
1:A:144:ARG:HG2	1:A:145:TYR:CE2	2.30	0.66
1:A:126:GLU:OE1	1:A:126:GLU:HA	1.95	0.66
1:A:159:PRO:CG	1:A:161:PHE:CE2	2.77	0.66
1:A:506:ASN:HD22	1:A:506:ASN:N	1.95	0.65
1:A:152:GLN:HE22	2:A:700:FAD:H5'2	1.63	0.62
1:A:163:GLY:HA3	1:A:366:ILE:HB	1.83	0.60
1:A:241:ARG:NH1	4:A:2124:HOH:O	2.32	0.60
1:A:372:ARG:HG3	1:A:372:ARG:HH11	1.67	0.59
1:A:163:GLY:CA	1:A:366:ILE:HB	2.35	0.57
1:A:159:PRO:HG3	1:A:387:THR:HG21	1.87	0.55
1:A:332:PRO:HD2	1:A:498:THR:CG2	2.37	0.54
1:A:241:ARG:HD3	4:A:2124:HOH:O	2.08	0.54
1:A:523:ARG:HA	1:A:526:GLN:HE21	1.75	0.52
1:A:506:ASN:H	1:A:506:ASN:ND2	2.01	0.51
1:A:30:TYR:CE2	1:A:451:GLU:HG2	2.47	0.49
1:A:142:ARG:HB2	1:A:142:ARG:NH1	2.28	0.48
1:A:126:GLU:CA	1:A:126:GLU:OE1	2.59	0.48
1:A:370:GLY:HA3	1:A:378:TYR:O	2.14	0.48
1:A:512:ARG:HH11	1:A:512:ARG:CG	2.27	0.47
1:A:246:ARG:HD2	4:A:2044:HOH:O	2.14	0.47
1:A:332:PRO:HD2	1:A:498:THR:HG22	1.96	0.47
1:A:79:GLU:OE1	1:A:106:LYS:HE3	2.14	0.46
1:A:142:ARG:HB2	1:A:142:ARG:CZ	2.46	0.46
1:A:226:ARG:HG3	1:A:284:GLY:HA3	1.98	0.46
1:A:525:ARG:NE	4:A:2229:HOH:O	2.31	0.45
1:A:471:ARG:NH1	1:A:473:GLU:OE1	2.43	0.44
1:A:89:ARG:HD3	1:A:90:TYR:CE2	2.52	0.44
1:A:452:GLN:HG3	1:A:525:ARG:HG3	1.98	0.44
1:A:316:THR:HG21	1:A:345:TYR:HB2	2.00	0.43
1:A:226:ARG:CG	1:A:284:GLY:HA3	2.48	0.43
1:A:23:GLY:O	1:A:28:GLY:HA3	2.20	0.42
1:A:231:ASP:HA	1:A:232:PRO:HD3	1.94	0.42
1:A:512:ARG:N	1:A:512:ARG:HD3	2.35	0.41
1:A:512:ARG:NE	4:A:2221:HOH:O	2.53	0.41
1:A:317:VAL:HB	1:A:323:ALA:HB2	2.02	0.41
1:A:346:GLU:CB	4:A:2162:HOH:O	2.68	0.41
1:A:412:LYS:NZ	1:A:479:GLU:OE2	2.50	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:506:ASN:ND2	1:A:506:ASN:N	2.64	0.40
1:A:495:TYR:HB3	1:A:514:PHE:CE1	2.56	0.40
1:A:306:ALA:O	1:A:310:ARG:HG3	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	530/542 (98%)	515 (97%)	15 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	440/453 (97%)	422 (96%)	18 (4%)	37	44

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	87	THR
1	A	126	GLU
1	A	144	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	201	VAL
1	A	237	ASP
1	A	240	LYS
1	A	242	TYR
1	A	264	SER
1	A	273	LEU
1	A	360	SER
1	A	372	ARG
1	A	392	LEU
1	A	484	GLU
1	A	498	THR
1	A	506	ASN
1	A	512	ARG
1	A	522	HIS
1	A	525	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	58	ASN
1	A	129	ASN
1	A	152	GLN
1	A	157	GLN
1	A	315	ASN
1	A	406	ASN
1	A	506	ASN
1	A	526	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FAD	A	700	-	48,58,58	1.47	6 (12%)	54,89,89	2.42	11 (20%)
3	NAP	A	701	-	42,52,52	1.75	6 (14%)	54,80,80	2.50	14 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	700	-	-	0/30/50/50	0/6/6/6
3	NAP	A	701	-	-	0/27/67/67	0/5/5/5

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	701	NAP	C7N-N7N	2.30	1.37	1.33
2	A	700	FAD	C8M-C8	2.48	1.56	1.51
3	A	701	NAP	C2N-C3N	2.54	1.42	1.39
2	A	700	FAD	C2A-N1A	2.59	1.38	1.33
2	A	700	FAD	C4-N3	2.64	1.38	1.33
3	A	701	NAP	P2B-O2B	2.76	1.68	1.60
3	A	701	NAP	P2B-O1X	3.37	1.62	1.51
2	A	700	FAD	C2A-N3A	3.91	1.39	1.32
2	A	700	FAD	C5X-N5	4.06	1.41	1.35
2	A	700	FAD	C4X-N5	4.73	1.40	1.33
3	A	701	NAP	O4B-C1B	5.39	1.48	1.41
3	A	701	NAP	O4D-C1D	5.59	1.48	1.41

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	701	NAP	N3A-C2A-N1A	-13.29	118.72	128.89
2	A	700	FAD	N3A-C2A-N1A	-10.78	120.64	128.89
2	A	700	FAD	C4-C4X-C10	-7.19	115.34	119.94
2	A	700	FAD	C4X-C4-N3	-3.90	118.25	123.59
3	A	701	NAP	O7N-C7N-N7N	-3.78	117.28	122.59
3	A	701	NAP	C3B-C2B-C1B	-3.24	96.46	102.73
2	A	700	FAD	C4A-C5A-N7A	-3.20	106.53	109.48
3	A	701	NAP	PN-O3-PA	-2.68	125.21	132.73
3	A	701	NAP	C1B-N9A-C4A	-2.59	123.03	126.94
2	A	700	FAD	C9A-C5X-N5	-2.58	118.53	122.36
3	A	701	NAP	O3X-P2B-O1X	-2.40	102.86	110.58
3	A	701	NAP	O5B-C5B-C4B	-2.27	100.76	109.12
2	A	700	FAD	O2'-C2'-C1'	-2.22	104.50	109.94
2	A	700	FAD	C4-C4X-N5	2.03	121.19	118.72
2	A	700	FAD	O4B-C1B-N9A	2.13	112.57	108.10
2	A	700	FAD	C5X-C9A-N10	2.40	119.44	117.62
3	A	701	NAP	O3X-P2B-O2X	2.40	116.53	107.38
3	A	701	NAP	C2A-N1A-C6A	2.51	123.26	118.77
3	A	701	NAP	O2X-P2B-O1X	2.53	118.73	110.58
3	A	701	NAP	O4B-C1B-N9A	2.65	113.65	108.10
3	A	701	NAP	C3N-C7N-N7N	3.33	121.47	117.82
2	A	700	FAD	C1'-N10-C9A	3.55	122.85	118.86
3	A	701	NAP	O3-PN-O5D	3.60	112.48	102.94
3	A	701	NAP	O4D-C1D-N1N	5.46	114.13	108.13
2	A	700	FAD	C4-N3-C2	6.69	121.03	115.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	700	FAD	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	532/542 (98%)	-0.59	0 100 100	13, 22, 36, 49	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	FAD	A	700	53/53	0.98	0.11	0.19	9,16,21,25	0
3	NAP	A	701	48/48	0.98	0.11	0.01	12,17,20,21	0

6.5 Other polymers [i](#)

There are no such residues in this entry.