



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 11:46 PM GMT

PDB ID : 1YMB
Title : HIGH RESOLUTION STUDY OF THE THREE-DIMENSIONAL STRUCTURE OF HORSE HEART METMYOGLOBIN
Authors : Evans, S.V.; Brayer, G.D.
Deposited on : 1993-09-27
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	5398 (1.90-1.90)
Ramachandran outliers	100387	5338 (1.90-1.90)
Sidechain outliers	100360	5339 (1.90-1.90)
RSRZ outliers	91569	4766 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	153	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 1401 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called METMYOGLOBIN.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	153	Total	C	N	O	S	0	0	0
			1199	769	210	218	2			

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: C₃₄H₃₂FeN₄O₄).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	154	Total	O	0	0
			154	154		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: METMYOGLOBIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	64.27Å 28.92Å 35.93Å 90.00° 107.10° 90.00°	Depositor
Resolution (Å)	(Not available) – 1.90 21.06 – 1.90	Depositor EDS
% Data completeness (in resolution range)	(Not available) ((Not available)-1.90) 85.4 (21.06-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.14 (at 1.90Å)	Xtriage
Refinement program	PROLSQ	Depositor
R, R_{free}	0.155 , (Not available) 0.222 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	12.9	Xtriage
Anisotropy	0.211	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 91.3	EDS
Estimated twinning fraction	0.045 for -h-l,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Outliers	0 of 10275 reflections	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	1401	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 13.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.39	2/1226 (0.2%)	2.06	41/1645 (2.5%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	85	GLU	CD-OE1	-6.40	1.18	1.25
1	A	18	GLU	CD-OE1	5.25	1.31	1.25

All (41) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	31	ARG	NE-CZ-NH1	14.78	127.69	120.30
1	A	31	ARG	NE-CZ-NH2	-12.37	114.12	120.30
1	A	103	TYR	CB-CG-CD1	10.14	127.09	121.00
1	A	102	LYS	CB-CG-CD	9.15	135.39	111.60
1	A	86	LEU	CB-CA-C	8.19	125.76	110.20
1	A	140	ASN	CA-CB-CG	7.61	130.14	113.40
1	A	103	TYR	CB-CG-CD2	-7.46	116.53	121.00
1	A	106	PHE	CB-CG-CD1	7.25	125.88	120.80
1	A	148	GLU	CA-CB-CG	7.06	128.94	113.40
1	A	41	GLU	CA-CB-CG	6.80	128.37	113.40
1	A	27	GLU	CG-CD-OE2	6.79	131.88	118.30
1	A	29	LEU	CB-CA-C	6.43	122.41	110.20
1	A	147	LYS	O-C-N	6.37	132.89	122.70
1	A	132	THR	CA-CB-CG2	6.27	121.18	112.40
1	A	44	ASP	C-N-CA	6.20	137.20	121.70
1	A	38	GLU	OE1-CD-OE2	-6.17	115.89	123.30
1	A	10	VAL	CA-CB-CG2	6.17	120.16	110.90
1	A	58	SER	N-CA-CB	-6.12	101.33	110.50
1	A	55	MET	CG-SD-CE	6.06	109.90	100.20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	99	ILE	CB-CG1-CD1	6.04	130.82	113.90
1	A	106	PHE	CB-CG-CD2	-6.03	116.58	120.80
1	A	22	ALA	CB-CA-C	5.98	119.06	110.10
1	A	37	PRO	N-CD-CG	-5.98	94.23	103.20
1	A	146	TYR	CB-CG-CD1	5.78	124.47	121.00
1	A	69	LEU	CB-CG-CD1	-5.62	101.44	111.00
1	A	145	LYS	CA-CB-CG	5.62	125.76	113.40
1	A	93	HIS	CA-CB-CG	-5.60	104.08	113.60
1	A	10	VAL	CG1-CB-CG2	-5.59	101.96	110.90
1	A	59	GLU	OE1-CD-OE2	5.52	129.93	123.30
1	A	51	THR	CA-C-O	5.32	131.27	120.10
1	A	141	ASP	CB-CG-OD1	5.31	123.08	118.30
1	A	44	ASP	CB-CG-OD1	5.30	123.07	118.30
1	A	139	ARG	NE-CZ-NH2	-5.30	117.65	120.30
1	A	145	LYS	CB-CG-CD	5.27	125.30	111.60
1	A	46	PHE	CB-CG-CD1	5.22	124.45	120.80
1	A	27	GLU	CG-CD-OE1	-5.17	107.96	118.30
1	A	11	LEU	CB-CA-C	5.15	119.99	110.20
1	A	113	HIS	CA-C-O	5.10	130.80	120.10
1	A	41	GLU	CG-CD-OE1	5.03	128.37	118.30
1	A	105	GLU	CA-C-O	5.03	130.67	120.10
1	A	91	GLN	CG-CD-OE1	-5.02	111.56	121.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1199	0	1211	87	0
2	A	5	0	0	1	0
3	A	43	0	30	3	0
4	A	154	0	0	36	1
All	All	1401	0	1241	88	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 35.

All (88) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1:GLY:HA2	4:A:276:HOH:O	1.61	1.00
1:A:98:LYS:HG3	4:A:225:HOH:O	1.76	0.85
1:A:145:LYS:HE3	4:A:218:HOH:O	1.76	0.85
1:A:21:ILE:HD12	1:A:66:THR:HG23	1.59	0.84
1:A:45:LYS:HE3	4:A:173:HOH:O	1.87	0.74
1:A:78:LYS:CG	4:A:261:HOH:O	2.36	0.73
1:A:78:LYS:HG2	4:A:261:HOH:O	1.86	0.73
1:A:13:VAL:HG13	1:A:16:LYS:HZ2	1.53	0.73
1:A:9:GLN:HG3	4:A:165:HOH:O	1.89	0.73
1:A:44:ASP:HA	1:A:47:LYS:HD2	1.71	0.70
1:A:98:LYS:HB2	4:A:237:HOH:O	1.93	0.69
1:A:98:LYS:HD3	1:A:151:PHE:HE1	1.58	0.69
1:A:21:ILE:CD1	1:A:66:THR:HG23	2.23	0.68
1:A:98:LYS:HD3	1:A:151:PHE:CE1	2.29	0.68
1:A:3:SER:HB2	4:A:284:HOH:O	1.95	0.66
1:A:98:LYS:CB	4:A:237:HOH:O	2.44	0.65
1:A:98:LYS:HE3	4:A:292:HOH:O	1.95	0.65
1:A:96:LYS:HE3	4:A:248:HOH:O	1.97	0.65
1:A:98:LYS:CE	4:A:292:HOH:O	2.46	0.63
1:A:75:ILE:O	1:A:78:LYS:HB2	1.99	0.63
1:A:119:HIS:N	1:A:120:PRO:HD3	2.14	0.63
1:A:123:PHE:HA	4:A:234:HOH:O	1.98	0.62
1:A:101:ILE:HG12	1:A:146:TYR:CD2	2.35	0.62
1:A:96:LYS:CE	4:A:248:HOH:O	2.48	0.62
1:A:118:LYS:C	1:A:120:PRO:HD3	2.21	0.61
1:A:3:SER:OG	1:A:6:GLU:HG3	2.02	0.59
1:A:126:ASP:HB2	4:A:227:HOH:O	2.02	0.59
1:A:30:ILE:HD13	1:A:56:LYS:HG3	1.83	0.59
1:A:131:MET:HA	4:A:264:HOH:O	2.03	0.58
1:A:78:LYS:HE2	4:A:158:HOH:O	2.02	0.58
1:A:99:ILE:HD12	3:A:154:HEM:HAC	1.86	0.57
1:A:101:ILE:H	1:A:152:GLN:HE22	1.52	0.57
1:A:98:LYS:NZ	4:A:292:HOH:O	2.36	0.57
1:A:83:GLU:CD	1:A:83:GLU:H	2.09	0.56
1:A:88:PRO:HD3	4:A:199:HOH:O	2.05	0.55
1:A:78:LYS:CE	4:A:261:HOH:O	2.53	0.55
1:A:82:HIS:HE1	1:A:137:LEU:HD23	1.70	0.55
1:A:51:THR:OG1	1:A:54:GLU:HB2	2.07	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:86:LEU:HD21	1:A:141:ASP:HB3	1.88	0.54
1:A:13:VAL:HG21	1:A:127:ALA:HB1	1.89	0.53
3:A:154:HEM:CGA	4:A:283:HOH:O	2.55	0.53
1:A:95:THR:HA	1:A:98:LYS:HE2	1.90	0.53
1:A:9:GLN:HG2	1:A:127:ALA:HA	1.91	0.52
1:A:2:LEU:HD12	1:A:133:LYS:HZ2	1.75	0.51
1:A:118:LYS:HG3	4:A:279:HOH:O	2.09	0.51
1:A:10:VAL:HG22	4:A:264:HOH:O	2.09	0.51
1:A:44:ASP:CA	1:A:47:LYS:HD2	2.42	0.49
1:A:3:SER:HB3	4:A:270:HOH:O	2.12	0.49
1:A:73:GLY:O	1:A:77:LYS:HG3	2.12	0.49
1:A:82:HIS:HE1	1:A:137:LEU:CD2	2.25	0.49
1:A:16:LYS:HE3	4:A:229:HOH:O	2.13	0.49
1:A:14:TRP:O	1:A:18:GLU:HG3	2.12	0.49
1:A:60:ASP:HB2	2:A:155:SO4:O4	2.13	0.49
1:A:101:ILE:H	1:A:152:GLN:NE2	2.10	0.49
1:A:146:TYR:HD2	1:A:152:GLN:HE21	1.59	0.49
1:A:131:MET:HE2	4:A:264:HOH:O	2.13	0.49
1:A:33:PHE:HE2	1:A:61:LEU:HD11	1.78	0.48
1:A:146:TYR:HD2	1:A:152:GLN:NE2	2.12	0.47
1:A:99:ILE:CD1	3:A:154:HEM:HAC	2.44	0.46
1:A:2:LEU:HA	1:A:6:GLU:OE1	2.14	0.46
1:A:116:HIS:NE2	1:A:128:GLN:NE2	2.64	0.46
1:A:52:GLU:HA	1:A:55:MET:HE3	1.97	0.46
1:A:101:ILE:HD13	1:A:104:LEU:HD12	1.98	0.46
1:A:95:THR:HG21	4:A:190:HOH:O	2.16	0.45
1:A:51:THR:O	1:A:55:MET:HE2	2.15	0.45
1:A:52:GLU:O	1:A:55:MET:HB2	2.17	0.45
1:A:107:ILE:O	1:A:110:ALA:HB3	2.17	0.45
1:A:112:ILE:HG12	4:A:198:HOH:O	2.17	0.45
1:A:14:TRP:CH2	1:A:17:VAL:HG11	2.52	0.44
1:A:123:PHE:HD1	4:A:234:HOH:O	1.99	0.44
1:A:133:LYS:HD2	4:A:161:HOH:O	2.17	0.44
1:A:117:SER:O	1:A:120:PRO:HG3	2.17	0.44
1:A:52:GLU:HA	1:A:55:MET:CE	2.47	0.44
1:A:151:PHE:C	1:A:152:GLN:HG3	2.39	0.43
1:A:10:VAL:HA	4:A:264:HOH:O	2.19	0.43
1:A:10:VAL:HG13	4:A:264:HOH:O	2.19	0.42
1:A:115:LEU:HD12	1:A:123:PHE:CZ	2.54	0.42
1:A:112:ILE:HD11	1:A:135:LEU:HD12	2.00	0.42
1:A:57:ALA:HB1	4:A:170:HOH:O	2.19	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:133:LYS:HD2	4:A:222:HOH:O	2.20	0.41
1:A:33:PHE:CE2	1:A:61:LEU:HD11	2.54	0.41
1:A:24:HIS:O	1:A:28:VAL:HG23	2.20	0.41
1:A:27:GLU:CD	1:A:118:LYS:HE3	2.40	0.41
1:A:146:TYR:CD2	1:A:152:GLN:NE2	2.88	0.41
1:A:2:LEU:HD11	1:A:133:LYS:HG2	2.02	0.41
1:A:87:LYS:NZ	4:A:282:HOH:O	2.53	0.40
1:A:6:GLU:OE1	1:A:133:LYS:HE3	2.20	0.40
1:A:102:LYS:HE3	1:A:106:PHE:CE1	2.57	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:168:HOH:O	4:A:309:HOH:O[2_656]	1.98	0.22

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	151/153 (99%)	137 (91%)	13 (9%)	1 (1%)	26 14

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	152	GLN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	123/123 (100%)	110 (89%)	13 (11%)	8 3

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9	GLN
1	A	27	GLU
1	A	41	GLU
1	A	50	LYS
1	A	54	GLU
1	A	78	LYS
1	A	83	GLU
1	A	87	LYS
1	A	118	LYS
1	A	133	LYS
1	A	140	ASN
1	A	145	LYS
1	A	152	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	26	GLN
1	A	82	HIS
1	A	128	GLN
1	A	140	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	HEM	A	154	1,4	30,50,50	2.20	6 (20%)	24,82,82	3.21	12 (50%)
2	SO4	A	155	-	4,4,4	1.25	1 (25%)	6,6,6	0.75	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HEM	A	154	1,4	-	0/10/54/54	0/0/8/8
2	SO4	A	155	-	-	0/0/0/0	0/0/0/0

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	154	HEM	C2D-C3D	-7.18	1.33	1.54
3	A	154	HEM	C3D-C4D	-5.02	1.45	1.51
3	A	154	HEM	C3B-C4B	-4.80	1.47	1.51
3	A	154	HEM	C2C-C1C	-2.28	1.48	1.52
2	A	155	SO4	O2-S	2.17	1.54	1.47
3	A	154	HEM	C4C-NC	2.44	1.39	1.36
3	A	154	HEM	CHD-C4C	2.50	1.42	1.36

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	154	HEM	CMA-C3A-C4A	-3.80	122.07	128.36
3	A	154	HEM	CAA-C2A-C1A	-3.52	123.18	127.01
3	A	154	HEM	C4B-CHC-C1C	-3.49	119.98	125.82
3	A	154	HEM	C3B-CAB-CBB	-2.53	120.58	124.46
3	A	154	HEM	CAD-C3D-C4D	2.65	121.81	112.47
3	A	154	HEM	CMA-C3A-C2A	2.68	130.84	125.24
3	A	154	HEM	CMD-C2D-C3D	2.77	126.61	114.35
3	A	154	HEM	C2D-C3D-C4D	3.09	106.74	101.50
3	A	154	HEM	CMB-C2B-C3B	4.17	126.95	116.53
3	A	154	HEM	CMC-C2C-C3C	5.87	131.19	116.53
3	A	154	HEM	CAD-C3D-C2D	6.32	131.38	113.22
3	A	154	HEM	C3C-CAC-CBC	8.13	136.93	124.46

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	154	HEM	3	0
2	A	155	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	153/153 (100%)	0.34	6 (3%)	43 47	5, 12, 26, 34	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	153	GLY	8.2
1	A	1	GLY	4.0
1	A	152	GLN	2.8
1	A	151	PHE	2.3
1	A	49	LEU	2.1
1	A	84	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	SO4	A	155	5/5	0.91	0.16	1.14	45,45,46,46	0
3	HEM	A	154	43/43	0.93	0.13	0.64	3,6,9,11	0

6.5 Other polymers [i](#)

There are no such residues in this entry.