



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 06:52 AM GMT

PDB ID : 2YNM
Title : Structure of the ADP \times AlF₃-Stabilized Transition State of the Nitrogenase-like Dark-Operative Protochlorophyllide Oxidoreductase Complex from *Prochlorococcus marinus* with Its Substrate Protochlorophyllide a
Authors : Krausze, J.; Lange, C.; Heinz, D.W.; Moser, J.
Deposited on : 2012-10-16
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3939 (2.10-2.10)
Clashscore	102246	4460 (2.10-2.10)
Ramachandran outliers	100387	4413 (2.10-2.10)
Sidechain outliers	100360	4414 (2.10-2.10)
RSRZ outliers	91569	3948 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	301	 2% 79% 10% 11%
1	B	301	 3% 81% 8% 11%
2	C	426	 6% 90% 7% 3%
3	D	530	 7% 84% 8% 8%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
10	GOL	A	1303	-	-	-	X
4	1PE	A	1296	-	-	-	X
5	EPE	D	1529	-	-	-	X

2 Entry composition

There are 13 unique types of molecules in this entry. The entry contains 11825 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LIGHT-INDEPENDENT PROTOCHLOROPHYLLIDE REDUCTASE IRON-SULFUR ATP-BINDING PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	267	Total	C	N	O	S	0	0	0
			2056	1304	347	393	12			
1	B	268	Total	C	N	O	S	0	1	0
			2069	1311	349	397	12			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	GLY	-	EXPRESSION TAG	UNP Q7VD39
A	-3	PRO	-	EXPRESSION TAG	UNP Q7VD39
A	-2	LEU	-	EXPRESSION TAG	UNP Q7VD39
A	-1	GLY	-	EXPRESSION TAG	UNP Q7VD39
A	0	SER	-	EXPRESSION TAG	UNP Q7VD39
B	-4	GLY	-	EXPRESSION TAG	UNP Q7VD39
B	-3	PRO	-	EXPRESSION TAG	UNP Q7VD39
B	-2	LEU	-	EXPRESSION TAG	UNP Q7VD39
B	-1	GLY	-	EXPRESSION TAG	UNP Q7VD39
B	0	SER	-	EXPRESSION TAG	UNP Q7VD39

- Molecule 2 is a protein called LIGHT-INDEPENDENT PROTOCHLOROPHYLLIDE REDUCTASE SUBUNIT N.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	417	Total	C	N	O	S	0	0	0
			3238	2061	564	599	14			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-7	GLY	-	EXPRESSION TAG	UNP Q7VD37
C	-6	PRO	-	EXPRESSION TAG	UNP Q7VD37

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-5	LEU	-	EXPRESSION TAG	UNP Q7VD37
C	-4	GLY	-	EXPRESSION TAG	UNP Q7VD37
C	-3	SER	-	EXPRESSION TAG	UNP Q7VD37
C	-2	PRO	-	EXPRESSION TAG	UNP Q7VD37
C	-1	GLU	-	EXPRESSION TAG	UNP Q7VD37
C	0	PHE	-	EXPRESSION TAG	UNP Q7VD37

- Molecule 3 is a protein called LIGHT-INDEPENDENT PROTOCHLOROPHYLLIDE REDUCTASE SUBUNIT B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	488	Total	C	N	O	S	0	2	0
			3854	2449	661	723	21			

- Molecule 4 is PENTAETHYLENE GLYCOL (three-letter code: 1PE) (formula: C₁₀H₂₂O₆).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			16	10	6		

- Molecule 5 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (three-letter code: EPE) (formula: C₈H₁₈N₂O₄S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	S	0	0
			15	8	2	4	1		
5	A	1	Total	C	N	O	S	0	0
			15	8	2	4	1		
5	B	1	Total	C	N	O	S	0	0
			15	8	2	4	1		
5	D	1	Total	C	N	O	S	0	0
			15	8	2	4	1		

- Molecule 6 is ALUMINUM FLUORIDE (three-letter code: AF3) (formula: AlF_3).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total 4	Al 1	F 3	0	0
6	B	1	Total 4	Al 1	F 3	0	0

- Molecule 7 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $\text{C}_{10}\text{H}_{15}\text{N}_5\text{O}_{10}\text{P}_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	A	1	Total 27	C 10	N 5	O 10	P 2	0	0
7	B	1	Total 27	C 10	N 5	O 10	P 2	0	0

- Molecule 8 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	B	1	Total Mg 1 1	0	0
8	A	1	Total Mg 1 1	0	0

- Molecule 9 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe_4S_4).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	A	1	Total	Fe	S	0	0
			8	4	4		
9	C	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 10 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	A	1	Total	C	O	0	0
			6	3	3		
10	C	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	D	1	Total	C	O	0	0
			6	3	3		
10	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 11 is PROTOCHLOROPHYLLIDE (three-letter code: PMR) (formula: $C_{35}H_{32}MgN_4O_5$).





Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
13	D	127	Total	O	0	0
			127	127		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: LIGHT-INDEPENDENT PROTOCHLOROPHYLLIDE REDUCTASE IRON-SULFUR ATP-BINDING PROTEIN




• Molecule 1: LIGHT-INDEPENDENT PROTOCHLOROPHYLLIDE REDUCTASE IRON-SULFUR ATP-BINDING PROTEIN




• Molecule 2: LIGHT-INDEPENDENT PROTOCHLOROPHYLLIDE REDUCTASE SUBUNIT N



• Molecule 3: LIGHT-INDEPENDENT PROTOCHLOROPHYLLIDE REDUCTASE SUBUNIT B





4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	308.40Å 74.11Å 74.23Å 90.00° 91.24° 90.00°	Depositor
Resolution (Å)	34.51 – 2.10 34.51 – 2.10	Depositor EDS
% Data completeness (in resolution range)	83.1 (34.51-2.10) 83.1 (34.51-2.10)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.74 (at 2.10Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE: 1.8.1_1168)	Depositor
R, R_{free}	0.199 , 0.233 0.208 , 0.240	Depositor DCC
R_{free} test set	4030 reflections (5.21%)	DCC
Wilson B-factor (Å ²)	23.6	Xtriage
Anisotropy	0.160	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 52.7	EDS
Estimated twinning fraction	0.022 for -h,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	1 of 81370 reflections (0.001%)	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	11825	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.69% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MG, ADP, PMR, SF4, 1PE, EPE, K, AF3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.22	0/2091	0.40	0/2821
1	B	0.23	0/2104	0.40	0/2838
2	C	0.22	0/3300	0.42	0/4469
3	D	0.22	0/3942	0.41	0/5335
All	All	0.22	0/11437	0.41	0/15463

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2056	0	2056	20	0
1	B	2069	0	2064	16	0
2	C	3238	0	3321	19	0
3	D	3854	0	3808	27	0
4	A	16	0	22	3	0
5	A	30	0	34	0	0
5	B	15	0	17	0	0
5	D	15	0	17	2	0
6	A	4	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	4	0	0	0	0
7	A	27	0	12	0	0
7	B	27	0	12	0	0
8	A	1	0	0	0	0
8	B	1	0	0	0	0
9	A	8	0	0	0	0
9	C	8	0	0	0	0
10	A	6	0	8	2	0
10	C	6	0	8	0	0
10	D	12	0	16	0	0
11	C	45	0	31	5	0
12	D	1	0	0	0	0
13	A	89	0	0	5	0
13	B	63	0	0	0	0
13	C	103	0	0	0	0
13	D	127	0	0	1	0
All	All	11825	0	11426	78	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 78 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:D:399:PRO:HG3	5:D:1529:EPE:H82	1.71	0.71
11:C:1414:PMR:HMAA	3:D:45:MET:HB3	1.73	0.70
1:A:202:ARG:HH22	4:A:1296:1PE:H121	1.56	0.69
3:D:147:GLU:HG3	3:D:148:ILE:HG13	1.79	0.65
3:D:514:ARG:O	3:D:515:GLU:HG2	1.99	0.62

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	265/301 (88%)	258 (97%)	7 (3%)	0	100	100
1	B	267/301 (89%)	259 (97%)	8 (3%)	0	100	100
2	C	415/426 (97%)	402 (97%)	13 (3%)	0	100	100
3	D	482/530 (91%)	468 (97%)	14 (3%)	0	100	100
All	All	1429/1558 (92%)	1387 (97%)	42 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	224/252 (89%)	222 (99%)	2 (1%)	84	89
1	B	225/252 (89%)	224 (100%)	1 (0%)	93	96
2	C	354/361 (98%)	352 (99%)	2 (1%)	90	94
3	D	412/443 (93%)	410 (100%)	2 (0%)	92	95
All	All	1215/1308 (93%)	1208 (99%)	7 (1%)	90	94

5 of 7 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	124	ASN
3	D	293	TYR
2	C	204	ILE
1	A	166	GLN
3	D	144	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	C	343	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 19 ligands modelled in this entry, 3 are monoatomic - leaving 16 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	1PE	A	1296	-	15,15,15	0.45	0	14,14,14	0.45	0
5	EPE	A	1297	-	14,15,15	0.37	0	18,20,20	1.81	5 (27%)
5	EPE	A	1298	-	14,15,15	0.36	0	18,20,20	1.94	6 (33%)
6	AF3	A	1299	13,7	0,3,3	0.00	-	0,3,3	0.00	-
7	ADP	A	1300	8,6	22,29,29	1.01	1 (4%)	27,45,45	1.84	5 (18%)
9	SF4	A	1302	1	0,12,12	0.00	-	0,24,24	0.00	-
10	GOL	A	1303	-	5,5,5	0.21	0	5,5,5	0.36	0
5	EPE	B	1296	-	14,15,15	0.41	0	18,20,20	1.87	5 (27%)
6	AF3	B	1298	8,13,7	0,3,3	0.00	-	0,3,3	0.00	-
7	ADP	B	1299	8,6	22,29,29	0.99	1 (4%)	27,45,45	1.84	4 (14%)
10	GOL	C	1412	-	5,5,5	0.17	0	5,5,5	0.32	0
9	SF4	C	1413	3,2	0,12,12	0.00	-	0,24,24	0.00	-

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
11	PMR	C	1414	13	36,53,53	2.44	15 (41%)	38,89,89	3.00	21 (55%)
5	EPE	D	1529	-	14,15,15	0.37	0	18,20,20	1.79	5 (27%)
10	GOL	D	1530	-	5,5,5	0.21	0	5,5,5	0.31	0
10	GOL	D	1532	-	5,5,5	0.18	0	5,5,5	0.20	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	1PE	A	1296	-	-	0/13/13/13	0/0/0/0
5	EPE	A	1297	-	-	0/9/19/19	0/1/1/1
5	EPE	A	1298	-	-	0/9/19/19	0/1/1/1
6	AF3	A	1299	13,7	-	0/0/0/0	0/0/0/0
7	ADP	A	1300	8,6	-	0/12/32/32	0/3/3/3
9	SF4	A	1302	1	-	0/0/48/48	0/6/5/5
10	GOL	A	1303	-	-	0/4/4/4	0/0/0/0
5	EPE	B	1296	-	-	0/9/19/19	0/1/1/1
6	AF3	B	1298	8,13,7	-	0/0/0/0	0/0/0/0
7	ADP	B	1299	8,6	-	0/12/32/32	0/3/3/3
10	GOL	C	1412	-	-	0/4/4/4	0/0/0/0
9	SF4	C	1413	3,2	-	0/0/48/48	0/6/5/5
11	PMR	C	1414	13	-	0/11/111/111	0/0/9/9
5	EPE	D	1529	-	-	0/9/19/19	0/1/1/1
10	GOL	D	1530	-	-	0/4/4/4	0/0/0/0
10	GOL	D	1532	-	-	0/4/4/4	0/0/0/0

The worst 5 of 17 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
11	C	1414	PMR	C3D-C4D	-4.23	1.35	1.41
11	C	1414	PMR	C4C-C3C	2.07	1.48	1.45
11	C	1414	PMR	C1A-C2A	2.13	1.49	1.45
11	C	1414	PMR	C4D-CHA	2.19	1.47	1.45
11	C	1414	PMR	CHC-C1C	2.43	1.46	1.41

The worst 5 of 51 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	1300	ADP	N3-C2-N1	-6.98	123.55	128.89
7	B	1299	ADP	N3-C2-N1	-6.61	123.83	128.89

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
11	C	1414	PMR	C4A-C3A-C2A	-3.50	102.73	106.91
11	C	1414	PMR	C4C-C3C-C2C	-3.29	101.60	106.94
11	C	1414	PMR	C1C-C2C-C3C	-3.29	102.98	106.91

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1296	1PE	3	0
10	A	1303	GOL	2	0
11	C	1414	PMR	5	0
5	D	1529	EPE	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	267/301 (88%)	0.05	7 (2%) 59 66	19, 35, 63, 78	0
1	B	268/301 (89%)	0.17	8 (2%) 54 62	18, 37, 64, 83	0
2	C	417/426 (97%)	0.21	25 (5%) 25 33	19, 42, 71, 96	0
3	D	488/530 (92%)	0.29	39 (7%) 15 21	22, 35, 83, 133	0
All	All	1440/1558 (92%)	0.20	79 (5%) 29 37	18, 38, 72, 133	0

The worst 5 of 79 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	D	512	GLY	6.7
3	D	513	CYS	6.5
3	D	514	ARG	6.1
3	D	515	GLU	5.7
3	D	479	ILE	5.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	EPE	D	1529	15/15	0.77	0.31	7.93	60,66,74,75	0
10	GOL	A	1303	6/6	0.60	0.44	7.13	63,69,71,71	0
4	1PE	A	1296	16/16	0.74	0.26	3.82	49,64,71,72	0
5	EPE	A	1297	15/15	0.93	0.19	1.51	32,46,53,54	0
5	EPE	A	1298	15/15	0.91	0.15	1.47	39,41,51,52	0
10	GOL	D	1532	6/6	0.88	0.17	1.23	38,42,45,46	0
11	PMR	C	1414	45/45	0.93	0.15	0.88	14,23,28,39	0
5	EPE	B	1296	15/15	0.92	0.17	0.85	35,58,60,60	0
6	AF3	B	1298	4/4	0.96	0.19	0.40	25,27,29,44	0
7	ADP	B	1299	27/27	0.97	0.14	-0.19	20,29,38,39	0
6	AF3	A	1299	4/4	0.97	0.17	-0.23	17,27,32,39	0
10	GOL	C	1412	6/6	0.97	0.12	-0.34	25,33,36,38	0
10	GOL	D	1530	6/6	0.97	0.13	-0.35	20,30,33,35	0
7	ADP	A	1300	27/27	0.96	0.12	-0.43	20,31,34,38	0
9	SF4	A	1302	8/8	0.99	0.11	-1.10	17,19,21,22	0
9	SF4	C	1413	8/8	0.99	0.11	-1.21	19,22,25,26	0
8	MG	A	1301	1/1	0.99	0.11	-1.54	26,26,26,26	0
8	MG	B	1297	1/1	0.92	0.11	-1.85	34,34,34,34	0
12	K	D	1531	1/1	0.73	0.10	-	66,66,66,66	0

6.5 Other polymers [i](#)

There are no such residues in this entry.