



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 06:57 AM GMT

PDB ID : 2YXL
Title : Crystal Structure of PH0851
Authors : Hikida, Y.; Kuratani, M.; Bessho, Y.; Ishii, R.; Sekine, S.; Yokoyama, S.;
RIKEN Structural Genomics/Proteomics Initiative (RSGI)
Deposited on : 2007-04-26
Resolution : 2.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4549 (2.58-2.50)
Clashscore	102246	5292 (2.58-2.50)
Ramachandran outliers	100387	5194 (2.58-2.50)
Sidechain outliers	100360	5196 (2.58-2.50)
RSRZ outliers	91569	4561 (2.58-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	450	

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 3659 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 450aa long hypothetical fmu protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	443	Total	C	N	O	S	0	0	0
			3574	2307	614	642	11			

- Molecule 2 is SINEFUNGIN (three-letter code: SFG) (formula: $C_{15}H_{23}N_7O_5$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			27	15	7	5		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	58	Total	O	0	0
			58	58		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 450aa long hypothetical fmu protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	116.62Å 116.62Å 93.68Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	38.18 – 2.55 38.17 – 2.55	Depositor EDS
% Data completeness (in resolution range)	99.3 (38.18-2.55) 99.3 (38.17-2.55)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	10.28 (at 2.54Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.212 , 0.246 0.210 , 0.245	Depositor DCC
R_{free} test set	1204 reflections (5.24%)	DCC
Wilson B-factor (Å ²)	56.9	Xtriage
Anisotropy	0.267	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 53.1	EDS
Estimated twinning fraction	0.024 for -h,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 24202 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3659	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.37% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SFG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.36	0/3653	0.53	1/4933 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	358	GLU	N-CA-C	5.53	125.92	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	408	HIS	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3574	0	3680	37	0
2	A	27	0	22	1	0
3	A	58	0	0	1	0
All	All	3659	0	3702	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:190:ARG:HH22	1:A:275:THR:HG23	1.44	0.82
1:A:183:ARG:HA	1:A:184:HIS:HB3	1.64	0.79
1:A:183:ARG:CA	1:A:184:HIS:HB3	2.19	0.72
1:A:257:PRO:HA	1:A:333:LYS:HD2	1.75	0.69
1:A:357:ARG:HB3	1:A:359:ASP:O	1.93	0.68
1:A:213:VAL:HG13	1:A:223:LYS:HB2	1.76	0.68
1:A:322:ALA:HB2	1:A:325:ILE:HB	1.79	0.64
1:A:271:GLY:HA3	1:A:300:LEU:HD13	1.84	0.59
1:A:32:GLN:H	1:A:32:GLN:HE21	1.56	0.54
1:A:358:GLU:O	1:A:361:ILE:HB	2.10	0.51
1:A:372:LEU:HG	1:A:404:PHE:CD2	2.46	0.51
1:A:417:LYS:O	1:A:417:LYS:HE3	2.11	0.50
1:A:217:ARG:HD3	1:A:419:PRO:HB2	1.93	0.50
1:A:358:GLU:N	1:A:359:ASP:C	2.67	0.48
1:A:183:ARG:HD3	3:A:5754:HOH:O	2.12	0.48
1:A:304:VAL:HG13	1:A:309:ILE:HB	1.95	0.48
1:A:183:ARG:HA	1:A:184:HIS:CB	2.35	0.46
1:A:23:ARG:C	1:A:23:ARG:HD3	2.37	0.46
1:A:79:PRO:HA	1:A:82:LEU:HD22	1.99	0.45
1:A:177:PHE:HA	1:A:180:VAL:HG22	1.98	0.45
1:A:32:GLN:H	1:A:32:GLN:NE2	2.14	0.45
1:A:30:PRO:HB2	1:A:32:GLN:NE2	2.32	0.45
1:A:294:LYS:HD3	1:A:316:VAL:HG11	1.99	0.44
1:A:266:LEU:HG	1:A:319:ALA:HB1	1.99	0.44
1:A:384:ARG:HG2	1:A:447:GLU:HB3	2.00	0.44
1:A:318:ASP:OD1	1:A:320:ARG:HD3	2.18	0.44
1:A:368:GLN:O	1:A:372:LEU:HB2	2.18	0.43
1:A:179:SER:HA	1:A:182:LYS:HE3	2.00	0.42
2:A:5748:SFG:H4'	2:A:5748:SFG:HG2	1.92	0.42
1:A:360:LYS:HB2	1:A:363:GLU:HB3	2.03	0.41
1:A:297:MET:HG2	1:A:316:VAL:HG22	2.02	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:20:GLU:OE2	1:A:42:HIS:HE1	2.03	0.41
1:A:287:ILE:HB	1:A:312:VAL:HG22	2.02	0.41
1:A:83:ASP:HA	1:A:84:PRO:HD3	1.96	0.41
1:A:110:LEU:O	1:A:114:ALA:HB3	2.20	0.41
1:A:322:ALA:HA	1:A:323:PRO:C	2.41	0.40
1:A:274:THR:HA	1:A:277:LEU:HD12	2.03	0.40
1:A:407:VAL:HG13	1:A:408:HIS:CD2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	441/450 (98%)	424 (96%)	13 (3%)	4 (1%)	21	36

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	409	PRO
1	A	328	GLU
1	A	184	HIS
1	A	322	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	388/396 (98%)	359 (92%)	29 (8%)	17	30

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	18	ILE
1	A	23	ARG
1	A	32	GLN
1	A	80	LEU
1	A	82	LEU
1	A	107	ILE
1	A	134	LEU
1	A	136	LYS
1	A	141	LYS
1	A	161	LEU
1	A	194	LEU
1	A	218	VAL
1	A	229	ASN
1	A	255	LEU
1	A	262	THR
1	A	274	THR
1	A	294	LYS
1	A	298	LYS
1	A	305	LYS
1	A	317	LYS
1	A	324	GLU
1	A	333	LYS
1	A	358	GLU
1	A	385	LEU
1	A	409	PRO
1	A	410	GLU
1	A	412	LYS
1	A	416	LEU
1	A	417	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	32	GLN
1	A	42	HIS
1	A	105	GLN
1	A	109	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	146	ASN
1	A	192	ASN
1	A	229	ASN
1	A	276	HIS
1	A	349	ASN
1	A	362	ASN
1	A	397	ASN
1	A	408	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SFG	A	5748	-	21,29,29	0.90	1 (4%)	19,42,42	2.52	4 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means

no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SFG	A	5748	-	-	0/9/33/33	0/3/3/3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	5748	SFG	O4'-C1'	2.59	1.44	1.41

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	5748	SFG	N3-C2-N1	-9.54	121.59	128.89
2	A	5748	SFG	C4'-O4'-C1'	-3.77	105.57	109.72
2	A	5748	SFG	O4'-C4'-C5'	-2.30	106.00	109.36
2	A	5748	SFG	C4-C5-N7	-2.15	107.50	109.48

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	5748	SFG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	443/450 (98%)	0.40	26 (5%)	26 30	22, 36, 46, 58	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	144	PRO	4.5
1	A	183	ARG	4.3
1	A	6	LYS	4.2
1	A	321	LYS	3.9
1	A	298	LYS	3.8
1	A	316	VAL	3.7
1	A	373	GLU	3.5
1	A	44	VAL	3.5
1	A	145	LYS	3.5
1	A	315	LEU	3.5
1	A	305	LYS	3.4
1	A	328	GLU	3.4
1	A	141	LYS	3.4
1	A	43	ASP	3.3
1	A	405	LEU	3.2
1	A	142	PRO	3.1
1	A	407	VAL	2.9
1	A	288	TYR	2.7
1	A	313	LYS	2.6
1	A	324	GLU	2.6
1	A	294	LYS	2.5
1	A	359	ASP	2.4
1	A	143	ASN	2.3
1	A	425	LEU	2.3
1	A	326	ILE	2.1
1	A	413	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	SFG	A	5748	27/27	0.86	0.19	0.02	75,78,79,79	0

6.5 Other polymers [i](#)

There are no such residues in this entry.