



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 11:54 PM GMT

PDB ID : 1YZW
Title : The 2.1Å Crystal Structure of the Far-red Fluorescent Protein HcRed: Inherent Conformational Flexibility of the Chromophore
Authors : Wilmann, P.G.; Petersen, J.; Pettikiriachchi, A.; Buckle, A.M.; Devenish, R.J.; Prescott, M.; Rossjohn, J.
Deposited on : 2005-02-28
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3939 (2.10-2.10)
Clashscore	102246	4460 (2.10-2.10)
Ramachandran outliers	100387	4413 (2.10-2.10)
Sidechain outliers	100360	4414 (2.10-2.10)
RSRZ outliers	91569	3948 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	225	<div> <div>13%</div> <div>92%</div> <div>6%</div> </div>
1	B	225	<div> <div>11%</div> <div>89%</div> <div>8%</div> </div>
1	C	225	<div> <div>8%</div> <div>94%</div> <div>• •</div> </div>
1	D	225	<div> <div>5%</div> <div>96%</div> <div>• •</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PEG	B	1001	-	-	-	X

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 7562 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GFP-like non-fluorescent chromoprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	221	Total	C	N	O	S	0	0	0
			1773	1128	304	327	14			
1	B	221	Total	C	N	O	S	0	1	0
			1781	1135	304	328	14			
1	C	221	Total	C	N	O	S	0	1	0
			1781	1135	304	328	14			
1	D	221	Total	C	N	O	S	0	0	0
			1773	1128	304	327	14			

There are 36 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	37	ALA	THR	SEE REMARK 999	UNP Q95W85
A	64	CRU	GLU	CHROMOPHORE	UNP Q95W85
A	64	CRU	TYR	CHROMOPHORE	UNP Q95W85
A	64	CRU	GLY	CHROMOPHORE	UNP Q95W85
A	123	HIS	LEU	SEE REMARK 999	UNP Q95W85
A	144	SER	CYS	SEE REMARK 999	UNP Q95W85
A	169	HIS	ARG	SEE REMARK 999	UNP Q95W85
A	174	HIS	LEU	SEE REMARK 999	UNP Q95W85
A	202	LEU	PRO	SEE REMARK 999	UNP Q95W85
B	37	ALA	THR	SEE REMARK 999	UNP Q95W85
B	64	CRU	GLU	CHROMOPHORE	UNP Q95W85
B	64	CRU	TYR	CHROMOPHORE	UNP Q95W85
B	64	CRU	GLY	CHROMOPHORE	UNP Q95W85
B	123	HIS	LEU	SEE REMARK 999	UNP Q95W85
B	144	SER	CYS	SEE REMARK 999	UNP Q95W85
B	169	HIS	ARG	SEE REMARK 999	UNP Q95W85
B	174	HIS	LEU	SEE REMARK 999	UNP Q95W85
B	202	LEU	PRO	SEE REMARK 999	UNP Q95W85
C	37	ALA	THR	SEE REMARK 999	UNP Q95W85
C	64	CRU	GLU	CHROMOPHORE	UNP Q95W85
C	64	CRU	TYR	CHROMOPHORE	UNP Q95W85

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	64	CRU	GLY	CHROMOPHORE	UNP Q95W85
C	123	HIS	LEU	SEE REMARK 999	UNP Q95W85
C	144	SER	CYS	SEE REMARK 999	UNP Q95W85
C	169	HIS	ARG	SEE REMARK 999	UNP Q95W85
C	174	HIS	LEU	SEE REMARK 999	UNP Q95W85
C	202	LEU	PRO	SEE REMARK 999	UNP Q95W85
D	37	ALA	THR	SEE REMARK 999	UNP Q95W85
D	64	CRU	GLU	CHROMOPHORE	UNP Q95W85
D	64	CRU	TYR	CHROMOPHORE	UNP Q95W85
D	64	CRU	GLY	CHROMOPHORE	UNP Q95W85
D	123	HIS	LEU	SEE REMARK 999	UNP Q95W85
D	144	SER	CYS	SEE REMARK 999	UNP Q95W85
D	169	HIS	ARG	SEE REMARK 999	UNP Q95W85
D	174	HIS	LEU	SEE REMARK 999	UNP Q95W85
D	202	LEU	PRO	SEE REMARK 999	UNP Q95W85

- Molecule 2 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			7	4	3		
2	C	1	Total	C	O	0	0
			7	4	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	105	Total 105	O 105	0	0
3	B	119	Total 119	O 119	0	0
3	C	105	Total 105	O 105	0	0
3	D	111	Total 111	O 111	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: GFP-like non-fluorescent chromoprotein




- Molecule 1: GFP-like non-fluorescent chromoprotein



- Molecule 1: GFP-like non-fluorescent chromoprotein



- Molecule 1: GFP-like non-fluorescent chromoprotein



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	76.22Å 120.73Å 118.47Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	84.51 – 2.10 19.89 – 2.10	Depositor EDS
% Data completeness (in resolution range)	98.0 (84.51-2.10) 98.1 (19.89-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.67 (at 2.09Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.223 , 0.266 0.240 , 0.280	Depositor DCC
R_{free} test set	3196 reflections (5.33%)	DCC
Wilson B-factor (Å ²)	36.2	Xtriage
Anisotropy	0.565	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 49.4	EDS
Estimated twinning fraction	0.000 for -h,l,k	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Outliers	0 of 63182 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7562	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 37.18 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 4.4621e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, CRU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.33	0/1796	0.52	0/2423
1	B	0.38	0/1796	0.53	0/2423
1	C	0.37	0/1796	0.52	0/2423
1	D	0.34	0/1796	0.54	0/2423
All	All	0.35	0/7184	0.53	0/9692

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	C	0	1
All	All	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	63	CYS	Mainchain
1	C	63	CYS	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1773	0	1703	6	0
1	B	1781	0	1704	15	0
1	C	1781	0	1704	7	0
1	D	1773	0	1703	4	0
2	B	7	0	10	0	0
2	C	7	0	10	0	0
3	A	105	0	0	1	0
3	B	119	0	0	0	0
3	C	105	0	0	0	0
3	D	111	0	0	0	0
All	All	7562	0	6834	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:83:GLN:HE22	1:D:188:THR:H	1.35	0.74
1:B:12:ILE:HD11	1:B:70:PHE:CZ	2.29	0.67
1:C:83:GLN:HE22	1:C:188:THR:H	1.41	0.65
1:A:127:PHE:CD1	1:A:133:VAL:HG11	2.33	0.64
1:A:200:GLN:HE22	1:B:226:LYS:H	1.48	0.61
1:A:226:LYS:H	1:B:200:GLN:HE22	1.49	0.60
1:C:60:ALA:HB1	1:C:199:LEU:HD21	1.82	0.59
1:C:12:ILE:HD11	1:C:70:PHE:CZ	2.37	0.59
1:B:64[B]:CRU:HD1	1:B:199:LEU:HD22	1.85	0.59
1:C:12:ILE:HD11	1:C:70:PHE:CE1	2.39	0.58
1:B:64[A]:CRU:HE1	1:B:159:ASN:HD22	1.72	0.54
1:C:226:LYS:H	1:D:200:GLN:HE22	1.56	0.53
1:B:10:MET:SD	1:B:114:ASN:ND2	2.83	0.52
1:B:40:GLN:HE22	1:B:67:SER:HB3	1.76	0.51
1:B:40:GLN:HE22	1:B:67:SER:CB	2.26	0.49
1:B:27:CYS:HB3	1:B:46:VAL:HA	1.95	0.48
1:B:64[A]:CRU:CE1	1:B:159:ASN:HD22	2.26	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:60:ALA:HB3	1:B:61:PRO:HD3	1.94	0.48
1:B:64[B]:CRU:HE1	1:B:199:LEU:HB3	1.97	0.46
1:B:64[B]:CRU:HE1	1:B:199:LEU:CB	2.48	0.44
1:A:165:VAL:HG22	3:A:320:HOH:O	2.17	0.43
1:A:147:VAL:HG13	1:A:192:PHE:CZ	2.53	0.43
1:C:60:ALA:CB	1:C:199:LEU:HD21	2.48	0.43
1:B:64[B]:CRU:CD2	1:B:197:ILE:HD13	2.50	0.42
1:D:103:LEU:HD21	1:D:122:VAL:HG13	2.02	0.42
1:A:127:PHE:HD1	1:A:133:VAL:HG11	1.82	0.41
1:B:35:PRO:HA	1:B:70:PHE:HA	2.03	0.41
1:D:83:GLN:NE2	1:D:188:THR:H	2.12	0.40
1:C:60:ALA:HB3	1:C:61:PRO:HD3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	218/225 (97%)	212 (97%)	6 (3%)	0	100	100
1	B	218/225 (97%)	211 (97%)	7 (3%)	0	100	100
1	C	218/225 (97%)	213 (98%)	5 (2%)	0	100	100
1	D	218/225 (97%)	212 (97%)	6 (3%)	0	100	100
All	All	872/900 (97%)	848 (97%)	24 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	188/190 (99%)	183 (97%)	5 (3%)	52	56
1	B	188/190 (99%)	184 (98%)	4 (2%)	61	66
1	C	188/190 (99%)	187 (100%)	1 (0%)	92	95
1	D	188/190 (99%)	187 (100%)	1 (0%)	92	95
All	All	752/760 (99%)	741 (98%)	11 (2%)	72	78

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	13	LYS
1	A	27	CYS
1	A	43	ARG
1	A	151	GLU
1	A	199	LEU
1	B	5	LEU
1	B	27	CYS
1	B	43	ARG
1	B	76	GLU
1	C	185	ARG
1	D	199	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	83	GLN
1	A	200	GLN
1	B	83	GLN
1	B	114	ASN
1	B	200	GLN
1	C	83	GLN
1	C	114	ASN
1	C	126	ASN
1	C	173	HIS
1	D	83	GLN
1	D	200	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

6 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CRU	A	64	1	21,25,26	2.96	11 (52%)	22,34,36	2.37	10 (45%)
1	CRU	B	64[A]	-	21,25,26	2.94	11 (52%)	22,34,36	2.83	9 (40%)
1	CRU	B	64[B]	-	21,25,26	2.95	11 (52%)	22,34,36	2.86	11 (50%)
1	CRU	C	64[A]	-	21,25,26	2.90	10 (47%)	22,34,36	2.68	8 (36%)
1	CRU	C	64[B]	-	21,25,26	2.95	10 (47%)	22,34,36	2.72	10 (45%)
1	CRU	D	64	1	21,25,26	2.94	10 (47%)	22,34,36	2.81	8 (36%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CRU	A	64	1	-	0/8/32/33	0/2/2/2
1	CRU	B	64[A]	-	-	0/8/32/33	0/2/2/2
1	CRU	B	64[B]	-	-	0/8/32/33	0/2/2/2
1	CRU	C	64[A]	-	-	0/8/32/33	0/2/2/2
1	CRU	C	64[B]	-	-	0/8/32/33	0/2/2/2
1	CRU	D	64	1	-	0/8/32/33	0/2/2/2

All (63) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	64[A]	CRU	CA3-C	-8.21	1.18	1.49
1	C	64[B]	CRU	CA3-C	-8.21	1.18	1.49

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	64	CRU	CA3-C	-8.17	1.18	1.49
1	A	64	CRU	CA3-C	-8.08	1.18	1.49
1	B	64[A]	CRU	CA3-C	-7.98	1.19	1.49
1	B	64[B]	CRU	CA3-C	-7.98	1.19	1.49
1	A	64	CRU	O-C	2.02	1.32	1.19
1	B	64[A]	CRU	O-C	2.05	1.33	1.19
1	B	64[B]	CRU	O-C	2.05	1.33	1.19
1	B	64[B]	CRU	CE1-CZ	2.24	1.43	1.38
1	D	64	CRU	CD2-CE2	2.28	1.42	1.38
1	C	64[B]	CRU	CD2-CE2	2.31	1.42	1.38
1	C	64[A]	CRU	CD2-CE2	2.31	1.42	1.38
1	B	64[B]	CRU	CD2-CE2	2.33	1.42	1.38
1	A	64	CRU	CD1-CG2	2.34	1.43	1.39
1	B	64[B]	CRU	CD1-CG2	2.36	1.43	1.39
1	D	64	CRU	C1-N3	2.39	1.43	1.38
1	C	64[A]	CRU	CE1-CZ	2.41	1.43	1.38
1	C	64[A]	CRU	CD1-CG2	2.42	1.44	1.39
1	B	64[A]	CRU	CD2-CE2	2.43	1.43	1.38
1	B	64[A]	CRU	CD1-CG2	2.43	1.44	1.39
1	A	64	CRU	CD2-CE2	2.45	1.43	1.38
1	B	64[B]	CRU	CE2-CZ	2.46	1.43	1.38
1	D	64	CRU	CD1-CG2	2.48	1.44	1.39
1	B	64[A]	CRU	CE1-CZ	2.48	1.43	1.38
1	C	64[B]	CRU	CD1-CG2	2.49	1.44	1.39
1	C	64[A]	CRU	C1-N3	2.50	1.43	1.38
1	C	64[B]	CRU	C1-N3	2.50	1.43	1.38
1	D	64	CRU	CE1-CZ	2.51	1.43	1.38
1	C	64[B]	CRU	CE2-CZ	2.51	1.44	1.38
1	C	64[A]	CRU	CE2-CZ	2.52	1.44	1.38
1	C	64[B]	CRU	CE1-CZ	2.55	1.44	1.38
1	D	64	CRU	CE2-CZ	2.56	1.44	1.38
1	A	64	CRU	CE1-CZ	2.57	1.44	1.38
1	B	64[A]	CRU	CE2-CZ	2.58	1.44	1.38
1	A	64	CRU	CE2-CZ	2.59	1.44	1.38
1	B	64[A]	CRU	C1-N3	2.67	1.44	1.38
1	B	64[B]	CRU	C1-N3	2.67	1.44	1.38
1	C	64[B]	CRU	CD2-CG2	2.73	1.44	1.39
1	B	64[B]	CRU	CD2-CG2	2.74	1.44	1.39
1	A	64	CRU	C1-N3	2.81	1.44	1.38
1	B	64[A]	CRU	CE1-CD1	2.85	1.43	1.38
1	D	64	CRU	CD2-CG2	2.92	1.45	1.39
1	C	64[A]	CRU	CD2-CG2	2.93	1.45	1.39

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	64	CRU	CE1-CD1	2.93	1.44	1.38
1	B	64[A]	CRU	CD2-CG2	2.94	1.45	1.39
1	C	64[A]	CRU	CE1-CD1	2.94	1.44	1.38
1	D	64	CRU	CE1-CD1	3.00	1.44	1.38
1	A	64	CRU	CD2-CG2	3.12	1.45	1.39
1	C	64[B]	CRU	CE1-CD1	3.15	1.44	1.38
1	B	64[B]	CRU	CE1-CD1	3.29	1.44	1.38
1	B	64[A]	CRU	CB2-CA2	4.53	1.39	1.35
1	C	64[A]	CRU	CB2-CA2	4.54	1.39	1.35
1	A	64	CRU	CB2-CA2	4.55	1.39	1.35
1	D	64	CRU	CB2-CA2	4.68	1.39	1.35
1	B	64[B]	CRU	CB2-CA2	4.85	1.39	1.35
1	C	64[B]	CRU	CB2-CA2	5.01	1.39	1.35
1	C	64[A]	CRU	CA3-N3	5.58	1.56	1.47
1	C	64[B]	CRU	CA3-N3	5.58	1.56	1.47
1	D	64	CRU	CA3-N3	5.82	1.57	1.47
1	B	64[A]	CRU	CA3-N3	6.02	1.57	1.47
1	B	64[B]	CRU	CA3-N3	6.02	1.57	1.47
1	A	64	CRU	CA3-N3	6.07	1.57	1.47

All (56) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	64	CRU	CB2-CA2-N2	-6.69	116.70	128.67
1	B	64[A]	CRU	CB2-CA2-N2	-6.23	117.54	128.67
1	C	64[A]	CRU	CB2-CA2-N2	-5.90	118.12	128.67
1	A	64	CRU	CB2-CA2-N2	-5.16	119.44	128.67
1	B	64[A]	CRU	O2-C2-CA2	-4.88	128.31	130.95
1	B	64[B]	CRU	O2-C2-CA2	-4.88	128.31	130.95
1	D	64	CRU	O2-C2-CA2	-3.57	129.02	130.95
1	B	64[A]	CRU	CB1-CA1-N	-3.55	118.42	124.94
1	B	64[B]	CRU	CB1-CA1-N	-3.55	118.42	124.94
1	B	64[B]	CRU	CE2-CZ-CE1	-3.47	114.85	119.79
1	C	64[A]	CRU	CB1-CA1-N	-3.22	119.02	124.94
1	C	64[B]	CRU	CB1-CA1-N	-3.22	119.02	124.94
1	A	64	CRU	CB1-CA1-N	-3.10	119.24	124.94
1	C	64[B]	CRU	CE2-CZ-CE1	-3.10	115.38	119.79
1	D	64	CRU	CB1-CA1-N	-3.07	119.29	124.94
1	C	64[A]	CRU	O2-C2-CA2	-2.99	129.33	130.95
1	C	64[B]	CRU	O2-C2-CA2	-2.99	129.33	130.95
1	A	64	CRU	CE2-CZ-CE1	-2.64	116.02	119.79
1	B	64[A]	CRU	CE2-CZ-CE1	-2.56	116.15	119.79

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	64[A]	CRU	CE2-CZ-CE1	-2.51	116.21	119.79
1	D	64	CRU	CE2-CZ-CE1	-2.47	116.28	119.79
1	B	64[B]	CRU	CD2-CG2-CD1	-2.37	114.03	117.64
1	C	64[A]	CRU	CA3-N3-C2	-2.34	120.19	123.99
1	C	64[B]	CRU	CA3-N3-C2	-2.34	120.19	123.99
1	A	64	CRU	N3-C1-N2	-2.31	110.64	113.26
1	C	64[B]	CRU	CD2-CG2-CD1	-2.19	114.30	117.64
1	A	64	CRU	CA3-N3-C2	-2.14	120.51	123.99
1	B	64[A]	CRU	N3-C1-N2	-2.11	110.87	113.26
1	B	64[B]	CRU	N3-C1-N2	-2.11	110.87	113.26
1	D	64	CRU	C-CA3-N3	2.01	117.40	113.00
1	C	64[B]	CRU	CE2-CD2-CG2	2.04	123.84	121.29
1	B	64[B]	CRU	CE2-CD2-CG2	2.05	123.85	121.29
1	A	64	CRU	CD1-CE1-CZ	2.11	122.31	119.87
1	B	64[A]	CRU	C-CA3-N3	2.14	117.68	113.00
1	B	64[B]	CRU	C-CA3-N3	2.14	117.68	113.00
1	D	64	CRU	CB1-CG1-CD3	2.34	117.03	112.75
1	A	64	CRU	CA2-C2-N3	2.35	104.58	103.40
1	A	64	CRU	C-CA3-N3	2.43	118.33	113.00
1	C	64[B]	CRU	CD1-CE1-CZ	2.43	122.68	119.87
1	B	64[A]	CRU	CB1-CG1-CD3	2.45	117.24	112.75
1	B	64[B]	CRU	CB1-CG1-CD3	2.45	117.24	112.75
1	A	64	CRU	CB1-CG1-CD3	2.62	117.55	112.75
1	B	64[B]	CRU	CD1-CE1-CZ	2.86	123.18	119.87
1	C	64[A]	CRU	CA2-C2-N3	3.05	104.93	103.40
1	C	64[B]	CRU	CA2-C2-N3	3.05	104.93	103.40
1	B	64[A]	CRU	CA2-C2-N3	3.15	104.98	103.40
1	B	64[B]	CRU	CA2-C2-N3	3.15	104.98	103.40
1	D	64	CRU	CA2-C2-N3	3.30	105.06	103.40
1	C	64[A]	CRU	CB1-CG1-CD3	3.84	119.78	112.75
1	C	64[B]	CRU	CB1-CG1-CD3	3.84	119.78	112.75
1	A	64	CRU	CB2-CA2-C2	6.00	131.15	122.36
1	C	64[A]	CRU	CB2-CA2-C2	6.68	132.14	122.36
1	B	64[A]	CRU	CB2-CA2-C2	7.09	132.74	122.36
1	D	64	CRU	CB2-CA2-C2	7.63	133.54	122.36
1	C	64[B]	CRU	CG2-CB2-CA2	8.41	141.14	130.22
1	B	64[B]	CRU	CG2-CB2-CA2	8.48	141.22	130.22

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	64[A]	CRU	2	0
1	B	64[B]	CRU	4	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PEG	B	1001	-	6,6,6	0.43	0	5,5,5	0.27	0
2	PEG	C	1002	-	6,6,6	0.47	0	5,5,5	0.22	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PEG	B	1001	-	-	0/4/4/4	0/0/0/0
2	PEG	C	1002	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	220/225 (97%)	0.72	29 (13%) 4 6	44, 53, 63, 68	0
1	B	220/225 (97%)	0.75	25 (11%) 7 9	43, 51, 61, 66	0
1	C	220/225 (97%)	0.49	18 (8%) 14 20	40, 50, 60, 71	0
1	D	220/225 (97%)	0.45	12 (5%) 29 37	38, 51, 62, 69	0
All	All	880/900 (97%)	0.60	84 (9%) 10 14	38, 51, 62, 71	0

All (84) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	4	GLY	4.8
1	A	203	ARG	4.7
1	B	4	GLY	4.6
1	A	209	TYR	4.4
1	D	203	ARG	4.0
1	B	69	THR	4.0
1	A	59	LEU	4.0
1	A	11	ARG	4.0
1	A	130	ASP	3.9
1	C	152	ASN	3.8
1	A	167	ASP	3.8
1	C	44	ILE	3.6
1	A	137	LYS	3.4
1	C	58	ILE	3.3
1	C	8	GLU	3.3
1	D	167	ASP	3.2
1	C	69	THR	3.2
1	B	5	LEU	3.2
1	B	8	GLU	3.2
1	A	129	ALA	3.1
1	B	186	ALA	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	155	LEU	3.1
1	D	59	LEU	3.1
1	D	209	TYR	3.0
1	B	59	LEU	3.0
1	B	152	ASN	3.0
1	A	58	ILE	2.9
1	C	76	GLU	2.9
1	B	188	THR	2.9
1	D	69	THR	2.9
1	A	168	ARG	2.9
1	A	148	VAL	2.9
1	B	20	VAL	2.9
1	A	81	PHE	2.8
1	D	130	ASP	2.8
1	C	81	PHE	2.8
1	C	56	PHE	2.8
1	B	44	ILE	2.8
1	B	184	VAL	2.8
1	C	55	ALA	2.8
1	A	205	LYS	2.7
1	A	69	THR	2.7
1	D	58	ILE	2.7
1	D	23	HIS	2.7
1	A	152	ASN	2.7
1	C	147	VAL	2.6
1	A	45	HIS	2.6
1	A	67	SER	2.6
1	B	148	VAL	2.6
1	B	187	LEU	2.5
1	A	206	LYS	2.5
1	B	81	PHE	2.5
1	B	29	GLY	2.4
1	B	56	PHE	2.4
1	B	116	LEU	2.4
1	C	71	VAL	2.4
1	B	124	GLY	2.3
1	A	56	PHE	2.3
1	A	76	GLU	2.3
1	B	89	PHE	2.3
1	A	60	ALA	2.3
1	A	165	VAL	2.3
1	A	44	ILE	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	59	LEU	2.3
1	A	24	TYR	2.3
1	B	58	ILE	2.3
1	C	114	ASN	2.3
1	D	212	LEU	2.2
1	D	60	ALA	2.2
1	C	116	LEU	2.2
1	D	11	ARG	2.2
1	B	167	ASP	2.2
1	B	76	GLU	2.2
1	A	70	PHE	2.1
1	C	130	ASP	2.1
1	C	167	ASP	2.1
1	B	162	ALA	2.1
1	C	60	ALA	2.1
1	B	109	THR	2.1
1	A	202	LEU	2.1
1	A	63	CYS	2.0
1	B	130	ASP	2.0
1	D	8	GLU	2.0
1	A	125	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	CRU	C	64[A]	24/25	0.84	0.24	-	45,48,54,58	8
1	CRU	B	64[A]	24/25	0.83	0.24	-	47,52,55,59	8
1	CRU	C	64[B]	24/25	0.84	0.24	-	46,48,54,58	8
1	CRU	A	64	24/25	0.78	0.24	-	53,56,58,59	0
1	CRU	D	64	24/25	0.83	0.22	-	52,56,58,61	0
1	CRU	B	64[B]	24/25	0.83	0.24	-	47,52,55,59	8

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	PEG	B	1001	7/7	0.83	0.18	6.04	57,58,62,63	0
2	PEG	C	1002	7/7	0.84	0.15	0.20	61,64,66,66	0

6.5 Other polymers [i](#)

There are no such residues in this entry.