



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 11:55 PM GMT

PDB ID : 1Z47  
Title : Structure of the ATPase subunit CysA of the putative sulfate ATP-binding cassette (ABC) transporter from Alicyclobacillus acidocaldarius  
Authors : Scheffel, F.; Demmer, U.; Warkentin, E.; Huelsmann, A.; Schneider, E.; Ermler, U.  
Deposited on : 2005-03-15  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*



The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	4755 (1.90-1.90)
Clashscore	102246	5398 (1.90-1.90)
Ramachandran outliers	100387	5338 (1.90-1.90)
Sidechain outliers	100360	5339 (1.90-1.90)
RSRZ outliers	91569	4766 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	355	
1	B	355	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5778 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called putative ABC-transporter ATP-binding protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	345	Total	C	N	O	S	0	0	0
			2734	1729	500	497	8			
1	B	342	Total	C	N	O	S	0	0	0
			2716	1719	497	493	7			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	EXPRESSION TAG	UNP Q9RHZ7
A	2	ARG	-	EXPRESSION TAG	UNP Q9RHZ7
A	3	GLY	-	EXPRESSION TAG	UNP Q9RHZ7
A	4	HIS	-	EXPRESSION TAG	UNP Q9RHZ7
A	5	HIS	-	EXPRESSION TAG	UNP Q9RHZ7
A	6	HIS	-	EXPRESSION TAG	UNP Q9RHZ7
A	7	HIS	-	EXPRESSION TAG	UNP Q9RHZ7
A	8	HIS	-	EXPRESSION TAG	UNP Q9RHZ7
A	9	HIS	-	EXPRESSION TAG	UNP Q9RHZ7
A	10	HIS	-	EXPRESSION TAG	UNP Q9RHZ7
A	11	GLY	-	EXPRESSION TAG	UNP Q9RHZ7
A	12	SER	-	EXPRESSION TAG	UNP Q9RHZ7
B	1	MET	-	EXPRESSION TAG	UNP Q9RHZ7
B	2	ARG	-	EXPRESSION TAG	UNP Q9RHZ7
B	3	GLY	-	EXPRESSION TAG	UNP Q9RHZ7
B	4	HIS	-	EXPRESSION TAG	UNP Q9RHZ7
B	5	HIS	-	EXPRESSION TAG	UNP Q9RHZ7
B	6	HIS	-	EXPRESSION TAG	UNP Q9RHZ7
B	7	HIS	-	EXPRESSION TAG	UNP Q9RHZ7
B	8	HIS	-	EXPRESSION TAG	UNP Q9RHZ7
B	9	HIS	-	EXPRESSION TAG	UNP Q9RHZ7
B	10	HIS	-	EXPRESSION TAG	UNP Q9RHZ7
B	11	GLY	-	EXPRESSION TAG	UNP Q9RHZ7
B	12	SER	-	EXPRESSION TAG	UNP Q9RHZ7

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total 1	Cl 1	0	0
2	A	1	Total 1	Cl 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	222	Total 222	O 222	0	0
3	B	104	Total 104	O 104	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: putative ABC-transporter ATP-binding protein



- Molecule 1: putative ABC-transporter ATP-binding protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	154.81Å 56.31Å 91.62Å 90.00° 91.72° 90.00°	Depositor
Resolution (Å)	19.97 – 1.90 19.97 – 1.90	Depositor EDS
% Data completeness (in resolution range)	92.9 (19.97-1.90) 93.0 (19.97-1.90)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.43 (at 1.90Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.225 , 0.277 0.225 , 0.275	Depositor DCC
$R_{free}$ test set	2944 reflections (5.07%)	DCC
Wilson B-factor (Å <sup>2</sup> )	30.5	Xtriage
Anisotropy	0.240	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 54.6	EDS
Estimated twinning fraction	0.014 for -h,-k,l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 58076 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	5778	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.39% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.64	0/2798	0.76	0/3792
1	B	0.47	0/2780	0.68	1/3769 (0.0%)
All	All	0.56	0/5578	0.73	1/7561 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	250	ASN	N-CA-C	-5.08	97.29	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2734	0	2708	37	0
1	B	2716	0	2691	76	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	222	0	0	3	0
3	B	104	0	0	3	0
All	All	5778	0	5399	113	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (113) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:12:SER:HB2	1:A:40:GLU:HB2	1.48	0.94
1:B:89:VAL:HG12	1:B:166:VAL:HB	1.53	0.91
1:B:320:LYS:HD3	1:B:320:LYS:H	1.45	0.82
1:B:161:ALA:HB3	1:B:162:PRO:HD3	1.61	0.82
1:A:297:ARG:HG3	1:A:297:ARG:HH11	1.48	0.79
1:B:320:LYS:HD3	1:B:320:LYS:N	1.97	0.78
1:B:133:MET:HG2	1:B:187:PHE:HZ	1.51	0.76
1:B:88:ASN:HB3	1:B:165:GLN:HB2	1.69	0.74
1:A:127:ARG:HH11	1:A:127:ARG:HG2	1.51	0.72
1:B:197:VAL:HG12	1:B:198:THR:H	1.56	0.69
1:A:178:THR:HG22	1:A:181:ARG:NH2	2.09	0.68
1:A:127:ARG:HG2	1:A:127:ARG:NH1	2.12	0.64
1:A:229:THR:OG1	1:A:232:GLU:HG3	1.98	0.64
1:B:197:VAL:HG12	1:B:198:THR:N	2.14	0.63
1:A:129:LEU:HD11	1:A:161:ALA:HA	1.81	0.63
1:B:39:ARG:HD2	3:B:591:HOH:O	1.97	0.63
1:A:96:TYR:CE2	1:A:152:GLN:HG3	2.34	0.62
1:A:182:ARG:HG2	1:A:182:ARG:HH11	1.65	0.62
1:B:129:LEU:HD11	1:B:161:ALA:HA	1.81	0.61
1:B:66:ARG:HG3	1:B:67:PRO:HD2	1.82	0.61
1:B:260:ARG:HD2	1:B:267:ALA:HB1	1.82	0.61
1:B:133:MET:HG2	1:B:187:PHE:CZ	2.34	0.60
1:A:58:LEU:HD22	1:A:202:VAL:HG12	1.83	0.59
1:B:136:GLU:CD	1:B:136:GLU:H	2.06	0.59
1:B:319:THR:HG22	1:B:345:MET:HE1	1.85	0.59
1:B:28:ALA:O	1:B:29:ARG:HB2	2.03	0.58
1:B:176:ILE:CG2	1:B:181:ARG:HB2	2.32	0.58
1:B:77:LYS:O	1:B:79:VAL:HG13	2.04	0.58
1:A:12:SER:O	1:A:40:GLU:N	2.36	0.58
1:B:14:THR:O	1:B:74:ILE:HA	2.04	0.57
1:B:263:VAL:HG21	1:B:268:LEU:HD22	1.86	0.57
1:B:110:PHE:O	1:B:113:ARG:HB3	2.04	0.57
1:B:301:ALA:HB2	1:B:345:MET:HE1	1.85	0.57
1:B:133:MET:HE1	1:B:156:LEU:HB3	1.85	0.56
1:B:88:ASN:OD1	1:B:165:GLN:HG3	2.05	0.56
1:B:173:PHE:O	1:B:176:ILE:CG2	2.54	0.56
1:B:320:LYS:H	1:B:320:LYS:CD	2.17	0.55

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:82:LEU:HD22	1:B:86:LYS:HG2	1.88	0.55
1:B:20:VAL:HG13	1:B:67:PRO:HB3	1.88	0.55
1:B:185:ARG:HG3	1:B:185:ARG:HH11	1.71	0.55
1:A:189:ARG:HB2	1:A:212:VAL:CG2	2.37	0.55
1:B:163:ARG:N	1:B:164:PRO:HD3	2.21	0.54
1:A:189:ARG:HB2	1:A:212:VAL:HG23	1.88	0.54
1:A:178:THR:HG22	1:A:181:ARG:HH22	1.71	0.54
1:A:182:ARG:HH11	1:A:182:ARG:CG	2.21	0.54
1:B:56:THR:HG22	1:B:60:LEU:CD2	2.38	0.53
1:A:13:MET:HG3	1:A:37:GLN:HB3	1.90	0.52
1:A:102:MET:HG2	1:A:106:ASP:CB	2.39	0.52
1:A:102:MET:HG2	1:A:106:ASP:HB2	1.91	0.52
1:B:176:ILE:HG23	1:B:181:ARG:HB2	1.92	0.51
1:B:60:LEU:HD12	1:B:67:PRO:HD3	1.91	0.51
1:A:178:THR:HG23	3:A:749:HOH:O	2.09	0.51
1:B:113:ARG:HH11	1:B:113:ARG:HG3	1.77	0.50
1:B:173:PHE:O	1:B:176:ILE:HG22	2.12	0.50
1:A:189:ARG:HD2	1:A:193:ASP:OD2	2.12	0.50
1:A:39:ARG:NH2	3:A:527:HOH:O	2.44	0.49
1:B:112:LEU:HD11	1:B:125:ARG:HB3	1.95	0.49
1:B:261:ILE:CD1	1:B:282:VAL:HG22	2.42	0.49
1:B:174:ALA:O	1:B:175:ALA:HB3	2.13	0.49
1:A:297:ARG:NH1	1:A:297:ARG:HG3	2.19	0.48
1:B:68:THR:O	1:B:69:LYS:HG3	2.13	0.48
1:B:88:ASN:O	1:B:165:GLN:HB2	2.14	0.48
1:B:94:GLN:OE1	1:B:174:ALA:HB3	2.13	0.48
1:B:178:THR:HG22	3:B:687:HOH:O	2.13	0.47
1:A:39:ARG:O	1:A:42:GLU:HG2	2.14	0.47
1:B:112:LEU:CD1	1:B:125:ARG:HB3	2.44	0.47
1:B:261:ILE:HD12	1:B:282:VAL:HG21	1.96	0.47
1:B:171:GLU:O	1:B:171:GLU:HG3	2.14	0.47
1:B:173:PHE:HB2	1:B:181:ARG:NH1	2.29	0.47
1:A:178:THR:HG21	1:A:309:LYS:O	2.15	0.47
1:B:105:TYR:OH	1:B:123:ASP:OD2	2.28	0.47
1:B:115:LYS:HB2	1:B:117:VAL:HG23	1.96	0.47
1:B:293:PRO:HD3	1:B:336:ARG:HH21	1.80	0.47
1:B:261:ILE:HD12	1:B:282:VAL:CG2	2.45	0.47
1:B:185:ARG:CG	1:B:185:ARG:HH11	2.29	0.46
1:B:123:ASP:O	1:B:127:ARG:HG3	2.15	0.46
1:B:28:ALA:O	1:B:29:ARG:CB	2.64	0.46
1:B:283:VAL:HB	1:B:351:PHE:HB2	1.98	0.46

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:295:SER:OG	1:A:298:GLU:HG2	2.16	0.46
1:B:219:LEU:HD22	1:B:219:LEU:N	2.31	0.45
1:B:185:ARG:NE	3:B:426:HOH:O	2.28	0.45
1:B:136:GLU:CD	1:B:136:GLU:N	2.69	0.45
1:A:234:TYR:OH	1:A:287:LYS:HE2	2.16	0.45
1:B:114:GLU:O	1:B:114:GLU:HG2	2.16	0.45
1:B:88:ASN:CG	1:B:165:GLN:HG3	2.37	0.45
1:A:260:ARG:HD3	1:A:267:ALA:HB1	1.98	0.45
1:A:163:ARG:HD3	3:A:685:HOH:O	2.18	0.44
1:A:182:ARG:NH1	1:A:182:ARG:CG	2.78	0.44
1:A:149:GLY:O	1:A:153:ARG:HG3	2.19	0.43
1:B:111:GLY:HA3	1:B:162:PRO:HG3	2.00	0.43
1:A:161:ALA:HB3	1:A:162:PRO:HD3	2.01	0.43
1:B:128:GLU:HA	1:B:131:ARG:HH21	1.83	0.43
1:A:12:SER:HB2	1:A:40:GLU:CB	2.34	0.43
1:B:211:GLU:HG3	1:B:211:GLU:O	2.17	0.43
1:B:112:LEU:HG	1:B:161:ALA:HB1	2.00	0.43
1:A:207:GLU:O	1:A:211:GLU:HG2	2.19	0.43
1:B:66:ARG:HD2	1:B:80:THR:O	2.19	0.42
1:B:133:MET:HE3	1:B:156:LEU:HD23	2.01	0.42
1:A:135:LEU:HD13	1:A:154:VAL:HG22	2.01	0.42
1:A:14:THR:O	1:A:74:ILE:HA	2.19	0.42
1:A:12:SER:O	1:A:39:ARG:HA	2.20	0.42
1:B:72:VAL:CG1	1:B:80:THR:HA	2.49	0.42
1:B:176:ILE:HG21	1:B:181:ARG:HB2	2.01	0.42
1:A:147:SER:OG	1:A:150:GLN:HG3	2.20	0.42
1:B:19:GLY:O	1:B:70:GLY:HA3	2.20	0.42
1:B:171:GLU:N	1:B:172:PRO:CD	2.83	0.41
1:B:334:ARG:HG3	1:B:335:HIS:CD2	2.55	0.41
1:B:41:GLY:HA2	1:B:197:VAL:O	2.20	0.41
1:B:319:THR:HG22	1:B:345:MET:CE	2.48	0.41
1:B:304:VAL:O	1:B:305:ARG:HB3	2.20	0.40
1:B:300:HIS:HB3	1:B:345:MET:CE	2.52	0.40
1:B:285:ARG:HA	1:B:286:PRO:HD3	1.98	0.40
1:B:132:PHE:CE1	1:B:194:GLU:HG2	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	343/355 (97%)	334 (97%)	8 (2%)	1 (0%)	46	35
1	B	340/355 (96%)	323 (95%)	17 (5%)	0	100	100
All	All	683/710 (96%)	657 (96%)	25 (4%)	1 (0%)	56	46

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	27	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	289/298 (97%)	281 (97%)	8 (3%)	51	41
1	B	287/298 (96%)	275 (96%)	12 (4%)	36	24
All	All	576/596 (97%)	556 (96%)	20 (4%)	43	31

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	13	MET
1	A	86	LYS
1	A	113	ARG
1	A	163	ARG
1	A	194	GLU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	249	SER
1	A	298	GLU
1	A	311	SER
1	B	14	THR
1	B	60	LEU
1	B	91	LEU
1	B	96	TYR
1	B	123	ASP
1	B	125	ARG
1	B	136	GLU
1	B	163	ARG
1	B	207	GLU
1	B	211	GLU
1	B	262	GLU
1	B	320	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	204	HIS
1	A	258	ASN
1	B	165	GLN
1	B	223	ASN
1	B	335	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	345/355 (97%)	0.28	14 (4%) 41 45	19, 30, 50, 68	0
1	B	342/355 (96%)	0.87	49 (14%) 4 4	30, 45, 74, 91	0
All	All	687/710 (96%)	0.57	63 (9%) 11 12	19, 38, 68, 91	0

All (63) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	96	TYR	7.7
1	A	11	GLY	7.0
1	B	174	ALA	6.7
1	B	95	ASN	6.1
1	B	76	GLY	6.0
1	A	12	SER	5.5
1	B	297	ARG	5.4
1	B	113	ARG	4.9
1	B	61	ILE	4.7
1	B	99	PHE	4.5
1	B	265	GLY	4.5
1	B	176	ILE	4.1
1	B	295	SER	3.9
1	B	110	PHE	3.9
1	B	340	GLY	3.7
1	B	94	GLN	3.7
1	B	73	TRP	3.7
1	B	342	TRP	3.6
1	B	39	ARG	3.4
1	B	173	PHE	3.2
1	B	298	GLU	3.1
1	B	175	ALA	3.1
1	B	69	LYS	3.1
1	A	61	ILE	3.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	272	PRO	2.9
1	A	297	ARG	2.9
1	B	85	GLN	2.8
1	B	299	ALA	2.8
1	B	321	ASP	2.8
1	A	144	HIS	2.8
1	B	273	ALA	2.7
1	B	332	ALA	2.7
1	A	273	ALA	2.6
1	B	75	GLY	2.6
1	B	40	GLU	2.5
1	A	176	ILE	2.5
1	B	172	PRO	2.5
1	B	317	ILE	2.4
1	B	66	ARG	2.4
1	B	93	PHE	2.4
1	B	322	GLY	2.4
1	B	294	ALA	2.4
1	B	335	HIS	2.3
1	B	246	ILE	2.3
1	A	108	VAL	2.3
1	A	134	ARG	2.3
1	B	134	ARG	2.3
1	A	57	ILE	2.3
1	B	101	HIS	2.3
1	B	154	VAL	2.3
1	B	86	LYS	2.3
1	B	301	ALA	2.2
1	B	82	LEU	2.2
1	B	83	PRO	2.2
1	A	129	LEU	2.2
1	B	132	PHE	2.1
1	A	311	SER	2.1
1	A	17	PHE	2.0
1	B	57	ILE	2.0
1	B	319	THR	2.0
1	B	296	GLU	2.0
1	B	292	GLN	2.0
1	B	289	VAL	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	CL	B	402	1/1	0.98	0.06	-1.31	37,37,37,37	0
2	CL	A	401	1/1	0.99	0.07	-1.57	25,25,25,25	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.