



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 22, 2016 – 07:09 AM EDT

PDB ID : 4Z49  
Title : Homo Sapiens Fatty Acid Synthetase, Thioesterase Domain at 1.7 Angstroms Resolution  
Authors : Spraggon, G.  
Deposited on : 2015-04-01  
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : unknown  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20027107  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0122  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027107


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3190 (1.70-1.70)
Clashscore	102246	3585 (1.70-1.70)
Ramachandran outliers	100387	3527 (1.70-1.70)
Sidechain outliers	100360	3527 (1.70-1.70)
RSRZ outliers	91569	3200 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	282	<div> <div>5%</div> <div> <div></div> <div>86%</div> <div>7%</div> <div>7%</div> </div> </div>
1	B	282	<div> <div>2%</div> <div> <div></div> <div>86%</div> <div>12%</div> <div>•</div> </div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4975 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fatty acid synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	262	Total	C	N	O	S	0	9	0
			2059	1303	352	390	14			
1	B	282	Total	C	N	O	S	0	7	0
			2216	1404	381	419	12			

- Molecule 2 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	K	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	320	Total	O	0	0
			320	320		
3	B	379	Total	O	0	0
			379	379		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Fatty acid synthase



#### • Molecule 1: Fatty acid synthase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	38.73Å 117.59Å 63.28Å 90.00° 105.58° 90.00°	Depositor
Resolution (Å)	24.89 – 1.70 31.50 – 1.70	Depositor EDS
% Data completeness (in resolution range)	96.6 (24.89-1.70) 96.6 (31.50-1.70)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	0.08	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.29 (at 1.70Å)	Xtriage
Refinement program	BUSTER 2.11.5	Depositor
R, $R_{free}$	0.171 , 0.212 0.173 , 0.216	Depositor DCC
$R_{free}$ test set	2872 reflections (5.23%)	DCC
Wilson B-factor (Å <sup>2</sup> )	14.7	Xtriage
Anisotropy	0.135	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 35.0	EDS
Estimated twinning fraction	0.032 for h,-k,-h-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 57767 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4975	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.56% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:  
K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.51	0/2130	0.62	0/2886
1	B	0.51	0/2283	0.65	1/3097 (0.0%)
All	All	0.51	0/4413	0.63	1/5983 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	B	2456	GLU	C-N-CA	5.33	135.03	121.70

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2059	0	2039	14	0
1	B	2216	0	2195	29	0
2	B	1	0	0	0	0
3	A	320	0	0	1	0
3	B	379	0	0	7	0
All	All	4975	0	4234	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 4.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:2265:SER:H	1:B:2498:HIS:HE1	1.20	0.87
1:B:2322:GLN:HE22	1:B:2440:HIS:H	1.25	0.84
1:A:2233:ARG:H	1:B:2295:GLN:HE21	1.25	0.84
1:A:2295:GLN:HE21	1:B:2233[B]:ARG:H	1.26	0.82
1:A:2295:GLN:HE21	1:B:2233[A]:ARG:H	1.26	0.81
1:A:2322:GLN:HE22	1:A:2440:HIS:H	1.26	0.80
1:B:2330:THR:HG23	3:B:2858:HOH:O	1.91	0.69
1:B:2475:HIS:HD2	3:B:3002:HOH:O	1.75	0.68
1:A:2258:HIS:HD2	3:A:2606:HOH:O	1.76	0.66
1:B:2283[B]:HIS:HD2	3:B:2817:HOH:O	1.81	0.63
1:A:2246:LEU:HD23	1:A:2305:ALA:HB3	1.81	0.63
1:B:2265:SER:H	1:B:2498:HIS:CE1	2.11	0.62
1:B:2322:GLN:NE2	1:B:2440:HIS:H	1.96	0.59
1:A:2322:GLN:NE2	1:A:2440:HIS:H	2.00	0.58
1:A:2283[A]:HIS:HD2	1:A:2316:GLU:OE2	1.89	0.55
1:B:2455:GLY:HA2	1:B:2458:LEU:HD22	1.89	0.55
1:A:2295:GLN:NE2	1:B:2233[B]:ARG:H	2.02	0.52
1:A:2295:GLN:NE2	1:B:2233[A]:ARG:H	2.03	0.52
1:B:2226:PRO:HB2	1:B:2275:ARG:HH12	1.76	0.51
1:B:2381:ASN:HD22	1:B:2382:ARG:H	1.59	0.51
1:B:2362:GLU:HG3	3:B:3019:HOH:O	2.11	0.51
1:A:2315:PHE:CG	1:A:2341:PRO:HG3	2.48	0.48
1:A:2309:TYR:HA	1:A:2344:VAL:HG13	1.95	0.48
1:A:2233:ARG:H	1:B:2295:GLN:NE2	2.04	0.47
1:B:2290:ILE:HD11	1:B:2317:MET:O	2.16	0.46
1:B:2458:LEU:HD21	1:B:2462:TYR:CD2	2.50	0.46
1:B:2283[A]:HIS:HD2	1:B:2316:GLU:OE2	1.99	0.45
1:B:2402:ASP:O	1:B:2406:LYS:HD3	2.18	0.44
1:B:2458:LEU:HG	1:B:2462:TYR:HA	1.98	0.44
1:B:2344:VAL:CG2	3:B:2809:HOH:O	2.65	0.43
1:B:2330:THR:HG21	3:B:2854:HOH:O	2.18	0.43
1:B:2459:GLY:H	1:B:2463:ASN:ND2	2.17	0.43
1:B:2381:ASN:HD22	1:B:2382:ARG:N	2.18	0.42
1:A:2312[A]:CYS:SG	1:A:2433:TYR:CD2	3.13	0.41
1:B:2467:VAL:HB	3:B:2705:HOH:O	2.19	0.41
1:B:2366:GLU:OE1	1:B:2380:HIS:NE2	2.54	0.41
1:B:2303:ARG:HG2	1:B:2501:LEU:HD22	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	265/282 (94%)	261 (98%)	4 (2%)	0	100	100
1	B	286/282 (101%)	277 (97%)	7 (2%)	2 (1%)	26	9
All	All	551/564 (98%)	538 (98%)	11 (2%)	2 (0%)	39	20

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	2456	GLU
1	B	2457	ASP

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	229/235 (97%)	224 (98%)	5 (2%)	60	39
1	B	241/235 (103%)	228 (95%)	13 (5%)	27	9
All	All	470/470 (100%)	452 (96%)	18 (4%)	38	17

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2221	SER
1	A	2225	ASN
1	A	2279	LEU
1	A	2377	ASP

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type
1	A	2384	LEU
1	B	2225	ASN
1	B	2236	SER
1	B	2266	ILE
1	B	2299	GLU
1	B	2303	ARG
1	B	2325	GLN
1	B	2381	ASN
1	B	2406	LYS
1	B	2456	GLU
1	B	2457	ASP
1	B	2458	LEU
1	B	2482	ARG
1	B	2484	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	2225	ASN
1	A	2258	HIS
1	A	2295	GLN
1	A	2320	GLN
1	A	2322	GLN
1	A	2331	HIS
1	A	2432	GLN
1	A	2466	GLN
1	A	2475	HIS
1	B	2225	ASN
1	B	2295	GLN
1	B	2320	GLN
1	B	2322	GLN
1	B	2381	ASN
1	B	2463	ASN
1	B	2466	GLN
1	B	2475	HIS
1	B	2498	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	262/282 (92%)	0.05	14 (5%) 30 32	8, 17, 43, 74	0
1	B	282/282 (100%)	-0.20	5 (1%) 71 76	8, 15, 35, 62	0
All	All	544/564 (96%)	-0.08	19 (3%) 48 52	8, 16, 40, 74	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2344	VAL	6.8
1	A	2378	MET	5.0
1	B	2238	GLN	4.3
1	A	2382	ARG	3.9
1	A	2343	TYR	3.9
1	A	2358	GLY	3.9
1	A	2381	ASN	3.8
1	B	2456	GLU	3.8
1	A	2379	GLU	3.6
1	A	2377	ASP	3.6
1	A	2376	THR	3.3
1	A	2406	LYS	2.6
1	B	2458	LEU	2.6
1	A	2384	LEU	2.5
1	B	2457	ASP	2.4
1	B	2454	TYR	2.3
1	A	2375	PHE	2.2
1	A	2383	VAL	2.2
1	A	2407	SER	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	K	B	2601	1/1	1.00	0.04	-4.77	19,19,19,19	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.