



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 13, 2016 – 06:43 PM EDT

PDB ID : 4Z4M  
Title : Crystal structure of GFP-TAX1BP1 UBZ2 domain fusion protein  
Authors : Rohaim, A.; Kawasaki, M.; Wakatsuki, S.  
Deposited on : 2015-04-02  
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20027790  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027790


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1045 (2.16-2.16)
Clashscore	102246	1152 (2.16-2.16)
Ramachandran outliers	100387	1131 (2.16-2.16)
Sidechain outliers	100360	1131 (2.16-2.16)
RSRZ outliers	91569	1050 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	260	 76% 19% . .
1	B	260	 77% 18% . . .

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4269 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Green fluorescent protein,Tax1-binding protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	254	Total	C	N	O	S	0	0	0
			2046	1303	345	388	10			
1	B	254	Total	C	N	O	S	0	0	0
			2046	1303	345	388	10			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP P42212
A	-1	SER	-	expression tag	UNP P42212
A	0	HIS	-	expression tag	UNP P42212
A	66	CR2	SER	CHROMOPHORE	UNP P42212
A	66	CR2	TYR	CHROMOPHORE	UNP P42212
A	66	CR2	GLY	CHROMOPHORE	UNP P42212
A	72	ALA	SER	engineered mutation	UNP P42212
A	233	GLY	-	linker	UNP P42212
A	234	SER	-	linker	UNP P42212
B	-2	GLY	-	expression tag	UNP P42212
B	-1	SER	-	expression tag	UNP P42212
B	0	HIS	-	expression tag	UNP P42212
B	66	CR2	SER	CHROMOPHORE	UNP P42212
B	66	CR2	TYR	CHROMOPHORE	UNP P42212
B	66	CR2	GLY	CHROMOPHORE	UNP P42212
B	72	ALA	SER	engineered mutation	UNP P42212
B	233	GLY	-	linker	UNP P42212
B	234	SER	-	linker	UNP P42212

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Zn	0	0
			1	1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	89	Total	O	0	0
			89	89		
3	B	86	Total	O	0	0
			86	86		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Green fluorescent protein,Tax1-binding protein 1



- Molecule 1: Green fluorescent protein,Tax1-binding protein 1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	67.73Å 88.77Å 91.95Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	63.87 – 2.15 45.97 – 2.15	Depositor EDS
% Data completeness (in resolution range)	97.3 (63.87-2.15) 97.3 (45.97-2.15)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	7.63 (at 2.16Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.234 , 0.298 0.233 , 0.299	Depositor DCC
$R_{free}$ test set	1504 reflections (5.30%)	DCC
Wilson B-factor (Å <sup>2</sup> )	24.0	Xtriage
Anisotropy	0.879	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 38.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.000 for -h,l,k	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4269	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 57.85 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 2.2293e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CR2, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.08	4/2076 (0.2%)	0.99	2/2804 (0.1%)
1	B	1.09	5/2076 (0.2%)	0.99	3/2804 (0.1%)
All	All	1.09	9/4152 (0.2%)	0.99	5/5608 (0.1%)

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	154	ALA	CA-CB	7.47	1.68	1.52
1	B	154	ALA	CA-CB	6.78	1.66	1.52
1	B	32	GLU	CG-CD	-6.39	1.42	1.51
1	B	99	PHE	CE1-CZ	6.38	1.49	1.37
1	A	124	GLU	CG-CD	5.72	1.60	1.51
1	B	72	ALA	CA-CB	5.67	1.64	1.52
1	A	95	GLU	CG-CD	-5.14	1.44	1.51
1	A	99	PHE	CE1-CZ	5.13	1.47	1.37
1	B	112	VAL	CB-CG1	-5.08	1.42	1.52

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	215	ARG	NE-CZ-NH2	-7.57	116.52	120.30
1	B	215	ARG	NE-CZ-NH2	-6.02	117.29	120.30
1	A	141	LEU	CB-CG-CD1	-5.98	100.83	111.00
1	B	7	LEU	CA-CB-CG	5.33	127.55	115.30
1	B	141	LEU	CA-CB-CG	5.25	127.38	115.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2046	0	1966	38	0
1	B	2046	0	1966	33	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	89	0	0	5	0
3	B	86	0	0	3	0
All	All	4269	0	3932	69	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:235:LYS:NZ	1:A:248:TYR:HB3	1.84	0.91
1:A:26:LYS:HE3	3:A:485:HOH:O	1.76	0.85
1:A:235:LYS:HZ2	1:A:248:TYR:HB3	1.45	0.80
1:A:42:LEU:HD21	1:A:71:PHE:HB2	1.67	0.76
1:B:78:MET:CE	1:B:229:ILE:HD12	2.16	0.75
1:A:79:LYS:NZ	3:A:401:HOH:O	2.18	0.75
1:A:15:LEU:HD23	1:A:118:THR:HG21	1.72	0.72
1:B:46:PHE:CZ	1:B:64:PHE:HB3	2.26	0.70
1:A:153:MET:CE	1:A:162:LYS:HE3	2.21	0.70
1:B:78:MET:HE3	1:B:229:ILE:HD12	1.74	0.70
1:B:221:LEU:HG	1:B:222:GLU:N	2.06	0.68
1:A:235:LYS:HZ1	1:A:248:TYR:HB3	1.57	0.68
1:A:235:LYS:N	3:A:404:HOH:O	2.29	0.65
1:B:256:HIS:O	1:B:259:THR:HB	1.96	0.65
1:B:221:LEU:HD21	1:B:223:PHE:CE2	2.33	0.64
1:B:81:HIS:CD2	1:B:229:ILE:HD13	2.33	0.64
1:B:238:PRO:HG2	1:B:239:MET:HE2	1.80	0.62
1:B:78:MET:HE2	1:B:229:ILE:HD12	1.82	0.62
1:B:4:GLY:HA2	1:B:7:LEU:HD22	1.83	0.61
1:A:46:PHE:CZ	1:A:64:PHE:HB3	2.38	0.59
1:B:4:GLY:HA3	1:B:85:LYS:O	2.02	0.58

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:29:VAL:HG11	1:A:64:PHE:CZ	2.39	0.57
1:A:203:THR:HG21	3:A:454:HOH:O	2.04	0.57
1:A:4:GLY:HA3	1:A:85:LYS:O	2.05	0.57
1:A:80:GLN:O	1:A:194:LEU:HG	2.05	0.57
1:A:42:LEU:HD21	1:A:71:PHE:CB	2.34	0.56
1:B:171:ILE:HD12	1:B:175:SER:HB3	1.87	0.55
1:A:199:HIS:HB2	1:A:227:ALA:O	2.07	0.53
1:A:76:ASP:OD1	1:A:79:LYS:HE3	2.08	0.53
1:B:163:VAL:HB	1:B:183:GLN:HB3	1.90	0.52
1:A:153:MET:HE2	1:A:162:LYS:HE3	1.89	0.52
1:B:58:PRO:HA	1:B:61:VAL:HG23	1.91	0.52
1:A:238:PRO:HB2	1:A:239:MET:HE2	1.92	0.51
1:B:210:ASP:OD1	1:B:210:ASP:C	2.49	0.51
1:A:221:LEU:HG	1:A:222:GLU:N	2.25	0.50
1:B:53:LEU:HD22	1:B:57:TRP:CE2	2.46	0.50
1:A:256:HIS:O	1:A:259:THR:HB	2.11	0.50
1:A:211:PRO:HD2	1:B:39:TYR:CE2	2.46	0.49
1:A:221:LEU:HD22	1:B:221:LEU:HD22	1.95	0.49
1:B:199:HIS:HB2	1:B:227:ALA:O	2.13	0.49
1:A:238:PRO:HG2	1:A:239:MET:HE3	1.96	0.47
1:A:4:GLY:O	1:A:7:LEU:HB2	2.15	0.47
1:B:235:LYS:N	3:B:407:HOH:O	2.48	0.46
1:A:153:MET:HE3	1:A:162:LYS:HE3	1.96	0.46
1:B:172:GLU:HG3	1:B:172:GLU:H	1.50	0.46
1:B:92:TYR:CE1	1:B:112:VAL:CG1	2.99	0.45
1:B:203:THR:HB	1:B:224:VAL:HG22	1.99	0.45
1:B:32:GLU:HG2	1:B:33:GLY:N	2.25	0.45
1:A:238:PRO:HG2	1:A:239:MET:CE	2.46	0.45
1:A:53:LEU:HD22	1:A:57:TRP:CD2	2.51	0.45
1:A:53:LEU:HD22	1:A:57:TRP:CE2	2.51	0.44
1:B:235:LYS:HG2	1:B:244:PHE:HB2	1.99	0.44
1:B:86:SER:HB3	1:B:194:LEU:HD22	2.00	0.43
1:A:255:ARG:NH2	3:A:412:HOH:O	2.51	0.43
1:B:238:PRO:CG	1:B:239:MET:HE2	2.47	0.43
1:A:171:ILE:HG22	1:A:172:GLU:N	2.33	0.43
1:A:173:ASP:OD2	1:A:175:SER:N	2.45	0.43
1:A:4:GLY:HA2	1:A:7:LEU:HD22	2.01	0.43
1:A:133:ASP:N	1:A:133:ASP:OD1	2.51	0.43
1:B:101:LYS:O	3:B:401:HOH:O	2.21	0.42
1:B:92:TYR:CE1	1:B:112:VAL:HG13	2.54	0.42
1:B:42:LEU:HD21	1:B:71:PHE:HB2	2.01	0.42

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:112:VAL:HG23	1:B:119:LEU:HD11	2.01	0.42
1:A:92:TYR:CE1	1:A:112:VAL:HG13	2.55	0.42
1:A:172:GLU:C	1:A:174:GLY:H	2.23	0.41
1:B:229:ILE:HG23	3:B:486:HOH:O	2.21	0.41
1:B:238:PRO:HG2	1:B:239:MET:CE	2.48	0.41
1:A:103:ASP:OD1	1:A:130:PHE:HA	2.20	0.41
1:A:240:CYS:HB3	1:A:260:HIS:CE1	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	247/260 (95%)	238 (96%)	9 (4%)	0	100	100
1	B	247/260 (95%)	240 (97%)	7 (3%)	0	100	100
All	All	494/520 (95%)	478 (97%)	16 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	224/228 (98%)	210 (94%)	14 (6%)	22	16

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	224/228 (98%)	210 (94%)	14 (6%)	22	16
All	All	448/456 (98%)	420 (94%)	28 (6%)	22	16

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	LEU
1	A	32	GLU
1	A	44	LEU
1	A	112	VAL
1	A	133	ASP
1	A	145	TYR
1	A	157	GLN
1	A	175	SER
1	A	178	LEU
1	A	194	LEU
1	A	203	THR
1	A	225	THR
1	A	229	ILE
1	A	259	THR
1	B	7	LEU
1	B	15	LEU
1	B	32	GLU
1	B	44	LEU
1	B	80	GLN
1	B	112	VAL
1	B	145	TYR
1	B	172	GLU
1	B	178	LEU
1	B	194	LEU
1	B	203	THR
1	B	225	THR
1	B	255	ARG
1	B	259	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	149	ASN
1	A	170	ASN
1	A	184	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	149	ASN
1	B	170	ASN
1	B	212	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	CR2	A	66	1	20,20,21	3.88	4 (20%)	24,27,29	5.09	8 (33%)
1	CR2	B	66	1	20,20,21	4.12	8 (40%)	24,27,29	6.33	11 (45%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CR2	A	66	1	-	0/6/25/26	0/2/2/2
1	CR2	B	66	1	-	0/6/25/26	0/2/2/2

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	66	CR2	CA2-C2	-4.13	1.44	1.48
1	B	66	CR2	C2-N3	-3.86	1.30	1.39
1	A	66	CR2	C2-N3	-3.39	1.31	1.39
1	B	66	CR2	CA2-N2	-3.17	1.31	1.38

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	66	CR2	C1-N3	-3.03	1.32	1.37
1	A	66	CR2	CA2-C2	-2.71	1.45	1.48
1	A	66	CR2	CE1-CD1	2.04	1.42	1.38
1	B	66	CR2	CE1-CD1	2.12	1.43	1.38
1	B	66	CR2	CA1-C1	2.15	1.51	1.49
1	B	66	CR2	CG2-CB2	3.27	1.53	1.46
1	B	66	CR2	CB2-CA2	16.09	1.49	1.35
1	A	66	CR2	CB2-CA2	16.13	1.49	1.35

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	66	CR2	O2-C2-CA2	-16.97	121.22	130.97
1	A	66	CR2	O2-C2-CA2	-13.09	123.45	130.97
1	B	66	CR2	C2-CA2-N2	-10.27	101.21	109.03
1	B	66	CR2	CG2-CB2-CA2	-8.20	119.89	130.27
1	A	66	CR2	C2-CA2-N2	-7.93	102.99	109.03
1	A	66	CR2	CG2-CB2-CA2	-6.58	121.94	130.27
1	A	66	CR2	C3-CA3-N3	-4.28	104.12	112.97
1	B	66	CR2	C3-CA3-N3	-3.82	105.08	112.97
1	B	66	CR2	CD2-CG2-CB2	-2.20	113.74	121.24
1	B	66	CR2	CA3-N3-C1	2.03	130.83	127.91
1	B	66	CR2	CB2-CA2-N2	2.09	132.43	128.71
1	B	66	CR2	CA1-C1-N3	2.12	125.81	122.89
1	A	66	CR2	CD1-CG2-CD2	2.28	121.03	117.62
1	A	66	CR2	CB2-CA2-N2	2.65	133.41	128.71
1	B	66	CR2	CB2-CA2-C2	2.93	126.36	122.24
1	A	66	CR2	CA2-N2-C1	3.41	108.27	105.56
1	B	66	CR2	CA2-N2-C1	6.02	110.35	105.56
1	A	66	CR2	CA2-C2-N3	16.92	111.74	103.37
1	B	66	CR2	CA2-C2-N3	20.16	113.35	103.37

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	253/260 (97%)	-0.09	2 (0%) 87 90	14, 23, 38, 56	0
1	B	253/260 (97%)	-0.09	1 (0%) 93 94	14, 23, 43, 60	0
All	All	506/520 (97%)	-0.09	3 (0%) 90 92	14, 23, 41, 60	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	229	ILE	3.2
1	A	190	ASP	2.2
1	A	1	MET	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
1	CR2	B	66	19/20	0.96	0.10	-	12,15,20,21	0
1	CR2	A	66	19/20	0.97	0.09	-	13,16,20,23	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	ZN	B	301	1/1	0.99	0.05	-2.33	23,23,23,23	0
2	ZN	A	301	1/1	0.96	0.05	-6.04	26,26,26,26	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.