



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2016 – 08:49 PM GMT

PDB ID : 4Z8N  
Title : Crystal structure of the erythrocyte-binding domain from Plasmodium vivax reticulocyte-binding protein 2a (PvRBP 2a)  
Authors : Gruszczyk, J.; Tham, W.H.  
Deposited on : 2015-04-09  
Resolution : 2.12 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : unknown  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026982  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20026982


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*



The reported resolution of this entry is 2.12 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	4587 (2.14-2.10)
Clashscore	102246	5132 (2.14-2.10)
Ramachandran outliers	100387	5080 (2.14-2.10)
Sidechain outliers	100360	5081 (2.14-2.10)
RSRZ outliers	91569	4597 (2.14-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	298	 4% 91% 9%
1	B	298	 7% 87% 8% 5%

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4945 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Reticulocyte-binding protein 2a.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	298	Total	C	N	O	S	0	0	0
			2424	1549	393	471	11			
1	B	284	Total	C	N	O	S	0	0	0
			2315	1482	374	448	11			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	158	GLY	-	cloning artifact	UNP A5JZ09
A	159	SER	-	cloning artifact	UNP A5JZ09
B	158	GLY	-	cloning artifact	UNP A5JZ09
B	159	SER	-	cloning artifact	UNP A5JZ09

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Cl	0	0
			1	1		

- Molecule 3 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	2	Total	Mg	0	0
			2	2		

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Ca	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	102	Total	O	0	0
			102	102		
6	B	93	Total	O	0	0
			93	93		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Reticulocyte-binding protein 2a



#### • Molecule 1: Reticulocyte-binding protein 2a



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	58.79 Å 93.45 Å 126.72 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.84 – 2.12 46.73 – 2.12	Depositor EDS
% Data completeness (in resolution range)	100.0 (43.84-2.12) 100.0 (46.73-2.12)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.67 (at 2.12 Å)	Xtriage
Refinement program	PHENIX	Depositor
R, $R_{free}$	0.203 , 0.229 0.212 , 0.235	Depositor DCC
$R_{free}$ test set	1031 reflections (2.64%)	DCC
Wilson B-factor (Å <sup>2</sup> )	39.5	Xtriage
Anisotropy	0.135	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 49.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 40144 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4945	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.40% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, PEG, MG, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.25	0/2470	0.39	0/3342
1	B	0.28	0/2358	0.42	0/3187
All	All	0.26	0/4828	0.40	0/6529

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2424	0	2407	19	0
1	B	2315	0	2302	21	0
2	A	1	0	0	0	0
3	A	7	0	10	0	0
4	B	2	0	0	0	0
5	B	1	0	0	0	0
6	A	102	0	0	7	0
6	B	93	0	0	3	0
All	All	4945	0	4719	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:200:GLU:HG2	1:B:296:SER:O	1.23	1.33
1:B:200:GLU:CG	1:B:296:SER:O	1.78	1.29
1:B:200:GLU:HG3	1:B:296:SER:O	1.74	0.85
1:A:317:GLU:OE2	6:A:601:HOH:O	1.97	0.82
1:A:288:GLU:OE1	6:A:603:HOH:O	2.08	0.71
1:B:309:GLU:O	6:B:601:HOH:O	2.07	0.71
1:B:195:ILE:HG23	1:B:202:LYS:HB2	1.73	0.69
1:A:376:GLU:OE2	6:A:604:HOH:O	2.11	0.68
1:A:230:LYS:NZ	1:A:230:LYS:O	2.27	0.64
1:A:163:ARG:NH2	1:A:436:GLU:OE1	2.31	0.62
1:A:196:THR:HA	1:A:202:LYS:HD3	1.82	0.61
1:A:179:TYR:O	1:A:425:LYS:NZ	2.30	0.60
1:B:175:ILE:HD12	1:B:273:GLN:HG2	1.86	0.57
1:B:200:GLU:HG3	1:B:296:SER:HB3	1.90	0.53
1:B:316:VAL:O	1:B:320:LYS:HG3	2.09	0.53
1:A:455:LYS:HE2	6:A:638:HOH:O	2.09	0.53
1:A:199:TYR:O	1:A:203:LYS:HG2	2.09	0.52
1:A:186:ASN:ND2	6:A:608:HOH:O	2.43	0.50
1:A:291:TYR:OH	6:A:602:HOH:O	2.04	0.49
1:B:205:TYR:HA	1:B:293:ILE:HD13	1.95	0.47
1:B:202:LYS:NZ	1:B:206:ASP:OD1	2.47	0.47
1:A:364:LEU:HD11	1:A:437:LYS:HB3	1.96	0.47
1:B:202:LYS:HA	1:B:205:TYR:HB3	1.97	0.46
1:B:200:GLU:OE1	1:B:298:LYS:NZ	2.48	0.46
1:B:364:LEU:HD12	1:B:444:LEU:HD12	1.97	0.46
1:A:347:LYS:HA	1:A:347:LYS:HD2	1.89	0.44
1:B:257:LYS:NZ	1:B:260:GLU:OE1	2.46	0.44
1:A:236:LYS:HD2	1:A:236:LYS:HA	1.78	0.43
1:B:327:TYR:OH	1:B:428:ASP:OD1	2.23	0.43
1:A:258:HIS:CD2	1:A:262:LYS:HE3	2.54	0.42
1:A:241:LEU:HD21	1:A:257:LYS:HB3	2.01	0.42
1:A:164:TYR:CD1	1:A:371:ILE:HG12	2.55	0.42
1:A:333:THR:HG22	1:B:330:TYR:CE1	2.55	0.41
1:B:248:ASN:ND2	6:B:603:HOH:O	2.41	0.41
1:B:408:LYS:HE3	1:B:412:ASP:OD2	2.21	0.41
1:B:436:GLU:HG2	6:B:613:HOH:O	2.21	0.41
1:B:200:GLU:HG3	1:B:296:SER:C	2.41	0.41
1:B:196:THR:HG22	1:B:202:LYS:HE2	2.02	0.41
1:A:277:GLU:HB3	6:A:680:HOH:O	2.22	0.40



There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	296/298 (99%)	290 (98%)	6 (2%)	0	100	100
1	B	280/298 (94%)	272 (97%)	8 (3%)	0	100	100
All	All	576/596 (97%)	562 (98%)	14 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	273/273 (100%)	273 (100%)	0	100	100
1	B	260/273 (95%)	260 (100%)	0	100	100
All	All	533/546 (98%)	533 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	365	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 5 ligands modelled in this entry, 4 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	PEG	A	502	-	6,6,6	0.62	0	5,5,5	0.36	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PEG	A	502	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	298/298 (100%)	0.17	12 (4%)	42 51	29, 49, 88, 110	0
1	B	284/298 (95%)	0.25	21 (7%)	17 23	32, 53, 99, 119	0
All	All	582/596 (97%)	0.21	33 (5%)	27 35	29, 51, 96, 119	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	311	MET	5.6
1	B	291	TYR	4.5
1	B	195	ILE	3.9
1	B	310	ILE	3.6
1	B	298	LYS	3.6
1	A	300	THR	3.5
1	B	314	ILE	3.4
1	B	299	CYS	3.4
1	B	406	LEU	3.2
1	B	203	LYS	3.2
1	B	200	GLU	3.0
1	A	398	PRO	3.0
1	B	201	ALA	2.9
1	A	453	SER	2.9
1	A	302	ALA	2.9
1	B	193	TYR	2.8
1	A	397	ILE	2.8
1	B	202	LYS	2.8
1	B	411	PHE	2.7
1	A	454	GLU	2.6
1	B	313	GLU	2.5
1	A	158	GLY	2.5
1	A	299	CYS	2.5
1	B	196	THR	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	194	TYR	2.4
1	B	315	TYR	2.4
1	A	396	ASN	2.3
1	A	170	SER	2.3
1	A	455	LYS	2.3
1	B	413	SER	2.2
1	B	302	ALA	2.1
1	B	204	ASN	2.0
1	A	327	TYR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	MG	B	502	1/1	0.93	0.13	1.38	53,53,53,53	0
3	PEG	A	502	7/7	0.70	0.20	0.36	74,77,81,81	0
5	CA	B	503	1/1	0.79	0.09	-1.18	95,95,95,95	0
4	MG	B	501	1/1	0.96	0.10	-	62,62,62,62	0
2	CL	A	501	1/1	0.93	0.13	-	84,84,84,84	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.