



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 07:03 AM GMT

PDB ID : 2ZAT  
Title : Crystal structure of a mammalian reductase  
Authors : Tanaka, N.; Aoki, K.; Nakamura, K.T.  
Deposited on : 2007-10-10  
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	2274 (1.50-1.50)
Ramachandran outliers	100387	2218 (1.50-1.50)
Sidechain outliers	100360	2216 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	260	
1	B	260	
1	C	260	
1	D	260	

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 8480 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dehydrogenase/reductase SDR family member 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	251	Total	C	N	O	S	0	0	0
			1879	1179	337	354	9			
1	B	251	Total	C	N	O	S	0	0	0
			1879	1179	337	354	9			
1	C	251	Total	C	N	O	S	0	0	0
			1879	1179	337	354	9			
1	D	251	Total	C	N	O	S	0	0	0
			1879	1179	337	354	9			

- Molecule 2 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula:  $C_{21}H_{28}N_7O_{17}P_3$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	B	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	C	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	D	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	178	Total	O	0	0
			178	178		
3	B	197	Total	O	0	0
			197	197		
3	C	196	Total	O	0	0
			196	196		
3	D	201	Total	O	0	0
			201	201		

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: Dehydrogenase/reductase SDR family member 4




- Molecule 1: Dehydrogenase/reductase SDR family member 4



- Molecule 1: Dehydrogenase/reductase SDR family member 4



- Molecule 1: Dehydrogenase/reductase SDR family member 4



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 42	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	109.61Å 109.61Å 94.31Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 1.50	Depositor
% Data completeness (in resolution range)	99.0 (40.00-1.50)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.10	Depositor
Refinement program	REFMAC 5.1.24	Depositor
R, $R_{free}$	0.161 , 0.187	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	8480	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	17.0	wwPDB-VP

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.64	0/1905	0.78	4/2582 (0.2%)
1	B	0.60	0/1905	0.76	2/2582 (0.1%)
1	C	0.60	0/1905	0.76	2/2582 (0.1%)
1	D	0.61	0/1905	0.77	4/2582 (0.2%)
All	All	0.61	0/7620	0.77	12/10328 (0.1%)

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	110	ASP	CB-CG-OD2	8.67	126.10	118.30
1	C	259	ARG	NE-CZ-NH2	-7.01	116.79	120.30
1	A	259	ARG	NE-CZ-NH2	-6.35	117.13	120.30
1	B	208	ASP	CB-CG-OD2	6.28	123.95	118.30
1	D	259	ARG	NE-CZ-NH2	-6.24	117.18	120.30
1	D	208	ASP	CB-CG-OD2	5.96	123.66	118.30
1	A	208	ASP	CB-CG-OD2	5.88	123.59	118.30
1	B	188	ARG	NE-CZ-NH1	5.53	123.07	120.30
1	D	32	ARG	NE-CZ-NH1	5.43	123.01	120.30
1	C	208	ASP	CB-CG-OD2	5.38	123.14	118.30
1	D	188	ARG	NE-CZ-NH1	5.26	122.93	120.30
1	A	259	ARG	NE-CZ-NH1	5.08	122.84	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1879	0	1937	7	0
1	B	1879	0	1937	11	0
1	C	1879	0	1937	11	0
1	D	1879	0	1937	13	0
2	A	48	0	25	2	0
2	B	48	0	25	5	0
2	C	48	0	25	3	0
2	D	48	0	25	2	0
3	A	178	0	0	3	0
3	B	197	0	0	8	0
3	C	196	0	0	4	0
3	D	201	0	0	2	0
All	All	8480	0	7848	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:3261:NAP:O3D	3:C:3448:HOH:O	1.78	0.99
2:B:2261:NAP:C5N	3:B:2392:HOH:O	2.37	0.71
1:B:104:PHE:O	3:B:2447:HOH:O	2.11	0.69
1:C:196:LEU:HD21	1:C:219:LEU:HG	1.80	0.64
1:D:196:LEU:HD13	1:D:224:LEU:HD22	1.78	0.64
1:A:139:GLU:OE1	1:A:185:ARG:NE	2.31	0.63
1:A:104:PHE:O	3:A:1437:HOH:O	2.15	0.63
1:B:161:LEU:HA	3:B:2447:HOH:O	2.01	0.60
2:B:2261:NAP:H5N	3:B:2444:HOH:O	2.00	0.60
2:B:2261:NAP:H5N	3:B:2392:HOH:O	1.97	0.59
2:C:3261:NAP:C5N	3:C:3447:HOH:O	2.51	0.58
1:A:98:ASN:OD1	2:A:1261:NAP:H4D	2.09	0.52
1:B:156:HIS:NE2	1:D:156:HIS:CE1	2.79	0.50
1:C:125:LYS:HE3	1:D:113:GLU:HG3	1.94	0.50
1:D:158:PHE:CB	1:D:161:LEU:HD12	2.42	0.49

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:125:LYS:CE	1:D:113:GLU:HG3	2.43	0.49
2:A:1261:NAP:C5N	3:A:1426:HOH:O	2.60	0.48
1:B:156:HIS:NE2	1:D:156:HIS:HE1	2.12	0.48
1:D:158:PHE:HB2	1:D:161:LEU:HD12	1.95	0.48
1:B:107:ASN:HD22	1:B:109:ILE:H	1.62	0.48
1:C:248:GLU:OE1	3:C:3454:HOH:O	2.20	0.47
1:B:196:LEU:HD13	1:B:224:LEU:HD22	1.98	0.46
1:D:98:ASN:OD1	2:D:4261:NAP:H4D	2.15	0.46
1:A:181:GLU:OE1	1:B:108:ILE:HG22	2.16	0.46
2:D:4261:NAP:C5N	3:D:4445:HOH:O	2.64	0.45
1:C:98:ASN:OD1	2:C:3261:NAP:H4D	2.16	0.45
1:A:161:LEU:HA	3:A:1437:HOH:O	2.15	0.45
1:C:152:VAL:HG22	1:C:251:VAL:HG13	1.98	0.45
1:C:137:GLU:HG2	3:C:3450:HOH:O	2.15	0.45
1:A:196:LEU:HD13	1:A:224:LEU:HD22	1.99	0.45
1:B:18:LEU:HD21	1:B:85:ALA:HB2	1.99	0.44
1:A:159:PRO:O	1:A:160:ASN:HB2	2.18	0.43
1:B:107:ASN:ND2	3:B:2407:HOH:O	2.50	0.43
1:B:107:ASN:CG	3:B:2407:HOH:O	2.57	0.42
1:C:113:GLU:HG2	1:D:129:LEU:HD11	2.01	0.42
2:B:2261:NAP:C5N	3:B:2444:HOH:O	2.63	0.42
1:C:28:LEU:HD13	1:C:54:THR:HG23	2.00	0.42
1:D:205:LEU:HD22	3:D:4362:HOH:O	2.19	0.41
1:D:18:LEU:HD21	1:D:85:ALA:HB2	2.02	0.41
1:B:199:THR:OG1	2:B:2261:NAP:N7N	2.53	0.41
1:C:181:GLU:OE1	1:D:107:ASN:HA	2.20	0.41
1:C:177:ASN:OD1	1:D:157:PRO:HB3	2.21	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	249/260 (96%)	243 (98%)	6 (2%)	0	100	100
1	B	249/260 (96%)	244 (98%)	5 (2%)	0	100	100
1	C	249/260 (96%)	241 (97%)	7 (3%)	1 (0%)	39	14
1	D	249/260 (96%)	243 (98%)	6 (2%)	0	100	100
All	All	996/1040 (96%)	971 (98%)	24 (2%)	1 (0%)	56	26

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	207	MET

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	204/211 (97%)	198 (97%)	6 (3%)	50	16
1	B	204/211 (97%)	198 (97%)	6 (3%)	50	16
1	C	204/211 (97%)	200 (98%)	4 (2%)	63	29
1	D	204/211 (97%)	199 (98%)	5 (2%)	55	20
All	All	816/844 (97%)	795 (97%)	21 (3%)	54	19

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	196	LEU
1	A	207	MET
1	A	208	ASP
1	A	213	GLU
1	A	220	ARG
1	A	259	ARG
1	B	107	ASN
1	B	196	LEU
1	B	211	ARG
1	B	212	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	213	GLU
1	B	259	ARG
1	C	79	GLU
1	C	209	LYS
1	C	213	GLU
1	C	259	ARG
1	D	114	GLU
1	D	196	LEU
1	D	207	MET
1	D	213	GLU
1	D	259	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	107	ASN
1	D	156	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NAP	A	1261	-	42,52,52	1.66	4 (9%)	54,80,80	1.77	6 (11%)
2	NAP	B	2261	-	42,52,52	1.74	6 (14%)	54,80,80	1.87	12 (22%)
2	NAP	C	3261	-	42,52,52	1.64	4 (9%)	54,80,80	1.64	12 (22%)
2	NAP	D	4261	-	42,52,52	1.59	2 (4%)	54,80,80	1.86	7 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAP	A	1261	-	-	0/27/67/67	0/5/5/5
2	NAP	B	2261	-	-	0/27/67/67	0/5/5/5
2	NAP	C	3261	-	-	0/27/67/67	0/5/5/5
2	NAP	D	4261	-	-	0/27/67/67	0/5/5/5

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	2261	NAP	PN-O2N	-2.37	1.44	1.54
2	B	2261	NAP	C3N-C7N	-2.16	1.47	1.50
2	C	3261	NAP	PN-O2N	-2.12	1.45	1.54
2	A	1261	NAP	O4B-C1B	2.11	1.43	1.41
2	C	3261	NAP	C2A-N1A	2.48	1.38	1.33
2	B	2261	NAP	C2A-N1A	2.56	1.38	1.33
2	A	1261	NAP	C2A-N1A	2.72	1.39	1.33
2	B	2261	NAP	O4B-C1B	2.81	1.44	1.41
2	B	2261	NAP	C2A-N3A	3.12	1.37	1.32
2	A	1261	NAP	C2A-N3A	3.34	1.38	1.32
2	D	4261	NAP	C2A-N3A	3.36	1.38	1.32
2	C	3261	NAP	C2A-N3A	3.61	1.38	1.32
2	C	3261	NAP	O7N-C7N	8.17	1.41	1.24
2	D	4261	NAP	O7N-C7N	8.28	1.41	1.24
2	A	1261	NAP	O7N-C7N	8.60	1.42	1.24
2	B	2261	NAP	O7N-C7N	8.63	1.42	1.24

All (37) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	4261	NAP	N3A-C2A-N1A	-9.81	121.38	128.89
2	A	1261	NAP	N3A-C2A-N1A	-9.52	121.60	128.89
2	C	3261	NAP	N3A-C2A-N1A	-6.60	123.84	128.89
2	B	2261	NAP	N3A-C2A-N1A	-6.45	123.96	128.89
2	B	2261	NAP	O7N-C7N-C3N	-4.63	114.53	119.59
2	B	2261	NAP	C3N-C2N-N1N	-4.40	115.29	120.36
2	C	3261	NAP	O7N-C7N-C3N	-3.92	115.30	119.59
2	D	4261	NAP	O7N-C7N-C3N	-3.64	115.61	119.59
2	B	2261	NAP	C4N-C3N-C7N	-3.28	112.42	121.09
2	B	2261	NAP	C4A-C5A-N7A	-3.02	106.70	109.48
2	B	2261	NAP	C4D-O4D-C1D	-2.59	106.87	109.72
2	A	1261	NAP	C4D-O4D-C1D	-2.41	107.07	109.72
2	C	3261	NAP	C4A-C5A-N7A	-2.34	107.32	109.48
2	B	2261	NAP	C1B-N9A-C4A	-2.34	123.41	126.94
2	C	3261	NAP	O3B-C3B-C4B	-2.30	104.14	111.05
2	C	3261	NAP	C3N-C2N-N1N	-2.29	117.72	120.36
2	C	3261	NAP	C1B-N9A-C4A	-2.25	123.55	126.94
2	A	1261	NAP	C4A-C5A-N7A	-2.25	107.41	109.48
2	D	4261	NAP	C3N-C2N-N1N	-2.19	117.84	120.36
2	D	4261	NAP	C4N-C3N-C7N	-2.15	115.41	121.09
2	C	3261	NAP	O4B-C1B-C2B	-2.11	102.78	106.60
2	B	2261	NAP	O3B-C3B-C4B	-2.02	105.00	111.05
2	D	4261	NAP	O2A-PA-O3	2.00	114.18	105.09
2	B	2261	NAP	O2A-PA-O3	2.07	114.47	105.09
2	C	3261	NAP	C3N-C7N-N7N	2.12	120.14	117.82
2	B	2261	NAP	O4D-C4D-C3D	2.13	109.43	105.15
2	C	3261	NAP	O2A-PA-O3	2.21	115.10	105.09
2	C	3261	NAP	C4B-O4B-C1B	2.26	112.20	109.72
2	C	3261	NAP	O4D-C1D-N1N	2.31	110.67	108.13
2	A	1261	NAP	O2A-PA-O3	2.42	116.06	105.09
2	A	1261	NAP	C2N-C3N-C4N	2.65	121.24	118.29
2	C	3261	NAP	C2N-C3N-C4N	2.91	121.53	118.29
2	B	2261	NAP	O7N-C7N-N7N	3.04	126.87	122.59
2	A	1261	NAP	O4D-C1D-N1N	3.12	111.56	108.13
2	D	4261	NAP	C2N-C3N-C4N	3.18	121.83	118.29
2	B	2261	NAP	C2N-C3N-C4N	3.32	121.98	118.29
2	D	4261	NAP	O4D-C1D-N1N	3.43	111.90	108.13

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1261	NAP	2	0
2	B	2261	NAP	5	0
2	C	3261	NAP	3	0
2	D	4261	NAP	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

### 6.3 Carbohydrates [i](#)

EDS was not executed - this section will therefore be empty.

### 6.4 Ligands [i](#)

EDS was not executed - this section will therefore be empty.

### 6.5 Other polymers [i](#)

EDS was not executed - this section will therefore be empty.