



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 11:59 PM GMT

PDB ID : 1ZBH
Title : 3'-end specific recognition of histone mRNA stem-loop by 3'-exonuclease
Authors : Cheng, Y.; Patel, D.J.
Deposited on : 2005-04-08
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1578 (3.00-3.00)
Clashscore	102246	1912 (3.00-3.00)
Ramachandran outliers	100387	1853 (3.00-3.00)
Sidechain outliers	100360	1856 (3.00-3.00)
RSRZ outliers	91569	1592 (3.00-3.00)
RNA backbone	2183	1036 (3.40-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	E	20	<div> <div>15%</div> <div>35%</div> <div>40%</div> <div>5%</div> <div>20%</div> </div>
1	F	20	<div> <div>5%</div> <div>35%</div> <div>30%</div> <div>15%</div> <div>20%</div> </div>
2	A	299	<div> <div>3%</div> <div>82%</div> <div>13%</div> <div>• •</div> </div>
2	B	299	<div> <div>64%</div> <div>10%</div> <div>•</div> <div>25%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	C	299	<div><div></div><div>67%</div><div>7%</div><div>25%</div><div></div></div>
2	D	299	<div>%<div><div></div><div>79%</div><div>15%</div><div></div></div><div></div></div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 9138 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called 5'-R(*CP*CP*GP*GP*CP*UP*CP*UP*UP*UP*CP*AP*GP*AP*GP*CP*CP*GP*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	F	16	Total	C	N	O	P	0	0	0
			333	150	55	113	15			
1	E	16	Total	C	N	O	P	0	0	0
			333	150	55	113	15			

- Molecule 2 is a protein called 3'-5' exonuclease ERI1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	289	Total	C	N	O	S	0	0	0
			2354	1503	395	439	17			
2	B	225	Total	C	N	O	S	0	0	0
			1823	1164	302	343	14			
2	C	225	Total	C	N	O	S	0	0	0
			1823	1164	302	343	14			
2	D	289	Total	C	N	O	S	0	0	0
			2354	1503	395	439	17			


There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	213	LEU	TRP	ENGINEERED	UNP Q8IV48
A	293	ASN	HIS	ENGINEERED	UNP Q8IV48
B	213	LEU	TRP	ENGINEERED	UNP Q8IV48
B	293	ASN	HIS	ENGINEERED	UNP Q8IV48
C	213	LEU	TRP	ENGINEERED	UNP Q8IV48
C	293	ASN	HIS	ENGINEERED	UNP Q8IV48
D	213	LEU	TRP	ENGINEERED	UNP Q8IV48
D	293	ASN	HIS	ENGINEERED	UNP Q8IV48

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	2	Total Mg 2 2	0	0
3	A	2	Total Mg 2 2	0	0
3	D	2	Total Mg 2 2	0	0
3	C	2	Total Mg 2 2	0	0

- Molecule 4 is ADENOSINE MONOPHOSPHATE (three-letter code: AMP) (formula: C₁₀H₁₄N₅O₇P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C N O P 23 10 5 7 1	0	0
4	B	1	Total C N O P 23 10 5 7 1	0	0
4	C	1	Total C N O P 23 10 5 7 1	0	0
4	D	1	Total C N O P 23 10 5 7 1	0	0

- Molecule 5 is water.




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	8	Total O 8 8	0	0
5	B	5	Total O 5 5	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	2	Total	O	0	0
			2	2		
5	D	2	Total	O	0	0
			2	2		
5	F	1	Total	O	0	0
			1	1		

● Molecule 1: 5'-R(*CP*CP*GP*GP*CP*UP*CP*UP*UP*UP*CP*AP*GP*AP*GP*CP*CP*GP*G)-3'



V210	I211	D212	L213	K217	D230	D234	Z236	N240	K256	M259	T260	L284	I305	R308	N319	E320	K321	M322	S330	S331	L334	P335	I336	M344	K348	LYS																				
SER	LVS	PHE	ILE	THR	SER	SER	ALA	SER	D60	F61	S62	Y66	I69	L84	K87	L98	E94	L106	K107	M116	I117	K118	F122	A123	D124	S125	Y126	Y127	D134	F154	H155	V156	V157	L158	L159	N160	T161	H162	T163	I166	F167	D168	T169	Q181	F203	P204

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	50.39Å 195.15Å 87.97Å 90.00° 92.13° 90.00°	Depositor
Resolution (Å)	19.92 – 3.00 19.91 – 3.00	Depositor EDS
% Data completeness (in resolution range)	95.8 (19.92-3.00) 95.7 (19.91-3.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.36 (at 2.98Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.209 , 0.260 0.272 , 0.308	Depositor DCC
R_{free} test set	3231 reflections (9.97%)	DCC
Wilson B-factor (Å ²)	41.9	Xtriage
Anisotropy	0.419	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 5.3	EDS
Estimated twinning fraction	0.056 for h,-k,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 33770 reflections	Xtriage
F_o, F_c correlation	0.84	EDS
Total number of atoms	9138	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.73% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: AMP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	E	0.66	0/370	1.27	3/574 (0.5%)
1	F	0.65	0/370	1.32	3/574 (0.5%)
2	A	0.33	0/2402	0.49	0/3235
2	B	0.33	0/1864	0.49	0/2521
2	C	0.33	0/1864	0.48	0/2521
2	D	0.35	0/2402	0.49	0/3235
All	All	0.37	0/9272	0.61	6/12660 (0.0%)

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	7	G	O3'-P-O5'	-11.54	82.08	104.00
1	E	7	G	O3'-P-O5'	-11.16	82.80	104.00
1	F	7	G	OP1-P-O3'	-9.85	83.52	105.20
1	E	7	G	OP1-P-O3'	-8.45	86.60	105.20
1	E	7	G	OP2-P-O3'	-8.43	86.65	105.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	333	0	173	0	0
1	F	333	0	173	3	0
2	A	2354	0	2381	21	0
2	B	1823	0	1817	13	0
2	C	1823	0	1817	17	0
2	D	2354	0	2381	31	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
3	C	2	0	0	0	0
3	D	2	0	0	0	0
4	A	23	0	12	0	0
4	B	23	0	12	0	0
4	C	23	0	12	0	0
4	D	23	0	12	0	0
5	A	8	0	0	0	0
5	B	5	0	0	0	0
5	C	2	0	0	0	0
5	D	2	0	0	0	0
5	F	1	0	0	0	0
All	All	9138	0	8790	81	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 81 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:317:ARG:HH11	2:C:317:ARG:HG3	1.18	1.03
2:D:259:ASN:H	2:D:319:ASN:HD21	1.13	0.93
2:D:127:TYR:O	2:D:161:THR:HG21	1.82	0.80
2:D:160:ASN:ND2	2:D:163:THR:H	1.81	0.78
2:C:317:ARG:NH1	2:C:317:ARG:HG3	1.97	0.72

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	A	287/299 (96%)	276 (96%)	10 (4%)	1 (0%)	46	84
2	B	223/299 (75%)	218 (98%)	5 (2%)	0	100	100
2	C	223/299 (75%)	217 (97%)	6 (3%)	0	100	100
2	D	287/299 (96%)	277 (96%)	10 (4%)	0	100	100
All	All	1020/1196 (85%)	988 (97%)	31 (3%)	1 (0%)	56	90

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	A	99	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	A	267/276 (97%)	247 (92%)	20 (8%)	17	51
2	B	208/276 (75%)	191 (92%)	17 (8%)	14	46
2	C	208/276 (75%)	201 (97%)	7 (3%)	44	81
2	D	267/276 (97%)	247 (92%)	20 (8%)	17	51
All	All	950/1104 (86%)	886 (93%)	64 (7%)	20	57

5 of 64 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	233	TRP
2	B	339	THR
2	D	240	ASN
2	B	240	ASN
2	B	276	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 20 such

sidechains are listed below:

Mol	Chain	Res	Type
2	C	172	GLN
2	C	205	GLN
2	D	240	ASN
2	B	343	GLN
2	C	143	ASN

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	E	15/20 (75%)	8 (53%)	1 (6%)
1	F	15/20 (75%)	6 (40%)	0
All	All	30/40 (75%)	14 (46%)	1 (3%)

5 of 14 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	F	8	C
1	F	9	U
1	F	12	U
1	F	13	U
1	F	15	C

All (1) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	E	14	U

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 8 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	AMP	A	1002	3	20,25,25	1.13	2 (10%)	22,38,38	2.10	3 (13%)
4	AMP	B	2002	3	20,25,25	1.07	1 (5%)	22,38,38	2.23	4 (18%)
4	AMP	C	3002	3	20,25,25	1.07	1 (5%)	22,38,38	2.26	4 (18%)
4	AMP	D	4002	3	20,25,25	1.12	1 (5%)	22,38,38	2.10	3 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	AMP	A	1002	3	-	0/6/26/26	0/3/3/3
4	AMP	B	2002	3	-	0/6/26/26	0/3/3/3
4	AMP	C	3002	3	-	0/6/26/26	0/3/3/3
4	AMP	D	4002	3	-	0/6/26/26	0/3/3/3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1002	AMP	O4'-C1'	2.11	1.43	1.41
4	C	3002	AMP	C5-C4	3.12	1.47	1.40
4	B	2002	AMP	C5-C4	3.12	1.47	1.40
4	A	1002	AMP	C5-C4	3.25	1.47	1.40
4	D	4002	AMP	C5-C4	3.34	1.48	1.40

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	2002	AMP	N3-C2-N1	-7.47	123.17	128.89
4	A	1002	AMP	N3-C2-N1	-7.34	123.28	128.89
4	C	3002	AMP	N3-C2-N1	-7.31	123.30	128.89
4	D	4002	AMP	N3-C2-N1	-7.14	123.42	128.89
4	C	3002	AMP	C2'-C1'-N9	-5.61	105.73	114.29

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	E	16/20 (80%)	1.20	3 (18%)	2 1	49, 49, 50, 51	0
1	F	16/20 (80%)	0.23	1 (6%)	23 9	49, 50, 50, 51	0
2	A	289/299 (96%)	0.27	10 (3%)	48 21	46, 50, 54, 68	0
2	B	225/299 (75%)	0.01	0	100 100	45, 49, 53, 55	0
2	C	225/299 (75%)	0.06	0	100 100	45, 49, 53, 54	0
2	D	289/299 (96%)	0.08	3 (1%)	84 60	46, 50, 54, 69	0
All	All	1060/1236 (85%)	0.13	17 (1%)	74 47	45, 50, 53, 69	0

The worst 5 of 17 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	6	G	3.5
2	A	88	LEU	3.4
2	D	123	ALA	2.9
2	D	124	ASP	2.7
2	A	100	ASP	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	MG	D	4001	1/1	0.94	0.18	-0.63	50,50,50,50	0
4	AMP	C	3002	23/23	0.95	0.20	-0.72	34,35,36,36	0
4	AMP	D	4002	23/23	0.93	0.21	-0.79	49,50,52,52	0
3	MG	C	3001	1/1	0.98	0.17	-1.09	10,10,10,10	0
4	AMP	B	2002	23/23	0.97	0.17	-1.15	30,32,34,34	0
4	AMP	A	1002	23/23	0.92	0.17	-1.61	46,47,47,48	0
3	MG	B	2001	1/1	0.96	0.13	-2.42	17,17,17,17	0
3	MG	A	1001	1/1	0.92	0.10	-2.99	28,28,28,28	0
3	MG	C	3000	1/1	0.96	0.18	-	14,14,14,14	0
3	MG	B	2000	1/1	0.97	0.20	-	20,20,20,20	0
3	MG	A	1000	1/1	0.96	0.09	-	21,21,21,21	0
3	MG	D	4000	1/1	0.90	0.12	-	38,38,38,38	0

6.5 Other polymers [i](#)

There are no such residues in this entry.