



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 07:03 AM GMT

PDB ID : 2ZC8
Title : Crystal structure of N-Acylamino Acid Racemase from *Thermus thermophilus* HB8
Authors : Hayashida, M.; Kim, S.H.; Takeda, K.; Hisano, T.; Miki, K.
Deposited on : 2007-11-05
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1833 (1.96-1.96)
Clashscore	102246	1953 (1.96-1.96)
Ramachandran outliers	100387	1936 (1.96-1.96)
Sidechain outliers	100360	1936 (1.96-1.96)
RSRZ outliers	91569	1835 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	369	<div> <div>3%</div> <div>90%</div> <div>9%</div> </div>
1	B	369	<div> <div>2%</div> <div>87%</div> <div>11%</div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 6213 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called N-acylamino acid racemase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	368	Total	C	N	O	S	0	0	0
			2872	1840	505	518	9			
1	B	368	Total	C	N	O	S	0	0	0
			2886	1847	508	522	9			

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	215	Total	O	0	0
			215	215		
2	B	240	Total	O	0	0
			240	240		

- Molecule 1: N-acylamino acid racemase



4 Data and refinement statistics

Property	Value	Source
Space group	I 4	Depositor
Cell constants a, b, c, α , β , γ	136.71Å 136.71Å 86.66Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	26.84 – 1.95 30.97 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.5 (26.84-1.95) 99.7 (30.97-1.95)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.88 (at 1.95Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, R_{free}	0.169 , 0.204 0.169 , 0.204	Depositor DCC
R_{free} test set	5900 reflections (10.19%)	DCC
Wilson B-factor (Å ²)	25.8	Xtriage
Anisotropy	0.081	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 53.6	EDS
Estimated twinning fraction	0.017 for -k,-h,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	1 of 57940 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6213	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.69% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.65	0/2931	0.76	0/3969
1	B	0.62	0/2945	0.77	1/3986 (0.0%)
All	All	0.63	0/5876	0.76	1/7955 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	186	THR	N-CA-CB	-5.02	100.76	110.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2872	0	2926	24	0
1	B	2886	0	2945	35	0
2	A	215	0	0	4	0
2	B	240	0	0	4	0
All	All	6213	0	5871	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:12:GLU:HB2	1:A:361:LEU:HD11	1.39	1.03
1:B:276:HIS:HE1	1:B:309:GLY:H	1.11	0.98
1:A:276:HIS:HE1	1:A:309:GLY:H	1.11	0.95
1:B:161:LYS:HA	1:B:186:THR:HG22	1.52	0.91
1:B:276:HIS:CE1	1:B:309:GLY:H	1.96	0.84
1:A:276:HIS:CE1	1:A:309:GLY:H	1.99	0.79
1:A:330:GLU:OE1	1:A:350:HIS:HD2	1.76	0.69
1:B:136:LEU:HD21	1:B:152:HIS:CE1	2.30	0.67
1:B:84:GLU:OE1	1:B:87:ARG:HD3	1.96	0.66
1:B:64:ARG:NH2	1:B:69:GLU:OE1	2.29	0.65
1:B:357:GLU:HG3	2:B:578:HOH:O	1.97	0.65
1:A:16:LYS:HG2	1:A:324:GLU:HG2	1.79	0.65
1:B:161:LYS:CA	1:B:186:THR:HG22	2.26	0.63
1:A:74:ARG:O	1:A:78:ARG:HD3	2.01	0.61
1:B:364:ARG:HD3	1:B:365:TYR:N	2.17	0.60
1:B:174:LYS:O	1:B:178:GLU:HG3	2.01	0.60
1:A:151:ARG:HD3	1:A:151:ARG:O	2.02	0.59
1:A:151:ARG:HD3	1:A:151:ARG:C	2.24	0.58
1:A:257:ARG:NH1	2:A:470:HOH:O	2.38	0.57
1:B:276:HIS:HE1	1:B:309:GLY:N	1.92	0.55
1:B:198:LEU:HD23	1:B:227:LEU:HD12	1.89	0.55
1:B:216:LEU:H	1:B:224:HIS:CE1	2.25	0.54
1:A:266:LEU:O	1:A:271:GLU:HG2	2.08	0.54
1:A:216:LEU:H	1:A:224:HIS:CE1	2.26	0.54
1:B:364:ARG:NE	1:B:366:MET:HG2	2.23	0.54
1:A:350:HIS:HE1	2:A:581:HOH:O	1.91	0.53
1:A:150:GLU:O	1:A:154:GLU:HG3	2.09	0.53
1:B:225:ALA:O	1:B:229:ARG:HG3	2.10	0.52
1:B:177:ARG:HD2	1:B:207:LEU:O	2.10	0.51
1:B:136:LEU:HB2	1:B:162:LEU:HD23	1.93	0.51
1:A:159:ARG:C	1:A:160:ILE:HD12	2.33	0.48
1:A:354:PRO:O	1:A:358:ARG:HG3	2.14	0.48
1:B:276:HIS:CE1	1:B:309:GLY:N	2.76	0.47
1:A:276:HIS:HD2	2:A:469:HOH:O	1.96	0.47
1:B:198:LEU:HD23	1:B:227:LEU:CD1	2.45	0.47
1:B:276:HIS:CE1	1:B:307:LEU:HB3	2.51	0.46
1:B:12:GLU:HB2	1:B:361:LEU:HD11	1.97	0.46
1:B:161:LYS:CB	1:B:186:THR:HG22	2.46	0.45
1:B:336:ASP:HB3	2:B:591:HOH:O	2.16	0.45
1:B:152:HIS:HD2	1:B:155:GLU:OE2	1.99	0.45
1:B:116:GLY:O	1:B:117:ARG:HD3	2.16	0.44
1:B:202:LYS:NZ	2:B:421:HOH:O	2.51	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:20:GLU:HA	1:A:25:VAL:HG22	1.99	0.43
1:B:9:ARG:HH12	1:B:357:GLU:CD	2.22	0.43
1:B:229:ARG:HE	1:B:229:ARG:HB2	1.30	0.43
1:B:97:PRO:O	1:B:267:GLY:HA2	2.19	0.42
1:A:324:GLU:HG3	2:A:452:HOH:O	2.18	0.42
1:B:317:SER:HB2	2:B:512:HOH:O	2.19	0.42
1:B:55:TYR:OH	1:B:265:ARG:HD3	2.19	0.42
1:A:23:PHE:CE1	1:B:74:ARG:NH1	2.86	0.42
1:A:136:LEU:HD21	1:A:152:HIS:CD2	2.54	0.42
1:B:189:ALA:HB3	1:B:215:PRO:HA	2.02	0.42
1:A:279:ALA:CB	1:A:286:LEU:HD21	2.49	0.42
1:B:329:GLU:HG2	1:B:352:LYS:HG2	2.02	0.41
1:A:276:HIS:CE1	1:A:307:LEU:HB3	2.55	0.41
1:A:91:ALA:N	1:A:92:PRO:CD	2.84	0.41
1:B:161:LYS:HD3	1:B:188:ASP:HB2	2.03	0.40
1:A:195:LEU:HD23	1:A:195:LEU:HA	1.94	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	366/369 (99%)	358 (98%)	8 (2%)	0	100	100
1	B	366/369 (99%)	358 (98%)	8 (2%)	0	100	100
All	All	732/738 (99%)	716 (98%)	16 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	294/298 (99%)	291 (99%)	3 (1%)	82	80
1	B	297/298 (100%)	292 (98%)	5 (2%)	68	63
All	All	591/596 (99%)	583 (99%)	8 (1%)	74	70

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	64	ARG
1	A	108	PHE
1	A	151	ARG
1	B	64	ARG
1	B	108	PHE
1	B	186	THR
1	B	198	LEU
1	B	364	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	26	GLN
1	A	152	HIS
1	A	224	HIS
1	A	276	HIS
1	A	301	ASN
1	A	350	HIS
1	B	121	GLN
1	B	152	HIS
1	B	224	HIS
1	B	276	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	368/369 (99%)	-0.09	12 (3%) 50 61	16, 25, 44, 75	0
1	B	368/369 (99%)	-0.05	7 (1%) 70 78	17, 26, 41, 53	0
All	All	736/738 (99%)	-0.07	19 (2%) 59 69	16, 26, 43, 75	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	23	PHE	4.8
1	A	25	VAL	4.0
1	A	79	ASP	3.2
1	A	22	SER	3.2
1	A	20	GLU	2.9
1	A	19	PHE	2.7
1	B	154	GLU	2.6
1	B	147	ARG	2.4
1	B	151	ARG	2.3
1	A	21	THR	2.3
1	A	24	GLY	2.3
1	B	150	GLU	2.2
1	A	355	PHE	2.2
1	A	15	LEU	2.2
1	B	101	ALA	2.1
1	B	335	LYS	2.1
1	A	147	ARG	2.1
1	A	17	PHE	2.0
1	B	336	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.