



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 20, 2016 – 11:44 AM EDT

PDB ID : 4ZE5  
Title : Structure of Gan1D-E170Q, a catalytic mutant of a putative 6-phospho-beta-galactosidase from Geobacillus stearothermophilus  
Authors : Lansky, S.; Zehavi, A.; Dvir, H.; Shoham, Y.; Shoham, G.  
Deposited on : 2015-04-20  
Resolution : 1.48 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20027790  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027790


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.48 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3129 (1.50-1.46)
Clashscore	102246	3380 (1.50-1.46)
Ramachandran outliers	100387	3310 (1.50-1.46)
Sidechain outliers	100360	3308 (1.50-1.46)
RSRZ outliers	91569	3133 (1.50-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	485	<div> <div>4%</div> <div> <div></div> <div>77%</div> <div>16%</div> <div>• 5%</div> </div> </div>
1	B	485	<div> <div>2%</div> <div> <div></div> <div>79%</div> <div>14%</div> <div>• 5%</div> </div> </div>
1	C	485	<div> <div>2%</div> <div> <div></div> <div>77%</div> <div>16%</div> <div>• 5%</div> </div> </div>
1	D	485	<div> <div>2%</div> <div> <div></div> <div>75%</div> <div>18%</div> <div>• 5%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	504	-	-	-	X
2	GOL	C	501	-	-	-	X
3	IMD	A	505	-	-	X	-

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 18228 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative 6-phospho-beta-galactobiosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	459	Total	C	N	O	S	0	21	0
			3902	2518	666	703	15			
1	B	459	Total	C	N	O	S	0	20	0
			3900	2515	662	709	14			
1	C	460	Total	C	N	O	S	0	23	0
			3923	2530	674	705	14			
1	D	460	Total	C	N	O	S	0	16	0
			3872	2497	664	699	12			

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-6	MET	-	initiating methionine	UNP W8QF82
A	-5	ILE	-	expression tag	UNP W8QF82
A	-4	HIS	-	expression tag	UNP W8QF82
A	-3	HIS	-	expression tag	UNP W8QF82
A	-2	HIS	-	expression tag	UNP W8QF82
A	-1	HIS	-	expression tag	UNP W8QF82
A	0	HIS	-	expression tag	UNP W8QF82
A	1	HIS	-	expression tag	UNP W8QF82
A	170	GLN	GLU	engineered mutation	UNP W8QF82
B	-6	MET	-	initiating methionine	UNP W8QF82
B	-5	ILE	-	expression tag	UNP W8QF82
B	-4	HIS	-	expression tag	UNP W8QF82
B	-3	HIS	-	expression tag	UNP W8QF82
B	-2	HIS	-	expression tag	UNP W8QF82
B	-1	HIS	-	expression tag	UNP W8QF82
B	0	HIS	-	expression tag	UNP W8QF82
B	1	HIS	-	expression tag	UNP W8QF82
B	170	GLN	GLU	engineered mutation	UNP W8QF82
C	-6	MET	-	initiating methionine	UNP W8QF82
C	-5	ILE	-	expression tag	UNP W8QF82
C	-4	HIS	-	expression tag	UNP W8QF82

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-3	HIS	-	expression tag	UNP W8QF82
C	-2	HIS	-	expression tag	UNP W8QF82
C	-1	HIS	-	expression tag	UNP W8QF82
C	0	HIS	-	expression tag	UNP W8QF82
C	1	HIS	-	expression tag	UNP W8QF82
C	170	GLN	GLU	engineered mutation	UNP W8QF82
D	-6	MET	-	initiating methionine	UNP W8QF82
D	-5	ILE	-	expression tag	UNP W8QF82
D	-4	HIS	-	expression tag	UNP W8QF82
D	-3	HIS	-	expression tag	UNP W8QF82
D	-2	HIS	-	expression tag	UNP W8QF82
D	-1	HIS	-	expression tag	UNP W8QF82
D	0	HIS	-	expression tag	UNP W8QF82
D	1	HIS	-	expression tag	UNP W8QF82
D	170	GLN	GLU	engineered mutation	UNP W8QF82

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	C	1	Total	C	O	0	0
			6	3	3		
2	C	1	Total	C	O	0	0
			6	3	3		
2	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is IMIDAZOLE (three-letter code: IMD) (formula:  $C_3H_5N_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	N	0	0
			5	3	2		
3	C	1	Total	C	N	0	0
			5	3	2		
3	D	1	Total	C	N	0	0
			5	3	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	616	Total 616	O 616	0	0
4	B	640	Total 640	O 640	0	0
4	C	690	Total 690	O 690	0	0
4	D	610	Total 610	O 610	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Putative 6-phospho-beta-galactobiosidase




- Molecule 1: Putative 6-phospho-beta-galactobiosidase





- Molecule 1: Putative 6-phospho-beta-galactobiosidase







• Molecule 1: Putative 6-phospho-beta-galactobiosidase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	101.55Å 97.44Å 105.24Å 90.00° 97.82° 90.00°	Depositor
Resolution (Å)	27.64 – 1.48 27.62 – 1.48	Depositor EDS
% Data completeness (in resolution range)	99.7 (27.64-1.48) 99.7 (27.62-1.48)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.50 (at 1.48Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, $R_{free}$	0.132 , 0.159 0.132 , 0.159	Depositor DCC
$R_{free}$ test set	16966 reflections (5.32%)	DCC
Wilson B-factor (Å <sup>2</sup> )	10.9	Xtriage
Anisotropy	0.482	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 50.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.000 for l,-k,h	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	18228	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 71.27 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 2.7039e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, IMD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.44	26/4091 (0.6%)	1.48	65/5559 (1.2%)
1	B	1.43	23/4076 (0.6%)	1.52	73/5543 (1.3%)
1	C	1.42	28/4117 (0.7%)	1.51	76/5597 (1.4%)
1	D	1.40	20/4041 (0.5%)	1.57	72/5496 (1.3%)
All	All	1.42	97/16325 (0.6%)	1.52	286/22195 (1.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	2
All	All	0	4

The worst 5 of 97 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	286	GLU	CD-OE2	10.28	1.36	1.25
1	C	238[A]	GLU	CD-OE1	9.94	1.36	1.25
1	C	238[B]	GLU	CD-OE1	9.94	1.36	1.25
1	B	238[A]	GLU	CG-CD	9.45	1.66	1.51
1	B	238[B]	GLU	CG-CD	9.45	1.66	1.51

The worst 5 of 286 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	180[A]	ARG	NE-CZ-NH2	-14.57	113.02	120.30
1	B	180[B]	ARG	NE-CZ-NH2	-14.57	113.02	120.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	458	ARG	NE-CZ-NH1	13.48	127.04	120.30
1	D	403	ARG	NE-CZ-NH1	13.33	126.96	120.30
1	C	411	ARG	NE-CZ-NH2	-12.60	114.00	120.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	476	ALA	Peptide
1	A	477	GLU	Peptide
1	B	349[A]	ASN	Peptide
1	B	349[B]	ASN	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3902	0	3768	34	0
1	B	3900	0	3731	22	0
1	C	3923	0	3783	38	0
1	D	3872	0	3716	46	0
2	A	24	0	32	3	0
2	B	18	0	24	0	0
2	C	12	0	16	0	0
2	D	6	0	8	0	0
3	A	5	0	5	5	0
3	C	5	0	5	0	0
3	D	5	0	5	1	0
4	A	616	0	0	15	0
4	B	640	0	0	9	0
4	C	690	0	0	21	0
4	D	610	0	0	15	0
All	All	18228	0	15093	139	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 139 close contacts within the same asymmetric unit are listed below, sorted by

their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:132[A]:MET:HE3	4:A:1057:HOH:O	1.42	1.20
1:D:132[A]:MET:HE3	4:D:1075:HOH:O	1.53	1.06
1:A:453:GLU:OE1	3:A:505:IMD:H2	1.59	1.01
1:C:132[A]:MET:HE3	4:C:1110:HOH:O	1.60	1.01
1:B:349[B]:ASN:O	1:B:350[B]:TRP:CD1	2.15	0.99

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	478/485 (99%)	462 (97%)	16 (3%)	0	100	100
1	B	476/485 (98%)	461 (97%)	15 (3%)	0	100	100
1	C	480/485 (99%)	462 (96%)	17 (4%)	1 (0%)	52	24
1	D	472/485 (97%)	456 (97%)	14 (3%)	2 (0%)	39	14
All	All	1906/1940 (98%)	1841 (97%)	62 (3%)	3 (0%)	52	24

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	310	PRO
1	C	310	PRO
1	D	167	THR

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	413/412 (100%)	409 (99%)	4 (1%)	82	60
1	B	411/412 (100%)	408 (99%)	3 (1%)	88	72
1	C	415/412 (101%)	411 (99%)	4 (1%)	82	60
1	D	407/412 (99%)	404 (99%)	3 (1%)	88	72
All	All	1646/1648 (100%)	1632 (99%)	14 (1%)	86	64

5 of 14 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	350[B]	TRP
1	C	93	ASN
1	D	93	ASN
1	B	350[A]	TRP
1	C	430	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 51 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	252	HIS
1	C	169	ASN
1	D	252	HIS
1	B	269	ASN
1	C	170	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	GOL	A	501	-	5,5,5	0.76	0	5,5,5	1.15	0
2	GOL	A	502	-	5,5,5	0.40	0	5,5,5	0.94	1 (20%)
2	GOL	A	503	-	5,5,5	0.45	0	5,5,5	1.04	0
2	GOL	A	504	-	5,5,5	0.60	0	5,5,5	0.95	0
3	IMD	A	505	-	3,5,5	0.62	0	4,5,5	1.61	1 (25%)
2	GOL	B	501	-	5,5,5	0.50	0	5,5,5	1.48	2 (40%)
2	GOL	B	502	-	5,5,5	0.66	0	5,5,5	0.82	0
2	GOL	B	503	-	5,5,5	0.47	0	5,5,5	0.81	0
2	GOL	C	501	-	5,5,5	0.63	0	5,5,5	1.01	1 (20%)
2	GOL	C	502	-	5,5,5	0.48	0	5,5,5	0.84	0
3	IMD	C	503	-	3,5,5	0.16	0	4,5,5	1.33	1 (25%)
2	GOL	D	501	-	5,5,5	0.44	0	5,5,5	0.77	0
3	IMD	D	502	-	3,5,5	0.41	0	4,5,5	0.61	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	501	-	-	0/4/4/4	0/0/0/0
2	GOL	A	502	-	-	0/4/4/4	0/0/0/0
2	GOL	A	503	-	-	0/4/4/4	0/0/0/0
2	GOL	A	504	-	-	0/4/4/4	0/0/0/0
3	IMD	A	505	-	-	0/0/0/0	0/1/1/1
2	GOL	B	501	-	-	0/4/4/4	0/0/0/0
2	GOL	B	502	-	-	0/4/4/4	0/0/0/0
2	GOL	B	503	-	-	0/4/4/4	0/0/0/0
2	GOL	C	501	-	-	0/4/4/4	0/0/0/0
2	GOL	C	502	-	-	0/4/4/4	0/0/0/0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	IMD	C	503	-	-	0/0/0/0	0/1/1/1
2	GOL	D	501	-	-	0/4/4/4	0/0/0/0
3	IMD	D	502	-	-	0/0/0/0	0/1/1/1

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	GOL	C3-C2-C1	-2.38	101.05	111.06
2	C	501	GOL	C3-C2-C1	-2.00	102.62	111.06
3	C	503	IMD	C5-C4-N3	2.02	111.46	107.75
2	A	502	GOL	O3-C3-C2	2.05	120.38	109.97
3	A	505	IMD	C4-C5-N1	2.06	111.55	107.75

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	504	GOL	3	0
3	A	505	IMD	5	0
3	D	502	IMD	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	459/485 (94%)	-0.14	17 (3%) 45 47	6, 11, 28, 67	0
1	B	459/485 (94%)	-0.21	12 (2%) 59 62	6, 11, 24, 61	0
1	C	460/485 (94%)	-0.21	10 (2%) 65 68	5, 9, 22, 80	0
1	D	460/485 (94%)	-0.14	12 (2%) 59 62	5, 11, 25, 90	0
All	All	1838/1940 (94%)	-0.17	51 (2%) 56 59	5, 11, 25, 90	0

The worst 5 of 51 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	350	TRP	9.6
1	D	312	ASP	8.5
1	D	329	THR	7.5
1	C	313	GLY	7.2
1	D	313	GLY	6.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	GOL	C	501	6/6	0.92	0.20	17.77	17,26,33,44	0
2	GOL	A	504	6/6	0.90	0.18	6.15	38,40,43,47	0
2	GOL	A	501	6/6	0.96	0.11	1.93	13,15,20,22	0
2	GOL	B	501	6/6	0.95	0.11	1.50	13,17,22,25	0
2	GOL	B	502	6/6	0.96	0.10	1.28	13,15,21,23	0
3	IMD	A	505	5/5	0.87	0.10	0.97	33,38,42,45	0
3	IMD	D	502	5/5	0.69	0.16	0.59	19,25,31,32	5
2	GOL	A	502	6/6	0.97	0.10	0.42	13,17,20,22	0
3	IMD	C	503	5/5	0.88	0.21	-	22,26,31,35	0
2	GOL	D	501	6/6	0.84	0.18	-	34,35,37,40	0
2	GOL	B	503	6/6	0.85	0.19	-	29,34,35,35	0
2	GOL	A	503	6/6	0.87	0.16	-	31,32,35,35	0
2	GOL	C	502	6/6	0.92	0.17	-	30,31,32,33	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.