



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 02:24 PM GMT

PDB ID : 3ZF4
Title : Phage dUTPases control transfer of virulence genes by a proto- oncogenic G protein-like mechanism. (Staphylococcus bacteriophage 80alpha dUTPase Y81A mutant with dUpNHpp).
Authors : Tormo-Mas, M.A.; Donderis, J.; Garcia-Caballer, M.; Alt, A.; Mir-Sanchis, I.; Marina, A.; Penades, J.R.
Deposited on : 2012-12-10
Resolution : 3.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1114 (3.14-3.06)
Clashscore	102246	1222 (3.14-3.06)
Ramachandran outliers	100387	1174 (3.14-3.06)
Sidechain outliers	100360	1174 (3.14-3.06)
RSRZ outliers	91569	1119 (3.14-3.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	204	 69% 10% • 20%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 1318 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DUTPASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	164	Total	C	N	O	S	0	0	0
			1275	803	222	248	2			

There are 35 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-33	MET	-	EXPRESSION TAG	UNP A4ZF98
A	-32	GLY	-	EXPRESSION TAG	UNP A4ZF98
A	-31	SER	-	EXPRESSION TAG	UNP A4ZF98
A	-30	SER	-	EXPRESSION TAG	UNP A4ZF98
A	-29	HIS	-	EXPRESSION TAG	UNP A4ZF98
A	-28	HIS	-	EXPRESSION TAG	UNP A4ZF98
A	-27	HIS	-	EXPRESSION TAG	UNP A4ZF98
A	-26	HIS	-	EXPRESSION TAG	UNP A4ZF98
A	-25	HIS	-	EXPRESSION TAG	UNP A4ZF98
A	-24	HIS	-	EXPRESSION TAG	UNP A4ZF98
A	-23	SER	-	EXPRESSION TAG	UNP A4ZF98
A	-22	SER	-	EXPRESSION TAG	UNP A4ZF98
A	-21	GLY	-	EXPRESSION TAG	UNP A4ZF98
A	-20	LEU	-	EXPRESSION TAG	UNP A4ZF98
A	-19	VAL	-	EXPRESSION TAG	UNP A4ZF98
A	-18	PRO	-	EXPRESSION TAG	UNP A4ZF98
A	-17	ARG	-	EXPRESSION TAG	UNP A4ZF98
A	-16	GLY	-	EXPRESSION TAG	UNP A4ZF98
A	-15	SER	-	EXPRESSION TAG	UNP A4ZF98
A	-14	HIS	-	EXPRESSION TAG	UNP A4ZF98
A	-13	MET	-	EXPRESSION TAG	UNP A4ZF98
A	-12	ALA	-	EXPRESSION TAG	UNP A4ZF98
A	-11	SER	-	EXPRESSION TAG	UNP A4ZF98
A	-10	MET	-	EXPRESSION TAG	UNP A4ZF98
A	-9	THR	-	EXPRESSION TAG	UNP A4ZF98
A	-8	GLY	-	EXPRESSION TAG	UNP A4ZF98
A	-7	GLY	-	EXPRESSION TAG	UNP A4ZF98

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-6	GLN	-	EXPRESSION TAG	UNP A4ZF98
A	-5	GLN	-	EXPRESSION TAG	UNP A4ZF98
A	-4	MET	-	EXPRESSION TAG	UNP A4ZF98
A	-3	GLY	-	EXPRESSION TAG	UNP A4ZF98
A	-2	ARG	-	EXPRESSION TAG	UNP A4ZF98
A	-1	GLY	-	EXPRESSION TAG	UNP A4ZF98
A	0	SER	-	EXPRESSION TAG	UNP A4ZF98
A	84	ALA	TYR	ENGINEERED MUTATION	UNP A4ZF98

- Molecule 2 is 2'-DEOXYURIDINE 5'-ALPHA,BETA-IMIDO-TRIPHOSPHATE (three-letter code: DUP) (formula: $C_9H_{16}N_3O_{13}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			28	9	3	13	3		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		

- Molecule 4 is NICKEL (II) ION (three-letter code: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Ni	0	0
			2	2		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	12	Total	O	0	0
			12	12		

- Molecule 1: DUTPASE

F105	L106	R107	M108	E112	K113	I114	F115	Y122	K123	L124	I129	Q150	E155	SER	VAL	SER	SER	E159	K163	F165	G169	VAL	MET	GLY	SER	SER	HIS	HIS	HIS	HIS	SER	SER	GLY	LEU	VAL	PRO	ARG	GLY	SER	HIS	SER	ALA	SER	MET	THR	GLY	GLY	GLN	GLN	MET	GLY	ARG	GLY	SER	SER	MET	T2	K12	S31	T34	T62	S63	R64	V67	L73	V74	I75	F76	I77	G78	K79	I92	I104
------	------	------	------	------	------	------	------	------	------	------	------	------	------	-----	-----	-----	-----	------	------	------	------	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	------

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 3	Depositor
Cell constants a, b, c, α , β , γ	87.47Å 87.47Å 87.47Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	61.85 – 3.10 61.85 – 3.10	Depositor EDS
% Data completeness (in resolution range)	99.5 (61.85-3.10) 99.6 (61.85-3.10)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.49 (at 3.13Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.240 , 0.291 0.239 , 0.288	Depositor DCC
R_{free} test set	521 reflections (14.12%)	DCC
Wilson B-factor (Å ²)	94.5	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.26 , 54.3	EDS
Estimated twinning fraction	0.068 for l,-k,h	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 4213 reflections	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	1318	wwPDB-VP
Average B, all atoms (Å ²)	106.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DUP, NI, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.27	0/1291	0.47	0/1737

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1275	0	1264	11	0
2	A	28	0	12	0	0
3	A	1	0	0	0	0
4	A	2	0	0	0	0
5	A	12	0	0	1	0
All	All	1318	0	1276	11	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 11 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:31:SER:HA	5:A:2004:HOH:O	1.94	0.67
1:A:104:ILE:H	1:A:104:ILE:HD13	1.76	0.51
1:A:108:ASN:HD21	1:A:112:GLU:HB2	1.78	0.47
1:A:62:THR:HB	1:A:77:THR:HG23	1.96	0.46
1:A:63:SER:HB3	1:A:75:ILE:HB	1.99	0.45

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	158/204 (78%)	150 (95%)	8 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	136/174 (78%)	127 (93%)	9 (7%)	21	56

5 of 9 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	104	ILE
1	A	150	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	115	PHE
1	A	79	LYS
1	A	106	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	13	ASN
1	A	72	HIS
1	A	102	GLN
1	A	111	ASN
1	A	150	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 3 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	DUP	A	777	3	23,29,29	1.68	4 (17%)	32,45,45	1.76	3 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DUP	A	777	3	-	0/13/34/34	0/2/2/2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	777	DUP	PA-O1A	2.07	1.48	1.46
2	A	777	DUP	PB-O1B	2.09	1.48	1.46
2	A	777	DUP	PB-N3A	4.45	1.75	1.63
2	A	777	DUP	PA-N3A	4.49	1.75	1.63

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	777	DUP	PG-O3B-PB	-3.41	121.24	132.67
2	A	777	DUP	O4'-C1'-N1	2.18	111.49	107.72
2	A	777	DUP	C4-N3-C2	7.21	121.28	114.14

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	164/204 (80%)	-0.08	0 100 100	74, 106, 138, 148	2 (1%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	DUP	A	777	28/28	0.96	0.15	-0.89	96,99,100,101	0
3	MG	A	888	1/1	0.94	0.12	-1.53	84,84,84,84	0
4	NI	A	1171	1/1	0.94	0.10	-	137,137,137,137	1
4	NI	A	1170	1/1	0.98	0.15	-	107,107,107,107	1

6.5 Other polymers [i](#)

There are no such residues in this entry.