



wwPDB X-ray Structure Validation Summary Report ⓘ

May 23, 2016 – 12:45 PM EDT

PDB ID : 4ZF3
Title : Crystal structure of Green Fluorescent Protein (GFP); S65T, H148D; circular permutant (50-51)
Authors : Oltrogge, L.M.; Boxer, S.G.
Deposited on : 2015-04-21
Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027674
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027674


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4755 (1.90-1.90)
Clashscore	102246	5398 (1.90-1.90)
Ramachandran outliers	100387	5338 (1.90-1.90)
Sidechain outliers	100360	5339 (1.90-1.90)
RSRZ outliers	91569	4766 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	252	<div> <div>2%</div> <div>80%</div> <div>6%</div> <div>14%</div> </div>
1	B	252	<div> <div>13%</div> <div>73%</div> <div>12%</div> <div>15%</div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3414 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Green fluorescent protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	217	Total	C	N	O	S	0	0	0
			1711	1091	291	325	4			
1	B	215	Total	C	N	O	S	0	0	0
			1677	1070	279	324	4			

There are 88 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-10	MET	-	initiating methionine	UNP P42212
A	-9	GLY	-	expression tag	UNP P42212
A	-8	HIS	-	expression tag	UNP P42212
A	-7	HIS	-	expression tag	UNP P42212
A	-6	HIS	-	expression tag	UNP P42212
A	-5	HIS	-	expression tag	UNP P42212
A	-4	HIS	-	expression tag	UNP P42212
A	-3	HIS	-	expression tag	UNP P42212
A	-2	SER	-	expression tag	UNP P42212
A	-1	SER	-	expression tag	UNP P42212
A	0	GLY	-	expression tag	UNP P42212
A	14	LEU	PHE	engineered mutation	UNP P42212
A	15	CRO	SER	chromophore	UNP P42212
A	?	CRO	TYR	chromophore	UNP P42212
A	?	CRO	GLY	chromophore	UNP P42212
A	30	ARG	GLN	engineered mutation	UNP P42212
A	49	SER	PHE	engineered mutation	UNP P42212
A	55	LYS	ASN	engineered mutation	UNP P42212
A	61	VAL	GLU	engineered mutation	UNP P42212
A	78	THR	ILE	engineered mutation	UNP P42212
A	95	PHE	TYR	engineered mutation	UNP P42212
A	98	ASP	HIS	engineered mutation	UNP P42212
A	103	THR	MET	engineered mutation	UNP P42212
A	106	ASN	LYS	engineered mutation	UNP P42212
A	113	ALA	VAL	engineered mutation	UNP P42212

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	116	THR	LYS	engineered mutation	UNP P42212
A	117	VAL	ILE	engineered mutation	UNP P42212
A	121	VAL	ILE	engineered mutation	UNP P42212
A	155	THR	SER	engineered mutation	UNP P42212
A	156	VAL	ALA	engineered mutation	UNP P42212
A	186	LEU	LEU	linker	UNP P42212
A	187	TYR	TYR	linker	UNP P42212
A	188	GLY	-	linker	UNP P42212
A	189	GLY	-	linker	UNP P42212
A	190	THR	-	linker	UNP P42212
A	191	GLY	-	linker	UNP P42212
A	192	GLY	-	linker	UNP P42212
A	193	SER	-	linker	UNP P42212
A	194	ALA	-	linker	UNP P42212
A	195	SER	-	linker	UNP P42212
A	196	GLN	-	linker	UNP P42212
A	223	ARG	SER	engineered mutation	UNP P42212
A	232	ILE	TYR	engineered mutation	UNP P42212
A	241	SER	CYS	engineered mutation	UNP P42212
B	-10	MET	-	initiating methionine	UNP P42212
B	-9	GLY	-	expression tag	UNP P42212
B	-8	HIS	-	expression tag	UNP P42212
B	-7	HIS	-	expression tag	UNP P42212
B	-6	HIS	-	expression tag	UNP P42212
B	-5	HIS	-	expression tag	UNP P42212
B	-4	HIS	-	expression tag	UNP P42212
B	-3	HIS	-	expression tag	UNP P42212
B	-2	SER	-	expression tag	UNP P42212
B	-1	SER	-	expression tag	UNP P42212
B	0	GLY	-	expression tag	UNP P42212
B	14	LEU	PHE	engineered mutation	UNP P42212
B	15	CRO	SER	chromophore	UNP P42212
B	?	CRO	TYR	chromophore	UNP P42212
B	?	CRO	GLY	chromophore	UNP P42212
B	30	ARG	GLN	engineered mutation	UNP P42212
B	49	SER	PHE	engineered mutation	UNP P42212
B	55	LYS	ASN	engineered mutation	UNP P42212
B	61	VAL	GLU	engineered mutation	UNP P42212
B	78	THR	ILE	engineered mutation	UNP P42212
B	95	PHE	TYR	engineered mutation	UNP P42212
B	98	ASP	HIS	engineered mutation	UNP P42212
B	103	THR	MET	engineered mutation	UNP P42212

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	106	ASN	LYS	engineered mutation	UNP P42212
B	113	ALA	VAL	engineered mutation	UNP P42212
B	116	THR	LYS	engineered mutation	UNP P42212
B	117	VAL	ILE	engineered mutation	UNP P42212
B	121	VAL	ILE	engineered mutation	UNP P42212
B	155	THR	SER	engineered mutation	UNP P42212
B	156	VAL	ALA	engineered mutation	UNP P42212
B	186	LEU	LEU	linker	UNP P42212
B	187	TYR	TYR	linker	UNP P42212
B	188	GLY	-	linker	UNP P42212
B	189	GLY	-	linker	UNP P42212
B	190	THR	-	linker	UNP P42212
B	191	GLY	-	linker	UNP P42212
B	192	GLY	-	linker	UNP P42212
B	193	SER	-	linker	UNP P42212
B	194	ALA	-	linker	UNP P42212
B	195	SER	-	linker	UNP P42212
B	196	GLN	-	linker	UNP P42212
B	223	ARG	SER	engineered mutation	UNP P42212
B	232	ILE	TYR	engineered mutation	UNP P42212
B	241	SER	CYS	engineered mutation	UNP P42212

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	18	Total O 18 18	0	0
2	B	8	Total O 8 8	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	46.37Å 68.26Å 57.87Å 90.00° 102.15° 90.00°	Depositor
Resolution (Å)	34.31 – 1.90 34.31 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.1 (34.31-1.90) 99.2 (34.31-1.90)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.84 (at 1.89Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.215 , 0.261 0.215 , 0.260	Depositor DCC
R_{free} test set	1391 reflections (5.03%)	DCC
Wilson B-factor (Å ²)	30.1	Xtriage
Anisotropy	0.567	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.46 , 64.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3414	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.50% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CRO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.44	0/1722	0.59	0/2327
1	B	0.37	0/1688	0.55	0/2287
All	All	0.40	0/3410	0.57	0/4614

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1711	0	1663	12	0
1	B	1677	0	1605	21	0
2	A	18	0	0	0	0
2	B	8	0	0	0	0
All	All	3414	0	3268	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 29 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:195:SER:OG	1:A:195:SER:O	2.09	0.70
1:A:156:VAL:HG13	1:B:156:VAL:HG13	1.75	0.68
1:A:171:LEU:HD22	1:B:171:LEU:HD23	1.84	0.59
1:A:171:LEU:HD21	1:A:173:PHE:HE2	1.68	0.59
1:B:26:ASP:HA	1:B:29:LYS:HG2	1.85	0.58

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	206/252 (82%)	199 (97%)	7 (3%)	0	100	100
1	B	204/252 (81%)	197 (97%)	6 (3%)	1 (0%)	34	21
All	All	410/504 (81%)	396 (97%)	13 (3%)	1 (0%)	52	42

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	84	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	184/215 (86%)	181 (98%)	3 (2%)	70	66

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	B	179/215 (83%)	174 (97%)	5 (3%)	51 41
All	All	363/430 (84%)	355 (98%)	8 (2%)	60 53

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	27	HIS
1	B	232	ILE
1	B	95	PHE
1	A	237	LEU
1	B	83	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CRO	A	15	1	23,23,24	2.86	9 (39%)	30,32,34	1.75	8 (26%)
1	CRO	B	15	1	23,23,24	2.91	9 (39%)	30,32,34	1.66	7 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CRO	A	15	1	-	0/12/31/32	0/2/2/2
1	CRO	B	15	1	-	0/12/31/32	0/2/2/2

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	15	CRO	OH-CZ	-4.76	1.25	1.37
1	B	15	CRO	OH-CZ	-4.62	1.26	1.37
1	A	15	CRO	CG2-CB2	-2.55	1.41	1.46
1	B	15	CRO	CA1-C1	-2.28	1.47	1.51
1	A	15	CRO	CA1-C1	-2.07	1.48	1.51

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	15	CRO	CA3-N3-C1	-2.98	124.09	127.38
1	B	15	CRO	CE2-CZ-CE1	-2.84	115.88	119.78
1	A	15	CRO	CE2-CZ-CE1	-2.62	116.18	119.78
1	A	15	CRO	CA1-C1-N3	-2.14	122.39	124.98
1	B	15	CRO	OG1-CB1-CG1	-2.10	103.31	109.56

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	216/252 (85%)	0.31	6 (2%) 56 60	19, 31, 48, 61	0
1	B	214/252 (84%)	0.94	33 (15%) 3 3	24, 44, 61, 68	0
All	All	430/504 (85%)	0.62	39 (9%) 11 13	19, 37, 59, 68	0

The worst 5 of 39 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	144	LEU	7.5
1	B	179	ILE	6.2
1	B	145	LEU	6.2
1	A	179	ILE	5.5
1	B	197	GLY	4.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	CRO	A	15	22/23	0.97	0.12	-	19,22,26,27	0
1	CRO	B	15	22/23	0.93	0.13	-	25,32,36,41	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.