



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 19, 2016 – 09:18 PM GMT

PDB ID : 4ZLW
Title : Crystal structure of the PmFTN variant E130A soaked in iron (overnight)
Authors : Pfaffen, S.; Murphy, M.E.P.
Deposited on : 2015-05-01
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026982
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20026982


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	168	<div> <div>2%</div> <div>90% 7%</div> </div>
1	B	168	<div> <div>4%</div> <div>90% 6%</div> </div>
1	C	168	<div> <div>4%</div> <div>83% 10% 7%</div> </div>
1	D	168	<div> <div>2%</div> <div>88% 5% 7%</div> </div>
1	E	168	<div> <div>5%</div> <div>83% 10% 6%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	168	 4% 82% 10% • 7%
1	G	168	 3% 81% 11% • 7%
1	H	168	 2% 85% 8% • 7%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	FE	A	202	-	-	-	X
2	FE	E	204	-	-	-	X
2	FE	G	202	-	-	-	X
2	FE	G	203	-	-	-	X

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 10566 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ferritin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	157	Total	C	N	O	S	0	3	0
			1263	793	211	253	6			
1	D	157	Total	C	N	O	S	0	2	0
			1249	786	208	249	6			
1	B	158	Total	C	N	O	S	0	3	0
			1270	800	211	253	6			
1	C	157	Total	C	N	O	S	0	2	0
			1255	789	210	250	6			
1	E	158	Total	C	N	O	S	0	2	0
			1255	789	209	251	6			
1	F	157	Total	C	N	O	S	0	1	0
			1243	783	207	248	5			
1	G	157	Total	C	N	O	S	0	1	0
			1247	790	207	245	5			
1	H	157	Total	C	N	O	S	0	0	0
			1237	780	206	246	5			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP B6DMH6
A	130	ALA	GLU	engineered mutation	UNP B6DMH6
D	0	GLY	-	expression tag	UNP B6DMH6
D	130	ALA	GLU	engineered mutation	UNP B6DMH6
B	0	GLY	-	expression tag	UNP B6DMH6
B	130	ALA	GLU	engineered mutation	UNP B6DMH6
C	0	GLY	-	expression tag	UNP B6DMH6
C	130	ALA	GLU	engineered mutation	UNP B6DMH6
E	0	GLY	-	expression tag	UNP B6DMH6
E	130	ALA	GLU	engineered mutation	UNP B6DMH6
F	0	GLY	-	expression tag	UNP B6DMH6
F	130	ALA	GLU	engineered mutation	UNP B6DMH6
G	0	GLY	-	expression tag	UNP B6DMH6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	130	ALA	GLU	engineered mutation	UNP B6DMH6
H	0	GLY	-	expression tag	UNP B6DMH6
H	130	ALA	GLU	engineered mutation	UNP B6DMH6

- Molecule 2 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	G	5	Total Fe 5 5	0	0
2	D	5	Total Fe 5 5	0	0
2	E	5	Total Fe 5 5	0	0
2	H	4	Total Fe 4 4	0	0
2	B	5	Total Fe 5 5	0	0
2	C	4	Total Fe 4 4	0	0
2	A	4	Total Fe 4 4	0	0
2	F	5	Total Fe 5 5	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	55	Total O 55 55	0	0
3	D	63	Total O 63 63	0	0
3	B	59	Total O 59 59	0	0
3	C	71	Total O 71 71	0	0
3	E	61	Total O 61 61	0	0
3	F	57	Total O 57 57	0	0
3	G	74	Total O 74 74	0	0
3	H	70	Total O 70 70	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ferritin




- Molecule 1: Ferritin




- Molecule 1: Ferritin




- Molecule 1: Ferritin



- Molecule 1: Ferritin





- Molecule 1: Ferritin







• Molecule 1: Ferritin



• Molecule 1: Ferritin



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 3	Depositor
Cell constants a, b, c, α , β , γ	175.05Å 175.05Å 175.05Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.55 – 2.00 46.78 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.9 (48.55-2.00) 99.9 (46.78-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.14 (at 2.00Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.187 , 0.227 0.193 , 0.229	Depositor DCC
R_{free} test set	6017 reflections (5.28%)	DCC
Wilson B-factor (Å ²)	35.3	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 48.7	EDS
Estimated twinning fraction	0.048 for l,-k,h	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	1 of 120017 reflections (0.001%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	10566	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 17.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: FE, YCM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.97	0/1268	0.88	0/1725
1	B	0.95	1/1280 (0.1%)	0.91	1/1741 (0.1%)
1	C	0.96	2/1264 (0.2%)	0.98	2/1720 (0.1%)
1	D	0.97	0/1262	0.93	3/1717 (0.2%)
1	E	0.94	0/1268	0.88	1/1725 (0.1%)
1	F	0.92	0/1262	0.92	3/1717 (0.2%)
1	G	0.99	1/1267 (0.1%)	0.91	4/1724 (0.2%)
1	H	0.95	0/1256	0.88	2/1709 (0.1%)
All	All	0.96	4/10127 (0.0%)	0.91	16/13778 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	6	ASP	CB-CG	-5.70	1.39	1.51
1	C	67	GLU	CG-CD	5.50	1.60	1.51
1	B	67	GLU	CG-CD	5.03	1.59	1.51
1	G	97	ASN	CB-CG	5.03	1.62	1.51

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	113	ASP	CB-CG-OD2	-7.23	111.79	118.30
1	G	152	ARG	NE-CZ-NH1	6.89	123.75	120.30
1	C	113	ASP	CB-CG-OD2	-6.63	112.34	118.30
1	G	146	ARG	NE-CZ-NH2	-6.18	117.21	120.30
1	H	99	ARG	NE-CZ-NH1	6.06	123.33	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1263	0	1197	7	0
1	B	1270	0	1202	3	0
1	C	1255	0	1190	10	0
1	D	1249	0	1185	5	0
1	E	1255	0	1193	11	0
1	F	1243	0	1182	12	0
1	G	1247	0	1191	16	0
1	H	1237	0	1178	10	0
2	A	4	0	0	0	0
2	B	5	0	0	0	0
2	C	4	0	0	0	0
2	D	5	0	0	0	0
2	E	5	0	0	0	0
2	F	5	0	0	0	0
2	G	5	0	0	0	0
2	H	4	0	0	0	0
3	A	55	0	0	1	0
3	B	59	0	0	0	0
3	C	71	0	0	1	0
3	D	63	0	0	1	0
3	E	61	0	0	1	0
3	F	57	0	0	1	0
3	G	74	0	0	0	0
3	H	70	0	0	1	0
All	All	10566	0	9518	69	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 69 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:6:ASP:HB3	1:G:10:ARG:NH1	1.92	0.83
1:A:157:VAL:HG11	1:H:157:VAL:HG22	1.70	0.74
1:G:94:GLU:HG3	1:G:134:ILE:CD1	2.20	0.70

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:130:ALA:O	1:C:134:ILE:HD12	1.92	0.69
1:E:157:VAL:HG22	1:F:157:VAL:HG11	1.78	0.66

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	156/168 (93%)	153 (98%)	2 (1%)	1 (1%)	30	22
1	B	157/168 (94%)	155 (99%)	2 (1%)	0	100	100
1	C	155/168 (92%)	151 (97%)	4 (3%)	0	100	100
1	D	155/168 (92%)	153 (99%)	1 (1%)	1 (1%)	30	22
1	E	156/168 (93%)	152 (97%)	3 (2%)	1 (1%)	30	22
1	F	155/168 (92%)	152 (98%)	2 (1%)	1 (1%)	30	22
1	G	155/168 (92%)	152 (98%)	1 (1%)	2 (1%)	15	7
1	H	154/168 (92%)	151 (98%)	2 (1%)	1 (1%)	30	22
All	All	1243/1344 (92%)	1219 (98%)	17 (1%)	7 (1%)	30	22

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	157	VAL
1	D	157	VAL
1	F	157	VAL
1	H	157	VAL
1	E	157	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	135/142 (95%)	135 (100%)	0	100	100
1	B	136/142 (96%)	134 (98%)	2 (2%)	72	75
1	C	134/142 (94%)	129 (96%)	5 (4%)	41	38
1	D	134/142 (94%)	134 (100%)	0	100	100
1	E	135/142 (95%)	131 (97%)	4 (3%)	48	47
1	F	134/142 (94%)	130 (97%)	4 (3%)	48	47
1	G	134/142 (94%)	132 (98%)	2 (2%)	72	75
1	H	133/142 (94%)	129 (97%)	4 (3%)	48	47
All	All	1075/1136 (95%)	1054 (98%)	21 (2%)	63	65

5 of 21 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	30	ASP
1	F	30	ASP
1	H	2	GLU
1	E	3	GLU
1	H	30	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	F	97	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

13 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	YCM	A	77[A]	1	7,9,10	2.16	2 (28%)	5,10,12	2.17	2 (40%)
1	YCM	A	77[B]	1	7,9,10	0.79	0	5,10,12	1.05	0
1	YCM	B	77[A]	1	7,9,10	1.86	1 (14%)	5,10,12	1.22	1 (20%)
1	YCM	B	77[B]	1	3,5,10	1.14	0	3,5,12	1.15	0
1	YCM	C	77[A]	1	7,9,10	1.58	1 (14%)	5,10,12	1.07	1 (20%)
1	YCM	C	77[B]	1	3,5,10	0.50	0	3,5,12	1.23	0
1	YCM	D	77[A]	1	3,5,10	2.32	1 (33%)	3,5,12	2.11	2 (66%)
1	YCM	D	77[B]	1	3,5,10	1.78	1 (33%)	3,5,12	1.33	0
1	YCM	E	77[A]	1	3,5,10	2.05	1 (33%)	3,5,12	2.20	1 (33%)
1	YCM	E	77[B]	1	3,5,10	0.66	0	3,5,12	1.50	1 (33%)
1	YCM	F	77	1	3,5,10	1.30	0	3,5,12	1.37	1 (33%)
1	YCM	G	77	1	3,5,10	1.21	0	3,5,12	1.32	0
1	YCM	H	77	1	3,5,10	1.70	1 (33%)	3,5,12	1.79	1 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	YCM	A	77[A]	1	-	0/6/8/10	0/0/0/0
1	YCM	A	77[B]	1	-	0/6/8/10	0/0/0/0
1	YCM	B	77[A]	1	-	0/6/8/10	0/0/0/0
1	YCM	B	77[B]	1	-	0/1/4/10	0/0/0/0
1	YCM	C	77[A]	1	-	0/6/8/10	0/0/0/0
1	YCM	C	77[B]	1	-	0/1/4/10	0/0/0/0
1	YCM	D	77[A]	1	-	0/1/4/10	0/0/0/0
1	YCM	D	77[B]	1	-	0/1/4/10	0/0/0/0
1	YCM	E	77[A]	1	-	0/1/4/10	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	YCM	E	77[B]	1	-	0/1/4/10	0/0/0/0
1	YCM	F	77	1	-	0/1/4/10	0/0/0/0
1	YCM	G	77	1	-	0/1/4/10	0/0/0/0
1	YCM	H	77	1	-	0/1/4/10	0/0/0/0

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	77[B]	YCM	CB-SG	2.26	1.86	1.81
1	H	77	YCM	CB-CA	2.53	1.55	1.52
1	E	77[A]	YCM	CB-CA	2.89	1.56	1.52
1	D	77[A]	YCM	CB-SG	3.34	1.88	1.81
1	C	77[A]	YCM	CB-SG	3.61	1.88	1.81

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	77[A]	YCM	O-C-CA	-2.48	119.08	125.72
1	E	77[B]	YCM	O-C-CA	-2.31	119.53	125.72
1	H	77	YCM	O-C-CA	-2.25	119.68	125.72
1	D	77[A]	YCM	O-C-CA	-2.14	119.98	125.72
1	C	77[A]	YCM	O-C-CA	-2.05	120.21	125.72

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

13 monomers are involved in 18 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	77[A]	YCM	1	0
1	A	77[B]	YCM	1	0
1	B	77[A]	YCM	2	0
1	B	77[B]	YCM	1	0
1	C	77[A]	YCM	2	0
1	C	77[B]	YCM	1	0
1	D	77[A]	YCM	2	0
1	D	77[B]	YCM	1	0
1	E	77[A]	YCM	1	0
1	E	77[B]	YCM	1	0
1	F	77	YCM	1	0
1	G	77	YCM	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	H	77	YCM	2	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 37 ligands modelled in this entry, 37 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	156/168 (92%)	-0.03	4 (2%) 59 60	27, 38, 62, 81	0
1	B	157/168 (93%)	-0.00	6 (3%) 44 45	27, 39, 60, 101	0
1	C	156/168 (92%)	-0.02	7 (4%) 37 38	27, 38, 60, 98	0
1	D	156/168 (92%)	-0.03	3 (1%) 70 70	26, 37, 59, 86	0
1	E	157/168 (93%)	-0.02	8 (5%) 32 33	29, 38, 69, 86	0
1	F	156/168 (92%)	-0.01	6 (3%) 44 45	28, 39, 62, 86	0
1	G	156/168 (92%)	-0.06	5 (3%) 51 52	26, 36, 61, 90	0
1	H	156/168 (92%)	0.03	4 (2%) 59 60	27, 37, 65, 82	0
All	All	1250/1344 (93%)	-0.02	43 (3%) 49 50	26, 38, 64, 101	0

The worst 5 of 43 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	157	VAL	7.8
1	H	156	VAL	6.4
1	F	154	LEU	5.7
1	H	157	VAL	5.3
1	E	153	SER	4.8

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
1	YCM	E	77[B]	6/11	0.92	0.12	-	40,45,47,47	6
1	YCM	G	77	6/11	0.88	0.12	-	41,47,53,66	0
1	YCM	E	77[A]	6/11	0.92	0.12	-	45,49,51,51	6
1	YCM	A	77[B]	10/11	0.85	0.23	-	42,45,52,53	10
1	YCM	A	77[A]	10/11	0.85	0.23	-	34,44,49,53	10
1	YCM	C	77[A]	10/11	0.93	0.15	-	29,39,43,46	10
1	YCM	B	77[B]	6/11	0.91	0.18	-	40,41,42,44	6
1	YCM	B	77[A]	10/11	0.91	0.18	-	38,41,42,44	10
1	YCM	D	77[B]	6/11	0.92	0.12	-	36,42,44,45	6
1	YCM	F	77	6/11	0.94	0.08	-	46,56,60,66	0
1	YCM	C	77[B]	6/11	0.93	0.15	-	39,40,43,45	6
1	YCM	H	77	6/11	0.95	0.09	-	40,51,53,56	0
1	YCM	D	77[A]	6/11	0.92	0.12	-	46,46,49,50	6

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	FE	A	202	1/1	0.93	0.24	9.10	58,58,58,58	1
2	FE	G	203	1/1	0.61	0.24	6.20	51,51,51,51	1
2	FE	E	204	1/1	0.93	0.25	6.05	54,54,54,54	1
2	FE	G	202	1/1	0.87	0.21	4.12	65,65,65,65	1
2	FE	C	203	1/1	0.85	0.18	1.49	55,55,55,55	1
2	FE	C	202	1/1	0.97	0.18	1.33	54,54,54,54	1
2	FE	F	205	1/1	0.98	0.13	1.23	42,42,42,42	1
2	FE	B	202	1/1	0.96	0.12	0.52	66,66,66,66	1
2	FE	B	204	1/1	0.94	0.12	0.28	48,48,48,48	1
2	FE	E	202	1/1	0.97	0.11	0.18	69,69,69,69	1
2	FE	F	203	1/1	0.98	0.11	-0.42	65,65,65,65	1
2	FE	D	203	1/1	0.94	0.11	-0.81	54,54,54,54	1
2	FE	C	204	1/1	0.99	0.08	-1.43	66,66,66,66	1
2	FE	D	202	1/1	0.93	0.16	-	63,63,63,63	1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	FE	H	202	1/1	0.91	0.16	-	62,62,62,62	1
2	FE	F	204	1/1	0.85	0.07	-	64,64,64,64	1
2	FE	E	205	1/1	0.87	0.11	-	58,58,58,58	1
2	FE	B	205	1/1	0.97	0.08	-	53,53,53,53	1
2	FE	E	201	1/1	0.97	0.17	-	63,63,63,63	1
2	FE	F	201	1/1	0.90	0.05	-	61,61,61,61	1
2	FE	F	202	1/1	0.99	0.17	-	52,52,52,52	1
2	FE	H	204	1/1	0.95	0.13	-	57,57,57,57	1
2	FE	B	203	1/1	0.89	0.16	-	60,60,60,60	1
2	FE	E	203	1/1	0.99	0.17	-	52,52,52,52	1
2	FE	D	204	1/1	0.90	0.07	-	62,62,62,62	1
2	FE	H	203	1/1	0.85	0.10	-	56,56,56,56	1
2	FE	A	201	1/1	0.96	0.15	-	49,49,49,49	1
2	FE	G	205	1/1	0.93	0.16	-	60,60,60,60	1
2	FE	D	205	1/1	0.89	0.19	-	55,55,55,55	1
2	FE	G	201	1/1	0.98	0.24	-	45,45,45,45	1
2	FE	D	201	1/1	0.98	0.14	-	53,53,53,53	1
2	FE	A	204	1/1	0.79	0.19	-	65,65,65,65	1
2	FE	H	201	1/1	0.97	0.22	-	49,49,49,49	1
2	FE	A	203	1/1	0.89	0.11	-	60,60,60,60	1
2	FE	G	204	1/1	0.80	0.14	-	57,57,57,57	1
2	FE	B	201	1/1	0.97	0.14	-	54,54,54,54	1
2	FE	C	201	1/1	0.89	0.16	-	65,65,65,65	1

6.5 Other polymers

There are no such residues in this entry.