



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 12:07 AM GMT

PDB ID : 1ZM2
Title : Structure of ADP-ribosylated eEF2 in complex with catalytic fragment of ETA
Authors : Joergensen, R.; Merrill, A.R.; Yates, S.P.; Marquez, V.E.; Schwan, A.L.; Boesen, T.; Andersen, G.R.
Deposited on : 2005-05-10
Resolution : 3.07 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.07 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1119 (3.12-3.04)
Clashscore	102246	1098 (3.10-3.06)
Ramachandran outliers	100387	1057 (3.10-3.06)
Sidechain outliers	100360	1057 (3.10-3.06)
RSRZ outliers	91569	1001 (3.10-3.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	842	<div> <div>53%</div> <div>42%</div> <div>..</div> </div>
1	C	842	<div> <div>3%</div> <div>47%</div> <div>47%</div> <div>..</div> </div>
1	E	842	<div> <div>14%</div> <div>46%</div> <div>48%</div> <div>..</div> </div>
2	B	207	<div> <div>62%</div> <div>32%</div> <div>5%</div> </div>
2	D	207	<div> <div>59%</div> <div>35%</div> <div>5%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	207	 53% 41% 6%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
1	DDE	E	699	-	-	X	-
3	APR	E	843	X	-	-	X

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 24024 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Elongation factor 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	823	Total	C	N	O	S	0	0	0
			6405	4075	1093	1207	30			
1	C	823	Total	C	N	O	S	0	0	0
			6405	4075	1093	1207	30			
1	E	823	Total	C	N	O	S	0	0	0
			6415	4082	1095	1208	30			


There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	699	DDE	HIS	MODIFIED RESIDUE	UNP P32324
C	699	DDE	HIS	MODIFIED RESIDUE	UNP P32324
E	699	DDE	HIS	MODIFIED RESIDUE	UNP P32324

- Molecule 2 is a protein called Exotoxin A.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	207	Total	C	N	O	0	0	0
			1588	1001	283	304			
2	D	207	Total	C	N	O	0	0	0
			1588	1001	283	304			
2	F	207	Total	C	N	O	0	0	0
			1588	1001	283	304			

- Molecule 3 is ADENOSINE-5-DIPHOSPHORIBOSE (three-letter code: APR) (formula: C₁₅H₂₃N₅O₁₄P₂).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	E	1	Total	C	N	O	P	0	0
			35	15	5	13	2		

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Elongation factor 2





• Molecule 1: Elongation factor 2






• Molecule 1: Elongation factor 2







• Molecule 2: Exotoxin A




• Molecule 2: Exotoxin A





● Molecule 2: Exotoxin A



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	330.63Å 67.84Å 191.50Å 90.00° 103.30° 90.00°	Depositor
Resolution (Å)	30.00 – 3.07 29.63 – 3.07	Depositor EDS
% Data completeness (in resolution range)	(Not available) (30.00-3.07) 99.4 (29.63-3.07)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.14	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.65 (at 3.06Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.231 , 0.265 0.229 , 0.256	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	52.7	Xtriage
Anisotropy	0.304	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 53.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.27$	Xtriage
Outliers	4 of 77946 reflections (0.005%)	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	24024	wwPDB-VP
Average B, all atoms (Å ²)	61.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 39.80 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 3.0079e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: APR, DDE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.47	0/6517	0.68	0/8823
1	C	0.43	0/6517	0.67	0/8823
1	E	0.41	0/6517	0.64	0/8823
2	B	0.60	0/1627	0.86	0/2216
2	D	0.58	0/1627	0.81	0/2216
2	F	0.57	0/1627	0.84	1/2216 (0.0%)
All	All	0.47	0/24432	0.70	1/33117 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	494	ARG	NE-CZ-NH1	6.14	123.37	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6405	0	6472	342	0
1	C	6405	0	6472	377	0
1	E	6415	0	6487	422	0
2	B	1588	0	1539	66	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	1588	0	1539	74	0
2	F	1588	0	1539	82	0
3	E	35	0	19	6	0
All	All	24024	0	24067	1350	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 28.

The worst 5 of 1350 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:546:GLU:HG3	2:D:547:GLU:HG3	1.29	1.14
2:B:546:GLU:HG3	2:B:547:GLU:HG3	1.32	1.11
1:C:404:THR:HG22	1:C:449:PRO:HA	1.34	1.05
1:E:68:ILE:HD12	1:E:390:ASP:HB2	1.37	1.03
1:C:231:LYS:HE3	1:C:232:LYS:HG3	1.42	1.02

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	818/842 (97%)	732 (90%)	74 (9%)	12 (2%)	13	47
1	C	818/842 (97%)	711 (87%)	87 (11%)	20 (2%)	7	33
1	E	818/842 (97%)	707 (86%)	97 (12%)	14 (2%)	11	43
2	B	205/207 (99%)	179 (87%)	24 (12%)	2 (1%)	19	58
2	D	205/207 (99%)	176 (86%)	24 (12%)	5 (2%)	7	33
2	F	205/207 (99%)	175 (85%)	27 (13%)	3 (2%)	13	47
All	All	3069/3147 (98%)	2680 (87%)	333 (11%)	56 (2%)	11	42

5 of 56 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	112	SER
1	C	235	VAL
1	C	460	ASP
1	C	639	ASP
1	C	761	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	699/714 (98%)	669 (96%)	30 (4%)	35	73
1	C	699/714 (98%)	651 (93%)	48 (7%)	19	55
1	E	699/714 (98%)	664 (95%)	35 (5%)	30	67
2	B	161/162 (99%)	145 (90%)	16 (10%)	10	35
2	D	161/162 (99%)	144 (89%)	17 (11%)	8	32
2	F	161/162 (99%)	146 (91%)	15 (9%)	11	39
All	All	2580/2628 (98%)	2419 (94%)	161 (6%)	23	59

5 of 161 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	528	HIS
1	C	813	SER
2	F	437	VAL
1	C	544	ASP
1	C	677	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 75 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	274	ASN
1	C	581	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	E	753	GLN
1	C	414	GLN
1	C	537	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	DDE	A	699	1	5,10,21	0.76	0	3,12,30	1.60	1 (33%)
1	DDE	C	699	1	5,10,21	1.20	1 (20%)	3,12,30	1.41	1 (33%)
1	DDE	E	699	1,3	13,20,21	1.97	4 (30%)	16,28,30	3.66	4 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	DDE	A	699	1	-	0/4/6/23	0/1/1/1
1	DDE	C	699	1	-	0/4/6/23	0/1/1/1
1	DDE	E	699	1,3	-	0/19/21/23	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	699	DDE	CAA-NCB	2.05	1.56	1.50
1	E	699	DDE	CBW-CBI	2.10	1.56	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	699	DDE	CB-CA	2.21	1.58	1.53
1	E	699	DDE	CAC-NCB	3.26	1.59	1.50
1	E	699	DDE	OAG-CBI	5.06	1.33	1.23

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	699	DDE	OAG-CBI-CBW	-6.76	112.63	120.72
1	C	699	DDE	CD2-NE2-CE1	2.04	108.93	105.71
1	A	699	DDE	CD2-NE2-CE1	2.07	108.97	105.71
1	E	699	DDE	OAG-CBI-NAD	5.50	131.24	123.08
1	E	699	DDE	CAU-CAT-CE1	7.07	150.71	112.58

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	699	DDE	1	0
1	C	699	DDE	1	0
1	E	699	DDE	11	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	APR	E	843	1	31,38,39	1.44	6 (19%)	39,58,60	1.82	9 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	APR	E	843	1	1/1/9/10	0/18/51/54	0/4/4/4

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	843	APR	O4D-C1D	2.01	1.48	1.43
3	E	843	APR	O4D-C4D	2.06	1.48	1.44
3	E	843	APR	C4-N3	2.17	1.38	1.35
3	E	843	APR	O2'-C2'	2.69	1.49	1.43
3	E	843	APR	C3'-C4'	2.76	1.60	1.53

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	843	APR	PB-O3A-PA	-4.98	118.74	132.73
3	E	843	APR	O2D-C2D-C3D	-4.87	101.89	111.23
3	E	843	APR	O4D-C4D-C5D	-2.97	103.04	109.53
3	E	843	APR	O4D-C1D-C2D	-2.68	100.68	106.16
3	E	843	APR	O2A-PA-O3A	2.11	114.67	105.09

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	E	843	APR	C1D

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	E	843	APR	6	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	822/842 (97%)	-0.43	0 100 100	5, 48, 81, 101	0
1	C	822/842 (97%)	-0.11	27 (3%) 50 26	11, 64, 123, 135	0
1	E	822/842 (97%)	0.57	117 (14%) 4 1	7, 106, 129, 147	0
2	B	207/207 (100%)	-0.71	0 100 100	5, 22, 55, 71	0
2	D	207/207 (100%)	-0.65	0 100 100	5, 25, 64, 78	0
2	F	207/207 (100%)	-0.63	0 100 100	7, 27, 64, 79	0
All	All	3087/3147 (98%)	-0.12	144 (4%) 35 16	5, 54, 123, 147	0

The worst 5 of 144 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	269	LEU	5.6
1	E	197	LEU	5.5
1	E	167	LEU	5.4
1	C	499	ASN	5.2
1	E	166	GLU	5.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	DDE	C	699	10/21	0.84	0.21	-	44,52,55,56	0
1	DDE	E	699	20/21	0.92	0.29	-	48,68,80,81	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
1	DDE	A	699	10/21	0.89	0.13	-	37,41,46,47	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	APR	E	843	35/36	0.85	0.33	6.43	64,71,85,86	0

6.5 Other polymers [i](#)

There are no such residues in this entry.