



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:29 PM GMT

PDB ID : 3ZNQ  
Title : IN VITRO AND IN VIVO INHIBITION OF HUMAN D-AMINO ACID OXIDASE: REGULATION OF D-SERINE CONCENTRATION IN THE BRAIN  
Authors : Hopkins, S.C.; Heffernan, M.L.R.; Saraswat, L.D.; Bowen, C.A.; Melnick, L.; Hardy, L.W.; Orsini, M.A.; Allen, M.S.; Koch, P.; Spear, K.L.; Foglesong, R.J.; Soukri, M.; Chytil, M.; Fang, Q.K.; Jones, S.W.; Varney, M.A.; Panatier, A.; Olier, S.H.R.; Pollegioni, L.; Piubelli, L.; Molla, G.; Nardini, M.; Large, T.H.  
Deposited on : 2013-02-15  
Resolution : 2.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3340 (2.80-2.72)
Clashscore	102246	3829 (2.80-2.72)
Ramachandran outliers	100387	3767 (2.80-2.72)
Sidechain outliers	100360	3770 (2.80-2.72)
RSRZ outliers	91569	3352 (2.80-2.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	347	 5% 79% 15% . . .
1	B	347	 12% 78% 17% . .

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 5623 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called D-AMINO-ACID OXIDASE.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	338	Total	C	N	O	S	0	1	0
			2727	1747	480	491	9			
1	B	338	Total	C	N	O	S	0	0	0
			2719	1742	477	491	9			

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula:  $C_{27}H_{33}N_9O_{15}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is 3-PHENETHYL-4H-FURO[3,2-B]PYRROLE-5-CARBOXYLIC ACID (three-letter code: SS8) (formula:  $C_{15}H_{11}NO_3$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			19	15	1	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	29	Total	O	0	0
			29	29		
4	B	23	Total	O	0	0
			23	23		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: D-AMINO-ACID OXIDASE



#### • Molecule 1: D-AMINO-ACID OXIDASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	84.47Å 84.47Å 188.67Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	27.65 – 2.75 27.65 – 2.75	Depositor EDS
% Data completeness (in resolution range)	90.1 (27.65-2.75) 90.4 (27.65-2.75)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.02 (at 2.76Å)	Xtriage
Refinement program	BUSTER 2.11.1	Depositor
R, $R_{free}$	0.199 , 0.255 0.216 , 0.268	Depositor DCC
$R_{free}$ test set	966 reflections (5.38%)	DCC
Wilson B-factor (Å <sup>2</sup> )	59.0	Xtriage
Anisotropy	0.448	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 67.0	EDS
Estimated twinning fraction	0.032 for -h,-k,l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 18907 reflections	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	5623	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	91.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.51% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SS8, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.49	1/2807 (0.0%)	0.73	2/3819 (0.1%)
1	B	0.43	0/2796	0.71	3/3805 (0.1%)
All	All	0.46	1/5603 (0.0%)	0.72	5/7624 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	126	PRO	N-CD	11.13	1.63	1.47

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	56	LEU	CB-CA-C	-14.78	82.11	110.20
1	B	55	TYR	CB-CA-C	9.40	129.20	110.40
1	B	56	LEU	CB-CA-C	-8.99	93.12	110.20
1	B	60	ASN	CB-CA-C	-7.88	94.65	110.40
1	A	56	LEU	N-CA-C	7.42	131.03	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	172	ARG	Sidechain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2727	0	2677	66	0
1	B	2719	0	2664	28	0
2	A	53	0	31	1	0
2	B	53	0	31	0	0
3	A	19	0	10	5	0
4	A	29	0	0	1	0
4	B	23	0	0	0	0
All	All	5623	0	5413	93	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (93) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:220:GLU:CG	1:A:221[B]:ARG:HH21	1.23	1.49
1:A:220:GLU:HG2	1:A:221[B]:ARG:NH2	1.37	1.37
1:A:220:GLU:CG	1:A:221[B]:ARG:NH2	1.87	1.31
1:A:220:GLU:CD	1:A:221[B]:ARG:NH2	1.83	1.30
1:A:22:ARG:HA	1:A:22:ARG:NE	1.63	1.05
1:A:220:GLU:CD	1:A:221[B]:ARG:HH22	1.49	1.03
1:A:22:ARG:HE	1:A:22:ARG:CA	1.73	1.01
1:A:22:ARG:HA	1:A:22:ARG:HE	0.85	1.00
1:A:220:GLU:HG2	1:A:221[B]:ARG:HH21	0.77	0.93
1:A:221[B]:ARG:CA	1:A:221[B]:ARG:HE	1.81	0.91
1:A:161:GLN:HG2	1:B:249:GLU:H	1.37	0.89
1:A:155:ARG:HG3	1:A:155:ARG:HH11	1.36	0.88
1:A:221[B]:ARG:NE	1:A:221[B]:ARG:HA	1.87	0.85
1:A:22:ARG:O	1:A:22:ARG:HD3	1.75	0.85
1:A:155:ARG:CG	1:A:155:ARG:HH11	1.89	0.83
1:A:61:ASN:ND2	1:A:63:GLN:HG3	1.98	0.78

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:220:GLU:OE1	1:A:221[B]:ARG:NH2	2.18	0.77
1:A:283:ARG:NE	3:A:357:SS8:O2	2.14	0.76
1:A:61:ASN:CG	1:A:63:GLN:HG3	2.06	0.76
1:A:21:GLU:HA	1:A:21:GLU:OE1	1.86	0.76
1:A:235:THR:HG21	1:A:276:ILE:HG12	1.70	0.74
1:A:53:GLN:HE22	1:A:96:ASN:HD21	1.34	0.74
1:A:221[B]:ARG:HA	1:A:221[B]:ARG:HE	1.48	0.72
3:A:357:SS8:H15	3:A:357:SS8:C6	2.19	0.71
1:A:20:HIS:NE2	1:A:155:ARG:HB3	2.05	0.70
1:A:21:GLU:CA	1:A:21:GLU:OE1	2.39	0.70
1:A:283:ARG:HE	3:A:357:SS8:C1	2.05	0.69
1:A:221[B]:ARG:NE	1:A:221[B]:ARG:CA	2.40	0.67
1:B:61:ASN:HD21	1:B:63:GLN:HG3	1.62	0.65
1:A:20:HIS:O	1:A:22:ARG:N	2.30	0.63
1:A:20:HIS:NE2	1:A:155:ARG:HD3	2.14	0.62
1:B:55:TYR:O	1:B:56:LEU:HB2	2.00	0.61
1:A:22:ARG:NE	1:A:22:ARG:CA	2.39	0.61
1:A:20:HIS:CE1	1:A:24:HIS:CG	2.89	0.61
1:A:3:VAL:HG11	1:A:330:PHE:HE1	1.65	0.60
1:A:155:ARG:NH1	1:A:155:ARG:CG	2.55	0.60
1:A:220:GLU:CB	1:A:221[B]:ARG:HH21	2.09	0.59
1:A:22:ARG:O	1:A:22:ARG:CD	2.49	0.59
1:A:20:HIS:HE1	1:A:24:HIS:ND1	2.02	0.58
1:A:61:ASN:OD1	1:A:63:GLN:HG3	2.04	0.57
1:B:79:VAL:HG13	1:B:80:HIS:HD2	1.69	0.57
1:A:104:ASP:HB3	1:A:108:LYS:HE3	1.86	0.57
1:A:224:TYR:HB2	1:A:242:PHE:HB2	1.86	0.56
1:A:104:ASP:HB3	1:A:108:LYS:CE	2.36	0.55
1:B:61:ASN:HD22	1:B:64:GLU:HG3	1.72	0.54
1:B:206:ASP:HB2	1:B:276:ILE:HD13	1.89	0.53
1:B:41:PRO:HB3	1:B:142:LYS:HD2	1.91	0.53
1:A:15:THR:HG21	1:A:179:VAL:HG11	1.92	0.52
1:B:61:ASN:ND2	1:B:63:GLN:HG3	2.24	0.52
1:B:92:ILE:HG12	1:B:213:PHE:HD2	1.74	0.51
1:A:20:HIS:C	1:A:22:ARG:H	2.13	0.51
1:A:53:GLN:HE22	1:A:96:ASN:ND2	2.07	0.50
1:B:291:LEU:HD21	1:B:325:GLU:HB3	1.93	0.50
1:B:15:THR:HG21	1:B:179:VAL:HG11	1.94	0.49
1:A:3:VAL:HG11	1:A:330:PHE:CE1	2.46	0.49
1:A:5:VAL:HB	1:A:34:VAL:HG22	1.94	0.49
1:B:227:PRO:HB2	1:B:266:LEU:HD11	1.95	0.48

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:161:GLN:HG3	1:B:161:GLN:O	2.13	0.48
1:A:249:GLU:H	1:B:161:GLN:HG2	1.79	0.48
1:A:42:LEU:HD22	1:B:42:LEU:HD23	1.96	0.48
1:B:3:VAL:HB	1:B:32:ILE:HG22	1.96	0.47
1:A:20:HIS:CD2	1:A:155:ARG:HD3	2.50	0.47
1:A:38:ARG:HE	2:A:356:FAD:H2B	1.80	0.47
3:A:357:SS8:H15	3:A:357:SS8:N5	2.29	0.47
1:B:27:LEU:HD22	1:B:30:LEU:HB2	1.96	0.47
1:A:56:LEU:HD13	3:A:357:SS8:H18	1.96	0.46
1:A:55:TYR:CD1	1:A:55:TYR:O	2.69	0.46
1:A:96:ASN:ND2	1:A:217:HIS:NE2	2.60	0.46
1:A:55:TYR:O	1:A:56:LEU:HB2	2.16	0.46
1:B:92:ILE:HG21	1:B:138:ILE:HG13	1.97	0.46
1:A:20:HIS:HE1	1:A:24:HIS:CE1	2.34	0.46
1:A:20:HIS:HE1	1:A:24:HIS:CG	2.34	0.46
1:A:266:LEU:O	1:A:268:PRO:HD3	2.17	0.45
1:A:296:LEU:HB3	1:A:303:THR:HG23	2.00	0.44
1:A:11:ILE:HG22	1:A:181:CYS:SG	2.58	0.43
1:A:197:PRO:HG3	1:A:247:TRP:CE2	2.53	0.43
1:B:284:PRO:HB2	1:B:310:GLY:HA2	2.00	0.43
1:B:13:LEU:HB3	1:B:148:LEU:HD13	1.99	0.43
1:B:50:GLY:HA2	1:B:316:LEU:HD13	1.99	0.43
1:A:61:ASN:HD21	1:A:63:GLN:HG3	1.81	0.43
1:B:20:HIS:CD2	1:B:155:ARG:HD3	2.54	0.43
1:A:55:TYR:CG	1:A:55:TYR:O	2.73	0.42
1:B:332:ARG:HG2	1:B:332:ARG:H	1.65	0.42
1:A:61:ASN:OD1	1:A:63:GLN:CG	2.66	0.42
1:A:204:LYS:HD3	1:A:276:ILE:HD11	2.02	0.42
1:B:63:GLN:HA	1:B:66:ASP:HB2	2.02	0.42
1:A:221[B]:ARG:HD3	4:A:2020:HOH:O	2.20	0.41
1:B:115:ARG:HG3	1:B:116:LYS:N	2.36	0.41
1:B:112:LEU:HB2	1:B:135:THR:HB	2.01	0.41
1:B:13:LEU:HD13	1:B:152:LEU:HD22	2.03	0.40
1:B:190:GLN:HE22	1:B:290:ARG:HH22	1.69	0.40
1:A:151:ARG:O	1:A:155:ARG:HG3	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	337/347 (97%)	307 (91%)	27 (8%)	3 (1%)	21	52
1	B	336/347 (97%)	311 (93%)	21 (6%)	4 (1%)	16	43
All	All	673/694 (97%)	618 (92%)	48 (7%)	7 (1%)	19	48

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	21	GLU
1	B	55	TYR
1	A	319	HIS
1	B	233	THR
1	A	20	HIS
1	B	240	GLY
1	B	319	HIS

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	291/299 (97%)	259 (89%)	32 (11%)	8	21
1	B	290/299 (97%)	261 (90%)	29 (10%)	9	24
All	All	581/598 (97%)	520 (90%)	61 (10%)	8	22

All (61) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	ARG
1	A	18	CYS
1	A	20	HIS
1	A	21	GLU
1	A	22	ARG
1	A	27	LEU
1	A	30	LEU
1	A	32	ILE
1	A	38	ARG
1	A	55	TYR
1	A	58	ASP
1	A	60	ASN
1	A	63	GLN
1	A	87	LEU
1	A	89	LEU
1	A	91	LEU
1	A	102	ILE
1	A	106	SER
1	A	108	LYS
1	A	116	LYS
1	A	120	ARG
1	A	142	LYS
1	A	145	LEU
1	A	151	ARG
1	A	154	GLU
1	A	155	ARG
1	A	173	GLU
1	A	206	ASP
1	A	235	THR
1	A	244	LEU
1	A	286	ARG
1	A	296	LEU
1	B	2	ARG
1	B	27	LEU
1	B	28	GLN
1	B	30	LEU
1	B	89	LEU
1	B	91	LEU
1	B	96	ASN
1	B	115	ARG
1	B	125	PHE
1	B	142	LYS
1	B	145	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	161	GLN
1	B	164	VAL
1	B	165	GLU
1	B	191	ARG
1	B	199	ARG
1	B	205	VAL
1	B	221	ARG
1	B	238	LEU
1	B	250	LEU
1	B	252	ASN
1	B	255	ASP
1	B	274	ARG
1	B	276	ILE
1	B	288	GLN
1	B	290	ARG
1	B	301	SER
1	B	332	ARG
1	B	333	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	20	HIS
1	A	69	GLN
1	A	80	HIS
1	A	96	ASN
1	A	180	ASN
1	A	252	ASN
1	A	295	GLN
1	B	20	HIS
1	B	53	GLN
1	B	61	ASN
1	B	80	HIS
1	B	83	ASN
1	B	96	ASN
1	B	134	HIS
1	B	180	ASN
1	B	246	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	FAD	A	356	-	48,58,58	1.96	15 (31%)	54,89,89	3.14	21 (38%)
3	SS8	A	357	-	14,21,21	2.72	7 (50%)	14,29,29	2.35	6 (42%)
2	FAD	B	356	-	48,58,58	2.07	13 (27%)	54,89,89	2.68	19 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	356	-	-	0/30/50/50	0/6/6/6
3	SS8	A	357	-	-	0/5/9/9	0/2/3/3
2	FAD	B	356	-	-	0/30/50/50	0/6/6/6

All (35) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	356	FAD	O2B-C2B	-2.21	1.37	1.43
2	A	356	FAD	P-O2P	-2.07	1.46	1.54
2	B	356	FAD	C5A-N7A	-2.03	1.32	1.39
2	A	356	FAD	C2A-N1A	2.05	1.37	1.33
3	A	357	SS8	C17-C16	2.11	1.43	1.38
2	A	356	FAD	C10-N1	2.11	1.39	1.35
2	A	356	FAD	C4X-C10	2.17	1.45	1.41
2	B	356	FAD	C9-C9A	2.20	1.45	1.40
2	B	356	FAD	C4A-N3A	2.28	1.39	1.35
2	B	356	FAD	C4-N3	2.38	1.37	1.33
2	A	356	FAD	O4'-C4'	2.39	1.48	1.43
2	A	356	FAD	C4-N3	2.41	1.37	1.33
2	B	356	FAD	C6-C7	2.64	1.45	1.37
2	B	356	FAD	C8A-N7A	2.67	1.39	1.34
2	A	356	FAD	C8A-N7A	2.77	1.39	1.34
2	A	356	FAD	C2A-N3A	2.89	1.37	1.32
2	B	356	FAD	C2A-N3A	3.01	1.37	1.32
3	A	357	SS8	C18-C17	3.07	1.45	1.38
3	A	357	SS8	C10-C11	3.08	1.40	1.37
2	B	356	FAD	O4B-C1B	3.12	1.45	1.41
2	B	356	FAD	C2A-N1A	3.14	1.39	1.33
2	A	356	FAD	C4A-N3A	3.28	1.40	1.35
2	A	356	FAD	C6-C7	3.33	1.47	1.37
2	A	356	FAD	O4B-C1B	3.34	1.45	1.41
2	B	356	FAD	C4X-C10	3.34	1.47	1.41
3	A	357	SS8	C14-C13	3.80	1.59	1.47
3	A	357	SS8	C19-C14	3.87	1.46	1.39
2	A	356	FAD	C4X-N5	3.91	1.39	1.33
2	A	356	FAD	C10-N10	4.00	1.43	1.39
3	A	357	SS8	C15-C14	4.01	1.47	1.39
2	B	356	FAD	C4X-N5	4.23	1.40	1.33
2	B	356	FAD	C10-N10	4.49	1.44	1.39
3	A	357	SS8	C13-C12	4.83	1.55	1.31
2	A	356	FAD	C9A-N10	6.73	1.48	1.38
2	B	356	FAD	C9A-N10	7.89	1.49	1.38

All (46) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	356	FAD	N3A-C2A-N1A	-13.82	118.31	128.89
2	B	356	FAD	N3A-C2A-N1A	-11.32	120.22	128.89
2	A	356	FAD	C4X-C10-N10	-6.14	116.90	120.52
2	B	356	FAD	C4X-C10-N10	-4.49	117.88	120.52

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	356	FAD	O3P-P-O5'	-4.08	92.10	102.94
2	B	356	FAD	C4X-C4-N3	-3.70	118.53	123.59
2	A	356	FAD	C4A-C5A-N7A	-3.55	106.21	109.48
2	B	356	FAD	P-O3P-PA	-3.52	122.85	132.73
2	A	356	FAD	O4B-C4B-C5B	-3.47	96.91	109.32
2	A	356	FAD	C1B-N9A-C4A	-3.30	121.96	126.94
2	A	356	FAD	C4X-C4-N3	-3.18	119.24	123.59
2	A	356	FAD	C9A-C5X-N5	-3.07	117.82	122.36
3	A	357	SS8	C11-C12-C13	-2.94	117.84	125.68
3	A	357	SS8	C17-C16-C15	-2.80	116.09	120.19
3	A	357	SS8	C6-C11-C12	-2.61	117.19	126.68
2	A	356	FAD	O2P-P-O5'	-2.57	95.51	108.46
2	B	356	FAD	C9A-C5X-N5	-2.49	118.67	122.36
2	B	356	FAD	O3P-P-O5'	-2.32	96.77	102.94
2	B	356	FAD	C4A-C5A-N7A	-2.21	107.45	109.48
2	A	356	FAD	P-O3P-PA	-2.14	126.72	132.73
2	B	356	FAD	O2P-P-O5'	-2.10	97.88	108.46
2	B	356	FAD	O2'-C2'-C3'	-2.07	103.82	109.02
2	B	356	FAD	O2'-C2'-C1'	-2.02	104.97	109.94
2	A	356	FAD	O2P-P-O1P	2.12	124.01	112.53
2	A	356	FAD	C5X-C9A-N10	2.13	119.23	117.62
2	A	356	FAD	C6-C5X-N5	2.17	121.75	118.96
2	B	356	FAD	C4-C4X-N5	2.21	121.40	118.72
2	A	356	FAD	C8M-C8-C7	2.39	125.98	120.73
2	B	356	FAD	C2A-N1A-C6A	2.41	123.07	118.77
2	B	356	FAD	O2P-P-O1P	2.60	126.59	112.53
2	B	356	FAD	O2B-C2B-C3B	2.62	120.34	111.83
2	B	356	FAD	O2P-P-O3P	2.62	116.99	105.09
3	A	357	SS8	C16-C15-C14	2.98	124.44	120.64
2	A	356	FAD	C2A-N1A-C6A	3.06	124.24	118.77
2	A	356	FAD	O2P-P-O3P	3.11	119.21	105.09
3	A	357	SS8	C10-C11-C12	3.24	137.47	125.14
2	A	356	FAD	O3'-C3'-C2'	3.39	117.29	108.75
2	B	356	FAD	O4B-C1B-N9A	3.40	115.23	108.10
2	A	356	FAD	O4B-C1B-N9A	4.16	116.80	108.10
2	A	356	FAD	C2B-C1B-N9A	4.35	120.94	114.29
2	B	356	FAD	O3'-C3'-C2'	4.37	119.75	108.75
3	A	357	SS8	C8-C7-C6	5.36	111.12	106.31
2	B	356	FAD	C4-N3-C2	5.69	120.17	115.25
2	A	356	FAD	C4-N3-C2	6.91	121.22	115.25
2	A	356	FAD	C4X-N5-C5X	7.39	125.27	116.76
2	B	356	FAD	C4X-N5-C5X	7.61	125.52	116.76

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	356	FAD	1	0
3	A	357	SS8	5	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	338/347 (97%)	0.13	17 (5%) 32 25	32, 71, 141, 173	0
1	B	338/347 (97%)	0.70	42 (12%) 5 3	67, 101, 155, 191	0
All	All	676/694 (97%)	0.42	59 (8%) 13 8	32, 88, 150, 191	0

All (59) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	300	PRO	9.0
1	A	27	LEU	8.2
1	A	26	VAL	8.1
1	B	299	GLY	7.1
1	B	25	SER	6.3
1	B	60	ASN	6.2
1	A	25	SER	5.2
1	B	57	SER	5.0
1	B	298	THR	5.0
1	B	59	PRO	4.8
1	A	338	LYS	4.8
1	B	26	VAL	4.5
1	B	194	LEU	4.4
1	B	101	ALA	4.3
1	A	57	SER	4.2
1	A	297	ARG	4.2
1	B	220	GLU	4.1
1	A	337	LYS	4.0
1	A	28	GLN	3.9
1	B	296	LEU	3.9
1	B	28	GLN	3.8
1	A	298	THR	3.8
1	B	338	LYS	3.8
1	A	58	ASP	3.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	60	ASN	3.5
1	B	193	PRO	3.4
1	B	55	TYR	3.3
1	B	181	CYS	3.3
1	B	188	ALA	3.1
1	B	30	LEU	3.0
1	B	244	LEU	3.0
1	B	100	GLU	3.0
1	B	243	GLN	2.9
1	B	253	ILE	2.8
1	A	219	PRO	2.8
1	B	58	ASP	2.8
1	B	1	MET	2.8
1	B	301	SER	2.7
1	B	337	LYS	2.7
1	A	9	GLY	2.6
1	B	85	GLU	2.6
1	B	195	LEU	2.6
1	B	175	ALA	2.5
1	B	105	PRO	2.5
1	B	99	HIS	2.5
1	A	11	ILE	2.4
1	A	336	GLU	2.4
1	B	297	ARG	2.4
1	B	295	GLN	2.4
1	B	102	ILE	2.4
1	B	56	LEU	2.3
1	A	181	CYS	2.3
1	B	82	PRO	2.2
1	B	103	PRO	2.2
1	B	22	ARG	2.2
1	B	192	ASP	2.1
1	B	24	HIS	2.1
1	A	301	SER	2.0
1	B	190	GLN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	SS8	A	357	19/19	0.93	0.20	0.08	60,62,72,72	0
2	FAD	A	356	53/53	0.97	0.17	-0.19	29,43,67,71	0
2	FAD	B	356	53/53	0.93	0.17	-0.59	61,69,81,86	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.