



Full wwPDB X-ray Structure Validation Report ⓘ

May 31, 2016 – 04:40 PM EDT

PDB ID : 4ZS5
Title : Human A20 OTU domain (WT) with alkylated C103
Authors : Langley, D.B.; Christ, D.; Grey, S.T.
Deposited on : 2015-05-13
Resolution : 3.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027674
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027674


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1124 (3.24-3.16)
Clashscore	102246	1024 (3.22-3.18)
Ramachandran outliers	100387	1004 (3.22-3.18)
Sidechain outliers	100360	1003 (3.22-3.18)
RSRZ outliers	91569	1129 (3.24-3.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	371	<div> <div>%</div> <div> <div></div> <div>74%</div> <div>6%</div> <div>•</div> <div>19%</div> </div> </div>
1	B	371	<div> <div>75%</div> <div>• •</div> <div>20%</div> </div>
1	C	371	<div> <div>70%</div> <div>6%</div> <div>23%</div> </div>
1	D	371	<div> <div>%</div> <div> <div></div> <div>75%</div> <div>5%</div> <div>•</div> <div>20%</div> </div> </div>
1	E	371	<div> <div>71%</div> <div>7%</div> <div>•</div> <div>21%</div> </div>
1	F	371	<div> <div>%</div> <div> <div></div> <div>73%</div> <div>•</div> <div>24%</div> </div> </div>

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 12290 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tumor necrosis factor alpha-induced protein 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	302	Total	C	N	O	S	0	0	0
			2195	1431	386	367	11			
1	B	295	Total	C	N	O	S	0	0	0
			2022	1302	364	346	10			
1	C	284	Total	C	N	O	S	0	0	0
			1996	1287	352	345	12			
1	D	298	Total	C	N	O	S	0	0	0
			2062	1326	370	357	9			
1	E	292	Total	C	N	O	S	0	0	0
			2077	1336	367	362	12			
1	F	281	Total	C	N	O	S	0	0	0
			1938	1243	343	341	11			

There are 30 discrepancies between the modelled and reference sequences:



Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	GLY	-	expression tag	UNP P21580
A	-3	PRO	-	expression tag	UNP P21580
A	-2	LEU	-	expression tag	UNP P21580
A	-1	GLY	-	expression tag	UNP P21580
A	0	SER	-	expression tag	UNP P21580
B	-4	GLY	-	expression tag	UNP P21580
B	-3	PRO	-	expression tag	UNP P21580
B	-2	LEU	-	expression tag	UNP P21580
B	-1	GLY	-	expression tag	UNP P21580
B	0	SER	-	expression tag	UNP P21580
C	-4	GLY	-	expression tag	UNP P21580
C	-3	PRO	-	expression tag	UNP P21580
C	-2	LEU	-	expression tag	UNP P21580
C	-1	GLY	-	expression tag	UNP P21580
C	0	SER	-	expression tag	UNP P21580
D	-4	GLY	-	expression tag	UNP P21580
D	-3	PRO	-	expression tag	UNP P21580

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-2	LEU	-	expression tag	UNP P21580
D	-1	GLY	-	expression tag	UNP P21580
D	0	SER	-	expression tag	UNP P21580
E	-4	GLY	-	expression tag	UNP P21580
E	-3	PRO	-	expression tag	UNP P21580
E	-2	LEU	-	expression tag	UNP P21580
E	-1	GLY	-	expression tag	UNP P21580
E	0	SER	-	expression tag	UNP P21580
F	-4	GLY	-	expression tag	UNP P21580
F	-3	PRO	-	expression tag	UNP P21580
F	-2	LEU	-	expression tag	UNP P21580
F	-1	GLY	-	expression tag	UNP P21580
F	0	SER	-	expression tag	UNP P21580

- Molecule 1: Tumor necrosis factor alpha-induced protein 3



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	97.75Å 80.73Å 153.14Å 90.00° 102.26° 90.00°	Depositor
Resolution (Å)	54.88 – 3.20 54.88 – 3.20	Depositor EDS
% Data completeness (in resolution range)	98.9 (54.88-3.20) 99.0 (54.88-3.20)	Depositor EDS
R_{merge}	0.20	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.38 (at 3.19Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.269 , 0.334 0.266 , 0.327	Depositor DCC
R_{free} test set	1879 reflections (5.15%)	DCC
Wilson B-factor (Å ²)	68.0	Xtriage
Anisotropy	0.671	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.24 , 72.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.87	EDS
Total number of atoms	12290	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: YCM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.56	0/2240	0.71	1/3071 (0.0%)
1	B	0.52	0/2054	0.67	1/2817 (0.0%)
1	C	0.54	0/2033	0.69	1/2787 (0.0%)
1	D	0.54	0/2102	0.71	1/2882 (0.0%)
1	E	0.54	0/2111	0.76	6/2885 (0.2%)
1	F	0.56	0/1970	0.68	1/2699 (0.0%)
All	All	0.54	0/12510	0.71	11/17141 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

There are no bond length outliers.

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	204	ARG	CG-CD-NE	10.98	134.86	111.80
1	E	141	ARG	NE-CZ-NH1	9.08	124.84	120.30
1	C	104	LEU	CA-CB-CG	8.76	135.45	115.30
1	E	104	LEU	CB-CG-CD1	8.28	125.08	111.00
1	E	141	ARG	NE-CZ-NH2	-7.79	116.41	120.30
1	E	104	LEU	CA-CB-CG	-7.00	99.20	115.30
1	D	123	ARG	NE-CZ-NH1	6.89	123.75	120.30
1	F	38	HIS	CB-CA-C	6.02	122.44	110.40
1	E	141	ARG	CD-NE-CZ	5.35	131.09	123.60
1	A	295	GLU	CG-CD-OE1	5.08	128.45	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	295	GLU	CG-CD-OE1	5.08	128.45	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	34	ASN	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2195	0	1888	13	0
1	B	2022	0	1614	8	0
1	C	1996	0	1591	13	0
1	D	2062	0	1610	7	0
1	E	2077	0	1703	14	0
1	F	1938	0	1515	3	0
All	All	12290	0	9921	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:104:LEU:HD12	1:E:104:LEU:O	1.65	0.95
1:C:241:GLN:CB	1:E:46:TYR:OH	2.19	0.89
1:B:106:HIS:O	1:B:110:GLN:HB2	1.99	0.61
1:E:104:LEU:CD2	1:E:196:ILE:HG12	2.33	0.58
1:D:94:ALA:HB1	1:D:258:VAL:HG11	1.87	0.57
1:C:94:ALA:HB1	1:C:258:VAL:HG11	1.86	0.56
1:A:60:PHE:CD1	1:A:63:ILE:HD12	2.40	0.56
1:E:104:LEU:HD12	1:E:104:LEU:C	2.25	0.56
1:E:39:HIS:HD2	1:E:204:ARG:NH1	2.05	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:104:LEU:HD21	1:C:196:ILE:HG12	1.88	0.54
1:A:94:ALA:HB1	1:A:258:VAL:HG11	1.90	0.54
1:E:47:THR:HG21	1:E:331:ASP:HB3	1.93	0.50
1:A:47:THR:HG21	1:A:331:ASP:HB3	1.93	0.50
1:B:47:THR:HG21	1:B:331:ASP:HB3	1.93	0.50
1:D:313:PRO:O	1:D:314:VAL:HG23	2.12	0.49
1:C:110:GLN:HG3	1:C:116:GLN:HG2	1.94	0.49
1:B:37:ILE:HG13	1:B:129:THR:CG2	2.43	0.49
1:D:141:ARG:NH2	1:D:295:GLU:OE2	2.46	0.49
1:C:141:ARG:NH2	1:C:295:GLU:OE2	2.45	0.48
1:F:102:ASN:OD1	1:F:123:ARG:NH1	2.42	0.48
1:B:102:ASN:OD1	1:B:123:ARG:NH1	2.42	0.48
1:C:11:TYR:HA	1:C:341:LEU:HD13	1.95	0.48
1:A:37:ILE:HG13	1:A:129:THR:CG2	2.44	0.48
1:E:102:ASN:OD1	1:E:123:ARG:NH1	2.41	0.48
1:E:110:GLN:HG3	1:E:116:GLN:HG2	1.95	0.48
1:F:141:ARG:NH2	1:F:295:GLU:OE2	2.47	0.47
1:B:254:SER:OG	1:B:255:HIS:N	2.47	0.47
1:C:102:ASN:OD1	1:C:123:ARG:NH1	2.42	0.47
1:C:242:GLU:HG2	1:E:336:PRO:CG	2.44	0.47
1:A:102:ASN:OD1	1:A:123:ARG:NH1	2.42	0.47
1:C:13:SER:OG	1:C:344:ASP:HB3	2.15	0.47
1:A:65:HIS:ND1	1:A:69:ILE:HD12	2.30	0.47
1:D:324:LEU:HD12	1:D:324:LEU:N	2.30	0.47
1:F:11:TYR:HA	1:F:341:LEU:HD13	1.97	0.46
1:D:11:TYR:HA	1:D:341:LEU:HD13	1.98	0.46
1:D:123:ARG:NH1	1:D:188:TYR:HB3	2.31	0.46
1:B:11:TYR:HA	1:B:341:LEU:HD13	1.98	0.45
1:E:11:TYR:HA	1:E:341:LEU:HD13	1.97	0.45
1:A:11:TYR:HA	1:A:341:LEU:HD13	1.99	0.45
1:E:40:PHE:O	1:E:204:ARG:NH1	2.48	0.45
1:A:276:VAL:HG12	1:A:285:ASP:HA	1.97	0.45
1:E:65:HIS:ND1	1:E:69:ILE:HD12	2.31	0.45
1:B:141:ARG:NH2	1:B:295:GLU:OE1	2.51	0.44
1:A:141:ARG:NH2	1:A:295:GLU:OE1	2.51	0.43
1:A:69:ILE:HG12	1:A:93:VAL:CG1	2.48	0.43
1:C:137:ASN:OD1	1:C:141:ARG:NH1	2.46	0.42
1:D:83:LEU:HD11	1:D:92:LEU:HB2	2.00	0.42
1:A:83:LEU:HD11	1:A:92:LEU:HB2	2.00	0.42
1:C:345:TYR:O	1:C:349:VAL:HG13	2.19	0.42
1:E:83:LEU:HD11	1:E:92:LEU:HB2	2.02	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:336:PRO:HD2	1:C:339:ILE:HD11	2.01	0.42
1:C:83:LEU:HD11	1:C:92:LEU:HB2	2.01	0.41
1:B:137:ASN:OD1	1:B:141:ARG:NH1	2.46	0.40
1:A:245:ARG:NH1	1:A:307:LEU:O	2.54	0.40
1:A:324:LEU:N	1:A:324:LEU:HD12	2.36	0.40
1:E:130:LEU:HD21	1:E:198:VAL:HG21	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	289/371 (78%)	271 (94%)	18 (6%)	0	100	100
1	B	278/371 (75%)	261 (94%)	17 (6%)	0	100	100
1	C	267/371 (72%)	254 (95%)	13 (5%)	0	100	100
1	D	285/371 (77%)	266 (93%)	18 (6%)	1 (0%)	39	80
1	E	275/371 (74%)	259 (94%)	14 (5%)	2 (1%)	26	72
1	F	262/371 (71%)	244 (93%)	17 (6%)	1 (0%)	39	80
All	All	1656/2226 (74%)	1555 (94%)	97 (6%)	4 (0%)	52	88

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	187	GLN
1	E	152	PHE
1	F	180	PRO
1	E	178	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	180/337 (53%)	175 (97%)	5 (3%)	51	84
1	B	142/337 (42%)	138 (97%)	4 (3%)	51	84
1	C	148/337 (44%)	144 (97%)	4 (3%)	52	85
1	D	143/337 (42%)	135 (94%)	8 (6%)	26	68
1	E	160/337 (48%)	153 (96%)	7 (4%)	35	74
1	F	136/337 (40%)	132 (97%)	4 (3%)	50	83
All	All	909/2022 (45%)	877 (96%)	32 (4%)	43	80

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	TYR
1	A	74	GLN
1	A	193	GLU
1	A	237	HIS
1	A	258	VAL
1	B	11	TYR
1	B	16	ARG
1	B	74	GLN
1	B	237	HIS
1	C	74	GLN
1	C	104	LEU
1	C	237	HIS
1	C	349	VAL
1	D	10	LEU
1	D	11	TYR
1	D	74	GLN
1	D	143	GLN
1	D	193	GLU
1	D	237	HIS
1	D	255	HIS
1	D	258	VAL
1	E	11	TYR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	E	16	ARG
1	E	74	GLN
1	E	104	LEU
1	E	143	GLN
1	E	193	GLU
1	E	237	HIS
1	F	16	ARG
1	F	62	GLU
1	F	74	GLN
1	F	237	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	277	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

6 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	YCM	A	103	1	7,9,10	1.43	2 (28%)	5,10,12	1.85	2 (40%)
1	YCM	B	103	1	7,9,10	1.38	2 (28%)	5,10,12	1.65	2 (40%)
1	YCM	C	103	1	7,9,10	1.40	2 (28%)	5,10,12	1.67	2 (40%)
1	YCM	D	103	1	7,9,10	1.55	2 (28%)	5,10,12	1.85	2 (40%)
1	YCM	E	103	1	7,9,10	1.39	2 (28%)	5,10,12	1.59	1 (20%)
1	YCM	F	103	1	7,9,10	1.44	2 (28%)	5,10,12	1.83	2 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	YCM	A	103	1	-	0/6/8/10	0/0/0/0
1	YCM	B	103	1	-	0/6/8/10	0/0/0/0
1	YCM	C	103	1	-	0/6/8/10	0/0/0/0
1	YCM	D	103	1	-	0/6/8/10	0/0/0/0
1	YCM	E	103	1	-	0/6/8/10	0/0/0/0
1	YCM	F	103	1	-	0/6/8/10	0/0/0/0

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	103	YCM	CB-SG	-2.91	1.75	1.81
1	A	103	YCM	CD-SG	-2.70	1.75	1.81
1	D	103	YCM	CD-SG	-2.67	1.75	1.81
1	E	103	YCM	CD-SG	-2.66	1.75	1.81
1	F	103	YCM	CD-SG	-2.57	1.76	1.81
1	F	103	YCM	CB-SG	-2.53	1.76	1.81
1	B	103	YCM	CB-SG	-2.53	1.76	1.81
1	A	103	YCM	CB-SG	-2.50	1.76	1.81
1	C	103	YCM	CD-SG	-2.49	1.76	1.81
1	C	103	YCM	CB-SG	-2.46	1.76	1.81
1	B	103	YCM	CD-SG	-2.31	1.76	1.81
1	E	103	YCM	CB-SG	-2.20	1.77	1.81

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	103	YCM	O-C-CA	-2.50	119.03	125.72
1	C	103	YCM	O-C-CA	-2.46	119.11	125.72
1	A	103	YCM	O-C-CA	-2.38	119.33	125.72
1	F	103	YCM	O-C-CA	-2.36	119.41	125.72
1	B	103	YCM	O-C-CA	-2.23	119.73	125.72
1	C	103	YCM	CD-CE-NZ2	2.32	118.04	115.48
1	E	103	YCM	CD-CE-NZ2	2.48	118.22	115.48
1	B	103	YCM	CD-CE-NZ2	2.52	118.25	115.48
1	D	103	YCM	CD-CE-NZ2	2.82	118.59	115.48
1	A	103	YCM	CD-CE-NZ2	2.87	118.64	115.48
1	F	103	YCM	CD-CE-NZ2	2.93	118.71	115.48

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	301/371 (81%)	-0.31	2 (0%) 89 83	24, 49, 73, 102	0
1	B	294/371 (79%)	-0.32	0 100 100	31, 58, 91, 102	0
1	C	283/371 (76%)	-0.30	0 100 100	29, 58, 88, 115	0
1	D	297/371 (80%)	-0.24	3 (1%) 84 75	31, 63, 95, 117	0
1	E	291/371 (78%)	-0.35	0 100 100	27, 55, 84, 93	0
1	F	280/371 (75%)	-0.22	5 (1%) 71 58	34, 61, 101, 129	0
All	All	1746/2226 (78%)	-0.29	10 (0%) 90 84	24, 57, 90, 129	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	133	THR	4.6
1	D	101	GLY	3.1
1	D	249	VAL	3.0
1	F	134	ASP	2.8
1	D	248	ILE	2.7
1	F	146	SER	2.7
1	A	313	PRO	2.6
1	F	117	ASP	2.5
1	A	252	TYR	2.2
1	F	132	GLU	2.2

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy

less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
1	YCM	B	103	10/11	0.90	0.36	-	57,71,92,94	0
1	YCM	D	103	10/11	0.86	0.29	-	70,76,95,97	0
1	YCM	E	103	10/11	0.90	0.20	-	50,53,67,75	0
1	YCM	A	103	10/11	0.91	0.26	-	54,63,111,126	0
1	YCM	C	103	10/11	0.85	0.20	-	64,75,78,80	0
1	YCM	F	103	10/11	0.90	0.24	-	43,60,102,109	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.