



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:33 PM GMT

PDB ID : 3ZUT
Title : THE STRUCTURE OF OST1 (D160A) KINASE
Authors : Yunta, C.; Martinez-Ripoll, M.; Albert, A.
Deposited on : 2011-07-20
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3553 (2.50-2.50)
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	362	<div> <div>2%</div> <div>65%</div> <div>10%</div> <div>••</div> <div>22%</div> </div>
1	B	362	<div> <div>%</div> <div>58%</div> <div>16%</div> <div>•</div> <div>22%</div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4582 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SERINE/THREONINE-PROTEIN KINASE SRK2E.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	281	Total	C	N	O	S	0	0	0
			2258	1444	393	409	12			
1	B	282	Total	C	N	O	S	0	0	0
			2265	1449	394	410	12			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	160	ALA	ASP	ENGINEERED MUTATION	UNP Q940H6
B	160	ALA	ASP	ENGINEERED MUTATION	UNP Q940H6

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	36	Total	O	0	0
			36	36		
2	B	23	Total	O	0	0
			23	23		

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Chain A:
-
- 2% 65% 10% 22%
- Y163
L163
L164
L165
L166
L167
L168
L169
L170
L171
L172
L173
L174
L175
L176
L177
L178
L179
L180
L181
L182
L183
L184
L185
L186
L187
L188
L189
L190
L191
L192
L193
L194
L195
L196
L197
L198
L199
L200
L201
L202
L203
L204
L205
L206
L207
L208
L209
L210
L211
L212
L213
L214
L215
L216
L217
L218
L219
L220
L221
L222
L223
L224
L225
L226
L227
L228
L229
L230
L231
L232
L233
L234
L235
L236
L237
L238
L239
L240
L241
L242
L243
L244
L245
L246
L247
L248
L249
L250
L251
L252
L253
L254
L255
L256
L257
L258
L259
L260
L261
L262
L263
L264
L265
L266
L267
L268
L269
L270
L271
L272
L273
L274
L275
L276
L277
L278
L279
L280
L281
L282
L283
L284
L285
L286
L287
L288
L289
L290
L291
L292
L293
L294
L295
L296
L297
L298
L299
L300
L301
L302
L303
L304
L305
L306
L307
L308
L309
L310
L311
L312
L313
L314
L315
L316
L317
L318
L319
L320
L321
L322
L323
L324
L325
L326
L327
L328
L329
L330
L331
L332
L333
L334
L335
L336
L337
L338
L339
L340
L341
L342
L343
L344
L345
L346
L347
L348
L349
L350
L351
L352
L353
L354
L355
L356
L357
L358
L359
L360
L361
L362
L363
L364
L365
L366
L367
L368
L369
L370
L371
L372
L373
L374
L375
L376
L377
L378
L379
L380
L381
L382
L383
L384
L385
L386
L387
L388
L389
L390
L391
L392
L393
L394
L395
L396
L397
L398
L399
L400
L401
L402
L403
L404
L405
L406
L407
L408
L409
L410
L411
L412
L413
L414
L415
L416
L417
L418
L419
L420
L421
L422
L423
L424
L425
L426
L427
L428
L429
L430
L431
L432
L433
L434
L435
L436
L437
L438
L439
L440
L441
L442
L443
L444
L445
L446
L447
L448
L449
L450
L451
L452
L453
L454
L455
L456
L457
L458
L459
L460
L461
L462
L463
L464
L465
L466
L467
L468
L469
L470
L471
L472
L473
L474
L475
L476
L477
L478
L479
L480
L481
L482
L483
L484
L485
L486
L487
L488
L489
L490
L491
L492
L493
L494
L495
L496
L497
L498
L499
L500
L501
L502
L503
L504
L505
L506
L507
L508
L509
L510
L511
L512
L513
L514
L515
L516
L517
L518
L519
L520
L521
L522
L523
L524
L525
L526
L527
L528
L529
L530
L531
L532
L533
L534
L535
L536
L537
L538
L539
L540
L541
L542
L543
L544
L545
L546
L547
L548
L549
L550
L551
L552
L553
L554
L555
L556
L557
L558
L559
L560
L561
L562
L563
L564
L565
L566
L567
L568
L569
L570
L571
L572
L573
L574
L575
L576
L577
L578
L579
L580
L581
L582
L583
L584
L585
L586
L587
L588
L589
L590
L591
L592
L593
L594
L595
L596
L597
L598
L599
L600
L601
L602
L603
L604
L605
L606
L607
L608
L609
L610
L611
L612
L613
L614
L615
L616
L617
L618
L619
L620
L621
L622
L623
L624
L625
L626
L627
L628
L629
L630
L631
L632
L633
L634
L635
L636
L637
L638
L639
L640
L641
L642
L643
L644
L645
L646
L647
L648
L649
L650
L651
L652
L653
L654
L655
L656
L657
L658
L659
L660
L661
L662
L663
L664
L665
L666
L667
L668
L669
L670
L671
L672
L673
L674
L675
L676
L677
L678
L679
L680
L681
L682
L683
L684
L685
L686
L687
L688
L689
L690
L691
L692
L693
L694
L695
L696
L697
L698
L699
L700
L701
L702
L703
L704
L705
L706
L707
L708
L709
L710
L711
L712
L713
L714
L715
L716
L717
L718
L719
L720
L721
L722
L723
L724
L725
L726
L727
L728
L729
L730
L731
L732
L733
L734
L735
L736
L737
L738
L739
L740
L741
L742
L743
L744
L745
L746
L747
L748
L749
L750
L751
L752
L753
L754
L755
L756
L757
L758
L759
L760
L761
L762
L763
L764
L765
L766
L767
L768
L769
L770
L771
L772
L773
L774
L775
L776
L777
L778
L779
L780
L781
L782
L783
L784
L785
L786
L787
L788
L789
L790
L791
L792
L793
L794
L

- [illegible]

4 Data and refinement statistics

Property	Value	Source
Space group	P 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	77.27Å 99.19Å 107.83Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.74 – 2.50 41.74 – 2.24	Depositor EDS
% Data completeness (in resolution range)	99.5 (41.74-2.50) 98.3 (41.74-2.24)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.18 (at 2.24Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.198 , 0.251 0.186 , 0.244	Depositor DCC
R_{free} test set	1470 reflections (5.03%)	DCC
Wilson B-factor (Å ²)	57.4	Xtriage
Anisotropy	0.080	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 35.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 40047 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4582	wwPDB-VP
Average B, all atoms (Å ²)	64.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.57	2/2310 (0.1%)	0.55	0/3125
1	B	0.49	2/2317 (0.1%)	0.55	0/3135
All	All	0.53	4/4627 (0.1%)	0.55	0/6260

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	159	CYS	CB-SG	13.56	2.05	1.82
1	A	137	CYS	CB-SG	10.21	1.99	1.82
1	B	159	CYS	CB-SG	9.91	1.99	1.82
1	B	137	CYS	CB-SG	5.31	1.91	1.82

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	137	CYS	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2258	0	2253	36	0
1	B	2265	0	2268	38	0
2	A	36	0	0	3	0
2	B	23	0	0	3	0
All	All	4582	0	4521	74	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (74) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:159:CYS:SG	1:A:159:CYS:CB	2.05	1.44
1:B:58:ILE:HB	1:B:316:THR:HG23	1.61	0.82
1:A:191:LYS:HE3	1:A:191:LYS:H	1.47	0.78
1:B:31:ASN:ND2	1:B:145:ASN:OD1	2.19	0.75
1:B:59:ASP:OD2	1:B:61:ASN:ND2	2.21	0.74
1:B:116:GLU:OE1	2:B:2008:HOH:O	2.08	0.72
1:B:49:VAL:HG22	1:B:93:VAL:HG22	1.73	0.70
1:A:84:ILE:HB	1:A:91:ALA:HB3	1.74	0.69
1:B:22:GLU:OE1	1:B:39:ARG:NH2	2.25	0.68
1:A:85:LEU:O	2:A:2005:HOH:O	2.12	0.68
1:B:72:LEU:O	2:B:2003:HOH:O	2.12	0.67
1:A:191:LYS:N	1:A:191:LYS:HE3	2.09	0.67
1:A:139:ARG:HG3	1:A:139:ARG:HH11	1.59	0.66
1:A:139:ARG:NH1	2:A:2017:HOH:O	2.29	0.66
1:B:305:ILE:O	1:B:309:MET:HG2	1.96	0.65
1:A:139:ARG:HG3	1:A:139:ARG:NH1	2.12	0.62
1:B:51:TYR:HA	1:B:90:LEU:O	2.01	0.60
1:A:190:LYS:HA	1:A:191:LYS:HE3	1.85	0.59
1:A:191:LYS:CE	1:A:191:LYS:H	2.15	0.59
1:A:39:ARG:HE	1:A:44:ASN:HA	1.68	0.58
1:A:12:LEU:HD12	1:A:12:LEU:H	1.69	0.57
1:A:217:PHE:O	1:A:232:ARG:HD2	2.07	0.55
1:B:181:ALA:HA	1:B:233:ILE:HD11	1.89	0.54
1:A:180:PRO:HG2	1:A:229:THR:HG21	1.89	0.54
1:B:63:LYS:O	1:B:67:ILE:HG12	2.08	0.54
1:A:313:ALA:O	1:A:316:THR:OG1	2.26	0.53
1:A:63:LYS:O	1:A:67:ILE:HG12	2.09	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:45:GLU:OE1	1:B:81:LYS:NZ	2.42	0.51
1:B:217:PHE:O	1:B:232:ARG:HD2	2.11	0.51
1:A:56:GLU:H	1:A:56:GLU:CD	2.14	0.51
1:B:317:VAL:HG23	1:B:319:PRO:HD3	1.93	0.50
1:A:54:ARG:HD3	1:A:315:ALA:HA	1.92	0.50
1:B:284:ASP:OD1	1:B:284:ASP:N	2.43	0.49
1:A:45:GLU:OE1	1:A:81:LYS:HE3	2.12	0.49
1:A:284:ASP:N	1:A:284:ASP:OD1	2.42	0.49
1:B:191:LYS:HE2	1:B:192:GLU:H	1.78	0.48
1:B:101:GLU:OE1	1:B:145:ASN:ND2	2.41	0.48
1:A:191:LYS:CD	1:A:191:LYS:H	2.27	0.47
1:B:307:GLU:O	1:B:311:ILE:HG13	2.14	0.47
1:A:101:GLU:OE1	1:A:145:ASN:ND2	2.39	0.47
1:A:144:GLU:OE1	1:A:144:GLU:N	2.43	0.47
1:B:187:VAL:HG22	1:B:193:TYR:CE2	2.51	0.46
1:A:42:GLN:N	1:A:42:GLN:OE1	2.49	0.46
1:A:58:ILE:HG13	1:A:315:ALA:O	2.16	0.46
1:A:55:GLY:HA2	1:A:316:THR:O	2.16	0.46
1:A:200:VAL:HG12	1:A:257:ILE:HG23	1.98	0.45
1:A:43:SER:O	1:A:44:ASN:CB	2.63	0.45
1:A:139:ARG:CG	1:A:139:ARG:HH11	2.29	0.44
1:B:85:LEU:HB2	1:B:311:ILE:HG23	1.99	0.44
1:A:67:ILE:HD11	1:A:309:MET:SD	2.57	0.44
1:B:236:VAL:HG12	1:B:236:VAL:O	2.17	0.44
1:A:229:THR:OG1	2:A:2026:HOH:O	2.21	0.44
1:B:204:GLY:HA3	1:B:257:ILE:HG21	1.99	0.43
1:B:138:HIS:CG	1:B:139:ARG:N	2.86	0.43
1:A:115:ASP:OD1	1:A:118:ARG:NH2	2.42	0.43
1:B:56:GLU:HG3	1:B:57:LYS:HD2	2.02	0.42
1:B:46:LEU:HA	1:B:46:LEU:HD12	1.82	0.42
1:B:144:GLU:OE1	2:B:2015:HOH:O	2.21	0.42
1:B:59:ASP:N	1:B:59:ASP:OD1	2.52	0.42
1:A:54:ARG:HD2	1:A:317:VAL:O	2.19	0.42
1:B:53:GLU:HA	1:B:89:HIS:CD2	2.55	0.42
1:A:314:GLU:O	1:A:317:VAL:HG22	2.19	0.41
1:B:138:HIS:HA	1:B:199:ASP:OD1	2.20	0.41
1:B:68:ASN:O	1:B:71:SER:OG	2.24	0.41
1:B:85:LEU:CD1	1:B:315:ALA:HB2	2.51	0.41
1:B:200:VAL:HG12	1:B:257:ILE:HG23	2.03	0.41
1:B:115:ASP:OD1	1:B:118:ARG:NH2	2.43	0.41
1:A:236:VAL:HG12	1:A:236:VAL:O	2.20	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:191:LYS:H	1:B:191:LYS:CE	2.34	0.41
1:B:206:THR:O	1:B:209:VAL:HG12	2.20	0.41
1:B:197:VAL:HG12	1:B:265:ARG:NH1	2.35	0.40
1:B:73:ARG:HB3	1:B:73:ARG:HE	1.38	0.40
1:A:118:ARG:HB2	1:A:276:TRP:CZ2	2.57	0.40
1:B:77:ILE:O	1:B:79:ARG:HG3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	275/362 (76%)	262 (95%)	12 (4%)	1 (0%)	39	61
1	B	276/362 (76%)	262 (95%)	11 (4%)	3 (1%)	17	31
All	All	551/724 (76%)	524 (95%)	23 (4%)	4 (1%)	26	46

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	318	PRO
1	B	161	PHE
1	B	53	GLU
1	A	318	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	246/321 (77%)	232 (94%)	14 (6%)	25	46
1	B	248/321 (77%)	226 (91%)	22 (9%)	12	23
All	All	494/642 (77%)	458 (93%)	36 (7%)	17	32

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	LEU
1	A	18	SER
1	A	31	ASN
1	A	39	ARG
1	A	56	GLU
1	A	83	VAL
1	A	137	CYS
1	A	139	ARG
1	A	155	ARG
1	A	159	CYS
1	A	191	LYS
1	A	227	ARG
1	A	284	ASP
1	A	316	THR
1	B	12	LEU
1	B	14	ILE
1	B	19	ASP
1	B	20	ARG
1	B	31	ASN
1	B	34	VAL
1	B	37	LEU
1	B	42	GLN
1	B	56	GLU
1	B	57	LYS
1	B	58	ILE
1	B	59	ASP
1	B	61	ASN
1	B	73	ARG
1	B	83	VAL
1	B	137	CYS
1	B	139	ARG
1	B	155	ARG
1	B	159	CYS
1	B	191	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	284	ASP
1	B	316	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	31	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	281/362 (77%)	-0.16	6 (2%) 67 71	34, 56, 95, 139	0
1	B	282/362 (77%)	-0.07	5 (1%) 71 75	37, 61, 113, 145	0
All	All	563/724 (77%)	-0.11	11 (1%) 68 72	34, 59, 108, 145	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	10	MET	4.8
1	A	11	ASP	3.9
1	A	159	CYS	3.3
1	A	34	VAL	2.9
1	B	13	PRO	2.8
1	A	137	CYS	2.7
1	B	90	LEU	2.4
1	B	301	PRO	2.4
1	B	117	ALA	2.4
1	A	319	PRO	2.1
1	B	176	THR	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.