



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 02:33 PM GMT

PDB ID : 3ZUT
Title : THE STRUCTURE OF OST1 (D160A) KINASE
Authors : Yunta, C.; Martinez-Ripoll, M.; Albert, A.
Deposited on : 2011-07-20
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3553 (2.50-2.50)
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	362	<div> <div>2%</div> <div>65%</div> <div>10%</div> <div>••</div> <div>22%</div> </div>
1	B	362	<div> <div>%</div> <div>58%</div> <div>16%</div> <div>•</div> <div>22%</div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4582 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SERINE/THREONINE-PROTEIN KINASE SRK2E.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	281	Total	C	N	O	S	0	0	0
			2258	1444	393	409	12			
1	B	282	Total	C	N	O	S	0	0	0
			2265	1449	394	410	12			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	160	ALA	ASP	ENGINEERED MUTATION	UNP Q940H6
B	160	ALA	ASP	ENGINEERED MUTATION	UNP Q940H6

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	36	Total	O	0	0
			36	36		
2	B	23	Total	O	0	0
			23	23		

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Chain A:

Sequence logo for Chain A. The top row shows conservation percentages (0% to 22%). The middle row shows amino acid conservation percentages. The bottom row shows the amino acid sequence.

Position	Amino Acid	Conservation (%)
1	Met	0.0
2	Asp	0.0
3	Arg	0.0
4	Pro	0.0
5	Ala	0.0
6	Val	0.0
7	Gly	0.0
8	Pro	0.0
9	His	0.0
10	Ser	0.0
11	Gln	0.0
12	Pro	0.0
13	Lys	0.0
14	Ser	0.0
15	Thr	0.0
16	Val	0.0
17	Gly	0.0
18	His	0.0
19	Ser	0.0
20	Gln	0.0
21	Pro	0.0
22	Lys	0.0
23	Ser	0.0
24	Thr	0.0
25	Val	0.0
26	Gln	0.0
27	Thr	0.0
28	Gly	0.0
29	Thr	0.0
30	Ala	0.0
31	Met	0.0
32	Met	0.0
33	Ala	0.0
34	Thr	0.0
35	Val	0.0
36	Ile	0.0
37	Asp	0.0
38	Ser	0.0
39	Ser	0.0
40	Asn	0.0
41	Met	0.0
42	Leu	0.0
43	Leu	0.0
44	Asp	0.0
45	Asp	0.0
46	Met	0.0
47	Leu	0.0
48	Asp	0.0
49	Leu	0.0
50	Asp	0.0
51	Leu	0.0
52	Asp	0.0
53	Leu	0.0
54	Asp	0.0
55	Leu	0.0
56	Asp	0.0
57	Leu	0.0
58	Asp	0.0
59	Leu	0.0
60	Asp	0.0
61	Leu	0.0
62	Asp	0.0
63	Leu	0.0
64	Asp	0.0
65	Leu	0.0
66	Asp	0.0
67	Leu	0.0
68	Asp	0.0
69	Leu	0.0
70	Asp	0.0
71	Leu	0.0
72	Asp	0.0
73	Leu	0.0
74	Asp	0.0
75	Leu	0.0
76	Asp	0.0
77	Leu	0.0
78	Asp	0.0
79	Leu	0.0
80	Asp	0.0
81	Leu	0.0
82	Asp	0.0
83	Leu	0.0
84	Asp	0.0
85	Leu	0.0
86	Asp	0.0
87	Leu	0.0
88	Asp	0.0
89	Leu	0.0
90	Asp	0.0
91	Leu	0.0
92	Asp	0.0
93	Leu	0.0
94	Asp	0.0
95	Leu	0.0
96	Asp	0.0
97	Leu	0.0
98	Asp	0.0
99	Leu	0.0
100	Asp	0.0
101	Leu	0.0
102	Asp	0.0
103	Leu	0.0
104	Asp	0.0
105	Leu	0.0
106	Asp	0.0
107	Leu	0.0
108	Asp	0.0
109	Leu	0.0
110	Asp	0.0
111	Leu	0.0
112	Asp	0.0
113	Leu	0.0
114	Asp	0.0
115	Leu	0.0
116	Asp	0.0
117	Leu	0.0
118	Asp	0.0
119	Leu	0.0
120	Asp	0.0
121	Leu	0.0
122	Asp	0.0
123	Leu	0.0
124	Asp	0.0
125	Leu	0.0
126	Asp	0.0
127	Leu	0.0
128	Asp	0.0
129	Leu	0.0
130	Asp	0.0
131	Leu	0.0
132	Asp	0.0
133	Leu	0.0
134	Asp	0.0
135	Leu	0.0
136	Asp	0.0
137	Leu	0.0
138	Asp	0.0
139	Leu	0.0
140	Asp	0.0
141	Leu	0.0
142	Asp	0.0
143	Leu	0.0
144	Asp	0.0
145	Leu	0.0
146	Asp	0.0
147	Leu	0.0
148	Asp	0.0
149	Leu	0.0
150	Asp	0.0
151	Leu	0.0
152		

Chain B:

58% 16% 22%

4 Data and refinement statistics

Property	Value	Source
Space group	P 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	77.27Å 99.19Å 107.83Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.74 – 2.50 41.74 – 2.24	Depositor EDS
% Data completeness (in resolution range)	99.5 (41.74-2.50) 98.3 (41.74-2.24)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.18 (at 2.24Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.198 , 0.251 0.186 , 0.244	Depositor DCC
R_{free} test set	1470 reflections (5.03%)	DCC
Wilson B-factor (Å ²)	57.4	Xtriage
Anisotropy	0.080	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 35.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 40047 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4582	wwPDB-VP
Average B, all atoms (Å ²)	64.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.57	2/2310 (0.1%)	0.55	0/3125
1	B	0.49	2/2317 (0.1%)	0.55	0/3135
All	All	0.53	4/4627 (0.1%)	0.55	0/6260

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	159	CYS	CB-SG	13.56	2.05	1.82
1	A	137	CYS	CB-SG	10.21	1.99	1.82
1	B	159	CYS	CB-SG	9.91	1.99	1.82
1	B	137	CYS	CB-SG	5.31	1.91	1.82

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	137	CYS	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2258	0	2253	36	0
1	B	2265	0	2268	38	0
2	A	36	0	0	3	0
2	B	23	0	0	3	0
All	All	4582	0	4521	74	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 74 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:159:CYS:SG	1:A:159:CYS:CB	2.05	1.44
1:B:58:ILE:HB	1:B:316:THR:HG23	1.61	0.82
1:A:191:LYS:HE3	1:A:191:LYS:H	1.47	0.78
1:B:31:ASN:ND2	1:B:145:ASN:OD1	2.19	0.75
1:B:59:ASP:OD2	1:B:61:ASN:ND2	2.21	0.74

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	275/362 (76%)	262 (95%)	12 (4%)	1 (0%)	39	61
1	B	276/362 (76%)	262 (95%)	11 (4%)	3 (1%)	17	31
All	All	551/724 (76%)	524 (95%)	23 (4%)	4 (1%)	26	46

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	318	PRO
1	B	161	PHE
1	B	53	GLU
1	A	318	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	246/321 (77%)	232 (94%)	14 (6%)	25	46
1	B	248/321 (77%)	226 (91%)	22 (9%)	12	23
All	All	494/642 (77%)	458 (93%)	36 (7%)	17	32

5 of 36 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	19	ASP
1	B	37	LEU
1	B	191	LYS
1	B	31	ASN
1	B	42	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	31	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	281/362 (77%)	-0.16	6 (2%) 67 71	34, 56, 95, 139	0
1	B	282/362 (77%)	-0.07	5 (1%) 71 75	37, 61, 113, 145	0
All	All	563/724 (77%)	-0.11	11 (1%) 68 72	34, 59, 108, 145	0

The worst 5 of 11 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	10	MET	4.8
1	A	11	ASP	3.9
1	A	159	CYS	3.3
1	A	34	VAL	2.9
1	B	13	PRO	2.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.